

MOLECULAR ROTATION IN SUPERFLUID ^4He NANODROPLETS: THE VALIDITY OF A HYDRODYNAMIC MODEL

CARLO CALLEGARI, ANDRÉ CONJESTEAU, IRENE REINHARD, KEVIN K. LEHMANN, and GIACINTO SCOLES, *Department of Chemistry, Princeton University, Princeton NJ 08544*; FRANCO DALFOVO, *Dipartimento di Matematica e Fisica, Università Cattolica del Sacro Cuore, I-25121 Brescia (Italy)*.

In order to accurately predict the effective moments of inertia (I_{eff}) of linear molecules rotating in superfluid ^4He droplets, we have recently devised a superfluid hydrodynamic model. In the present implementation, the helium density profile induced by the He-molecule interaction potential is first calculated at the Density Functional level, and then used as the input of the hydrodynamic equation for the irrotational motion of a viscousless fluid. The kinetic energy of the fluid (E_k) is then used to calculate I_{eff} via: $E_k = \frac{1}{2} \Delta I \omega^2$, where ω is the angular velocity of the molecule, and ΔI is the difference between I_{eff} and the moment of inertia of the bare molecule.^a The model relies on the assumption that the density of the fluid in the rotating frame of reference is independent of ω and can therefore be calculated in the limit of a static molecule (adiabatic following approximation). The validity of this approximation, and its first-order corrections will be discussed. The advantages and limitations of Density Functional Theory for calculating the true helium density will also be addressed, in a comparison with Quantum Monte Carlo results which recently became available for HCN and its oligomers.^b

^aC. Callegari, A. Conjusteau, I. Reinhard, K.K. Lehmann, G. Scoles, and F. Dalfovo, Phys. Rev. Lett. **83**, 5058 (1999); *ibid.* **84**, 1848(E) (2000).

^bE. Draeger and D. M. Ceperley, private communication.