

AGGREGATED MOLECULAR FORMATIONS IN POLAR LIQUIDS AND ITS DISPLAY ON RAMAN SPECTRA

U. N. TASHKENBAEV, F. H. TUKHVATULLIN, A. JUMABAEV, Z. MAMATOV and H. HUSHVAKTOV,
Samarkand State University, 703004 Samarkand, Univ. Blvd, 15, Uzbekistan (email: info@samarkand.uz).

On a base of Raman spectra study of C=O vibrations band for some ketones (cyclohexanone, methylethylketone, acetone) the possibility of existence of dipole-dipole aggregated formations is analyzed. For these substances, the noncoincidence of the C=O vibrations band maxima frequencies for parallel ($I_{\parallel}(\nu)$) and perpendicular ($I_{\perp}(\nu)$) polarized components is typical. The difference $\Delta\nu$ is 5-6 cm^{-1} . Our studies show that, in a large degree, the presence of $\Delta\nu$ for C=O vibration and its change with temperature and with the dilution of substance is a result of imposition of near-placed bands with different polarization ratio belonging to monomers and dipole-dipole aggregated formations. In case of pure methylethylketone and acetone, the C=O vibrations band complexity is displayed in the form of different asymmetry of resulting bands in $I_{\parallel}(\nu)$ and $I_{\perp}(\nu)$, in the peculiarities in trend of depolarization ratio within the band and in the form of typical change of a band's shape at dilution in non-polar and polar solvents. For cyclohexanone the band complexity can evidently be seen from the shape of experimental spectra for $I_{\parallel}(\nu)$ and $I_{\perp}(\nu)$. The behavior of cyclohexanone complicated C=O vibrations band at dilution in non-polar and polar solvents is analogous to that for acetone or methylethylketone.