

## ELECTRONIC SPECTROSCOPY OF WC FROM 17,500 to 24,000 $\text{cm}^{-1}$

A. W. SMITH, S. M. SICKAFOOSE, and M. D. MORSE, *Department of Chemistry, University of Utah, Salt Lake City, UT, 84112.*

Resonant two-photon ionization spectroscopy has been used to study the diatomic molecule WC. A low resolution scan revealed a five member vibrational progression beginning at 17 585  $\text{cm}^{-1}$  with the  $\nu_{00}$  transition. Analysis of this progression yielded a vibrational frequency of  $\omega'_e(^{184}W^{12}C) = 752.6(4.9) \text{ cm}^{-1}$  and a bond length of  $r'_e(^{184}W^{12}C) = 1.747(4) \text{ \AA}$ . In addition, several unassigned bands were rotationally resolved. Interestingly, all of the observed excited states have  $\Omega' = 2$ . All of the rotationally resolved transitions were fit simultaneously to produce the best possible fit of the ground state. Assignment of these bands confirmed a ground state of  $^3\Delta_1$  from a  $14\sigma^2 8\pi^4 15\sigma^2 4\delta^1 16\sigma^1$  configuration and determined the ground state as  $r''_0(^{184}W^{12}C) = 1.7143(2) \text{ \AA}$ . Dispersed fluorescence studies to elucidate the ground and low-lying excited states will also be reported. These results on WC are compared to the results of studies on MoC and other transition metal carbides.