

## FREQUENCY MODULATION SPECTROSCOPY OF THE TiO E-X TRANSITION

KAORI KOBAYASHI, GREGORY E. HALL, and TREVOR J. SEARS, *Department of Chemistry, Brookhaven National Laboratory, Upton, New York 11973-5000*; and ANTHONY J. MERER, *Department of Chemistry, University of British Columbia, 2036 Main Mall, Vancouver, British Columbia V6T 1Z1 Canada*.

Frequency modulation spectroscopy has been successfully used to measure the spectra of jet-cooled transition-metal containing radicals. This paper reports the recent results on the  $E^3\Pi-X^3\Delta$  (0,0) and (1,0) bands of TiO. This band system is known to be 10 times weaker than the B-X or A-X bands and illustrates the high sensitivity of our spectrometer. TiO radicals were generated by the reaction of laser-ablated metal atoms with O<sub>2</sub> or CO<sub>2</sub> seeded in a supersonic expansion of Ar. All 3 sub-bands of the (0,0) band were recorded at high resolution and previously unresolved lambda-type doubling of  $\Omega=1$  of the upper state was also resolved. More precise and some previously undetermined molecular constants are now available. The still weaker (1,0) band was also observed and an equilibrium bond length in the  $E^3\Pi$  was determined.

*Acknowledgments:* This work at Brookhaven National Laboratory was carried out under Contract No. DE-AC02-98CH10886 with the U.S. Department of Energy and supported by its Division of Chemical Sciences, Office of Basic Energy Sciences.