

INFRARED PHOTODISSOCIATION SPECTROSCOPY OF $M^+-(C_2H_2)_n$ CLUSTERS^a

T. JAEGER, R. S. WALTERS, and M. A. DUNCAN, *University of Georgia, Department of Chemistry, Athens, GA, 30602-2556.*

Metal ion-acetylene complexes of the general form $M^+-(C_2H_2)_n$ are synthesized in the gas phase by a pulsed nozzle laser ablation cluster source. These ions are then mass selected in a reflectron time-of-flight mass spectrometer where they are excited and undergo multiphoton absorption from a tunable infrared OPO/OPA laser system. Efficient fragmentation occurs by the loss of whole C_2H_2 units and this process is enhanced on vibrational resonances near the symmetric and asymmetric C-H stretch frequencies of acetylene. Infrared photofragmentation spectra are obtained by monitoring the fragment yield versus energy of the tunable infrared laser in the 3000 cm^{-1} to 3400 cm^{-1} region. The observed infrared band positions are red-shifted from those of free acetylene consistent with the weakening of the carbon-carbon bond energy by π electron withdrawal. Interpretation of the spectra for $M^+ = Ni^+$ and Co^+ will be discussed.

^aResearch sponsored by the U.S. Department of Energy