

WAVE NUMBERS, STRENGTHS, WIDTHS AND SHIFTS WITH PRESSURE OF LINES OF  $^{16}\text{O}_2$  IN SYSTEMS  $a$   
 $^1\Delta_g - X^3\Sigma_g^-$  AND  $b$   $^1\Sigma_g^+ - X^3\Sigma_g^-$

SWEE-LAN CHEAH, YUAN-PERN LEE, *Department of Chemistry, National Tsing Hua University, Hsinchu 30013, Taiwan*; J. F. OGILVIE, *Centre for Experimental and Constructive Mathematics, Department of Mathematics, Simon Fraser University, 8888 University drive, Burnaby, British Columbia V5A 1S6 Canada.*

Analysis of newly measured spectral data<sup>a</sup> for wave number and intensities of transitions of  $^{16}\text{O}_2$  in absorption in systems  $a$   $^1\Delta_g, v=0 - X^3\Sigma_g^-, v=0$  and  $b$   $^1\Sigma_g^+, v=0, 1, 2 - X^3\Sigma_g^-, v=0$  with proper statistical treatment yields band parameters for excited states, based on parameters for the ground vibrational state valid up to  $J=41$ , that are the most precise ( $\sigma = 0.021, 0.038, 0.043, 0.50 \text{ m}^{-1}$  respectively) in published form, and strengths of lines that are internally consistent and in satisfactory agreement with other published values.

---

<sup>a</sup>S.-L. Cheah, Y.-P. Lee and J. F. Ogilvie, *J. Quant. Spectrosc. Radiative Transfer*, **64**, 467–482 (2000)