

AUTOIONIZING STATES OF CALCIUM MONOFLUORIDE

XING JIANG, JASON O. CLEVINGER, SERGEY I. PANOV AND ROBERT W. FIELD, *Department of Chemistry, Massachusetts Institute of Technology, Cambridge, MA 02139.*

We will report experimental results from the pulsed optical-optical double resonance study of Calcium Monofluoride(CaF). These spectra are recorded via D $^2\Sigma^+$ state of CaF in the autoionization region above $v^+=0$ ionization potential with $n^*=15 \sim 45$. Compared to the OODR spectra of CaCl obtained previously in the same region, more $^2\Sigma^+$ and $^2\Pi$ states are assigned for CaF. This can be explained by the predissociation of CaCl. In this autoionization region the disappearance or shift of some members of the predicted CaF core-penetrating series reveals the existence of perturbation between certain states, which is not the case for CaCl. The OODR spectra of CaF in the high- n^* region indicate that the vibrationally autoionizing states of CaF converge to $v^+ > 0$ vibrational levels of CaF^+ ion. This work completes previous studies of CaF by exploiting different intermediate states, e.g. the reverse-polarized C $^2\Pi$ state, to investigate the vibrational constant ω_e for each Rydberg series and the corresponding quantum defect derivative with respect to the internuclear distance R^+ , which controls the autoionization process.