

CO/He GLOW DISCHARGE TEMPERATURE MEASUREMENTS USING CO 4th POSITIVE BAND LIF

R. J. LEIWEKE and W. R. LEMPert, *Nonequilibrium Thermodynamics Laboratory, Department of Mechanical Engineering and Department of Chemistry, The Ohio State University, Columbus, OH 43210.*

Single-photon Laser Induced Fluorescence has been used to measure the trans-rotational temperature of mildly vibrationally excited carbon monoxide ($v'' \leq 9$) within the positive column of a CO/He normal d.c. glow discharge.^a A tunable, narrowband ArF excimer laser at 193nm was used to pump the strongly allowed CO $X^1\Sigma^+(v''=7) \rightarrow A^1\Pi(v'=1)$ transition with subsequent collection of the 200.8nm (1,8) 4th Positive emission.^b The resulting rotational spectral peaks were assigned and a subset was used to estimate the rotational temperature to be $432 \pm 72\text{K}$ (3σ). Fourier Transform – InfraRed spectroscopic measurements of this plasma were in good agreement at $395 \pm 10\text{K}$. These results indicate that within a normal glow discharge environment, CO 4th Positive LIF is well suited as a temperature diagnostic.

^a S. DeBenedictis, *Chem. Phys.*, **71**, p.247, 1982.

^b G.L. Wolk and J.W. Rich, *Chem. Phys. Lett.* **87**, p. 117, 1982.