

FOURIER TRANSFORM MICROWAVE SPECTROSCOPIC AND AB INITIO STUDIES OF THE Kr-CH<sub>4</sub> AND Ar-CH<sub>4</sub> VAN DER WAALS COMPLEXES

YAQIAN LIU and WOLFGANG JÄGER, *Department of Chemistry, University of Alberta, Edmonton, AB, Canada, T6G 2G2.*

Rotational spectra of various isotopomers of the Kr-CH<sub>4</sub> van der Waals complex were recorded using a pulsed molecular beam Fourier transform microwave spectrometer. Transitions corresponding to two excited internal rotor states, namely the  $j=1, K=0$  and  $j=2, K=1$  states, were measured and assigned, in addition to the previously reported ground state transitions.<sup>a</sup> The  $j=1, K=0$  state was found to be Coriolis perturbed by a nearby  $j=1, K=1$  state. The spectra of the isotopomers containing <sup>83</sup>Kr showed nuclear quadrupole hyperfine structure. Rotational, centrifugal distortion, and quadrupole coupling constants were determined and were used to obtain structural parameters. *Ab initio* potential energy surfaces of Kr-CH<sub>4</sub> and Ar-CH<sub>4</sub> were constructed at the CCSD(T) level of theory with the aug-cc-pVTZ basis set supplemented with bond functions. The dipole moments of the two complexes were also computed at various configurations. The obtained results from the *ab initio* calculations were compared with those from the experiment.

---

<sup>a</sup>Y. Liu and W. Jäger, 55th International Symposium on Molecular Spectroscopy, paper RE05 (2000).