

INTENSITIES, SELF- AND AIR-BROADENING, AND PRESSURE-INDUCED LINE SHIFTS IN THE $2\nu_2$ BAND OF HCN

V. MALATHY DEVI, D. CHRIS BENNER, *Department of Physics, The College of William and Mary, Box 8795, Williamsburg, VA 23187-8795*; M. A. H. SMITH, C. P. RINSLAND, *Atmospheric Sciences, NASA Langley Research Center, Mail Stop 401A, Hampton, VA 23681-2199*; S. W. SHARPE and R. L. SAMS, *Mail Stop K8-88, Battelle Blvd., Pacific Northwest National Laboratory, P. O. Box 999, Richland, WA 99352*.

High-resolution infrared absorption spectra of HCN in the $2\nu_2$ band region near 1411 cm^{-1} have been recorded with the Bruker IFS 120 HR Fourier transform spectrometer located at Pacific Northwest National Laboratory. The measurements were all made at ambient temperatures between 24.7 and 24.9°C , although the variation during the course of recording each spectrum was less than $\pm 0.01^\circ\text{C}$. Two stainless steel absorption cells that were electroplated and gold coated to minimize reactivity were used to obtain the spectra; their path lengths were 0.958 and 19.95 cm . Four spectra of high-purity (99.8%) HCN together with three spectra of lean mixtures ($\sim 3\%$) of HCN in dry air were simultaneously fit using a multispectrum nonlinear least squares procedure.^a The analysis yielded room-temperature values for absolute intensities, self- and air-broadening coefficients and self- and air-induced pressure shift coefficients for numerous lines in the $2\nu_2$ band of $\text{H}^{12}\text{C}^{14}\text{N}$. Since there are no previous measurements of broadening and shift parameters in the $2\nu_2$ band, our results will be compared with values recently determined in the ν_1 band of $\text{H}^{12}\text{C}^{14}\text{N}$ ^b and with current HITRAN values.^c

^aD. Chris Benner, C. P. Rinsland, V. Malathy Devi, M. A. H. Smith and D. Atkins, *JQSRT* **53**, 705-721 (1995).

^bC. P. Rinsland et al., *JQSRT*, in press (2003).

^cL. S. Rothman et al., *JQSRT*, in press (2003).