

## VIBRONIC STRUCTURES OF THE MgNC $\tilde{A}^2\Pi$ AND $\tilde{X}^2\Sigma^+$ STATES

MASARU FUKUSHIMA and TAKASHI ISHIWATA, *Faculty of Information Sciences, Hiroshima City University, Asa-Minami, Hiroshima 731-3194, Japan.*

We have generated MgNC in supersonic free jet expansions, and measured the laser induced fluorescence excitation spectra of the vibronic bands of the  $\tilde{A}^2\Pi - \tilde{X}^2\Sigma^+$  transition. We have found some new features in the spectra, though part of the spectra were already reported by Wright and Miller<sup>a</sup>. The new features are consisting of the  $\nu_2$  bending vibronic bands. From the rotational analyses of some of the vibronic bands, we have obtained information on the vibronic structure of the  $\tilde{A}^2\Pi$  state of MgNC. The assignments of the  $\nu_2$  vibronic bands were carried out utilizing the vibrational analyses of the dispersed fluorescence spectra obtained by the excitation of the  $\nu_2$  vibronic bands. On the basis of the vibrational structures of the dispersed fluorescence spectra which are not only those of the  $\nu_2$  vibronic bands but also the spectra obtained by the excitation of the other vibronic bands reported by us<sup>b</sup>, the vibronic structure of the  $\tilde{X}^2\Sigma^+$  state of MgNC has been analysed. The vibronic structures of the  $\tilde{A}^2\Pi$  and  $\tilde{X}^2\Sigma^+$  states of MgNC will be discussed comparing with the computational results by Hirano group<sup>c</sup>.

---

<sup>a</sup>R. R. Wright and T. A. Miller, *J. Mol. Spectrosc.* **194**, 219 (1999).

<sup>b</sup>M. Fukushima and T. Ishiwata, *J. Mol. Spectrosc.* **216**, 159 (2002).

<sup>c</sup>T. E. Odaka, T. Taketsugu, T. Hirano and U. Nagashima, *J. Chem. Phys.* **115**, 1349 (2001); T. E. Odaka, T. Hirano and P. Jansen, *J. Mol. Spectrosc.* **211**, 147 (2002); T. Hirano, K. Ishii T. E. Odaka and P. Jansen, *J. Mol. Spectrosc.* **215**, 42 (2002); T. E. Odaka, T. Hirano and P. Jansen, *J. Mol. Spectrosc.* **216**, 379 (2002).