

## MICROWAVE SPECTRAL STUDIES OF BENZYL ALCOHOL AND SEC-PHENETHANOL

KARISSA UTZAT, ROBERT K. BOHN, AND EMILY CHU, *Department of Chemistry, University of Connecticut, Storrs, CT 06269-3060.*

Benzyl compounds ( $C_6H_5C_\alpha H_2X$ ), generally fall into two conformational classes. With  $X=CH_3$ ,  $C_2H_5$ , F, or Cl, the C-C-C $_\alpha$ -X dihedral angle is  $90^\circ$ ; with  $X=CN$  and CCH, the dihedral angle is  $0^\circ$ ; however, benzyl alcohol ( $C_6H_5CH_2OH$ ) and benzyl amine ( $C_6H_5CH_2NH_2$ ) have asymmetric structures. Two conformers of benzyl amine have been observed,<sup>a</sup> one with a dihedral angle approximately  $50^\circ$  and a higher energy form with dihedral angle  $90^\circ$ . Benzyl alcohol has a single asymmetric conformation with dihedral, C-C-C $_\alpha$ -O about  $60^\circ$ .<sup>b</sup> The microwave rotational spectrum displays tunneling doublets for all transitions indicating a low barrier path between two of the four equivalent energy minima. These splittings are significantly smaller in the  $C_6H_5CH_2OD$  spectrum. The rotational spectrum of sec-phenethanol, ( $C_6H_5C_\alpha H(C_\beta H_3)OH$ ), displays a single conformation with C-C-C $_\alpha$ -C $_\beta$  dihedral angle approximately  $90^\circ$  and the C-C-C $_\alpha$ -O dihedral angle approximately  $30^\circ$ , and no tunneling split transitions. *ab initio* calculations on sec-phenethanol and benzyl alcohol will also be discussed.

---

<sup>a</sup>S. Melandri, A. Maris, P.G. Favero, W. Caminati, CHEMPHYSICHEM 2(3), 172-177 (2001).

<sup>b</sup>M. Trættemberg, H. Østensen, R. Seip, Acta Chemica Scandinavica A 34, 449-454 (1980).