

ELECTRONIC STATE OF CoH: A PROBLEM RELATED TO THE SIZE-INCONSISTENCY OF THE MR-SDCI METHOD AND THE MIXING OF $^3\Phi$ AND $^3\Pi$

REI FUKUI, UMPEI NAGASHIMA, and TSUNEO HIRANO, *Grid Technology Research Center, Institute of Advanced Industrial Science and Technology, 6-9-3 Ueno, Taito-ku, Tokyo 110-0015, Japan*; HIROSHI TATEWAKI, *Computation Center and Institute of Natural Sciences, Nagoya City University, Nagoya 467-8501, Japan*; TAKESHI NORO AND KIYOSHI TANAKA, *Division of Chemistry, Graduate School of Science, Hokkaido University, Kita 10, Nishi 8, Kita-ku, Sapporo, 060-0810, Japan*.

The ground state of CoH has been determined experimentally as the $^3\Phi$ by the far-infrared magnetic resonance study,^a and theoretically by *ab initio* methods.^b However, MR-SDCI and MR-SDCI + Q *ab initio* molecular orbital calculations predicted that the lowest electronic state is $^5\Phi$. The size-consistent MR-ACPF method correctly predicted that $^3\Phi$ should be the ground state. As is previously reported by Tanaka *et al.*^c on FeH, the controversial results in MR-SDCI may arise through truncation error inherent in the size-inconsistent MR-SDCI method.

Another difficulty is that we cannot clearly separate the Φ and Π states when CoH is treated under C_{2v} symmetry, not under its real symmetry of $C_{\infty v}$. The best results we obtained at the moment are: $r_e = 1.528 \text{ \AA}$, $B_e = 7.291 \text{ cm}^{-1}$, and $\omega_e = 2005 \text{ cm}^{-1}$ at the level of the MR-ACPF + relativistic correction using the Roos ANO (Co) and Dunning aug-cc-pVQZ (H) basis sets.

^aS.P. Beaton, K.M. Evenson, T. Nelis, and J.M. Brown, *J. Chem. Phys.*, **89**, 4446-4448 (1988).

^bD.P. Chong, S.R. Langhoff, C.W. Bauschlicher, Jr., S.P. Walch, and H. Partridge, *J. Chem. Phys.*, **85**, 2850-2860 (1986); M. Freindorf, C. M. Marian, and B. A. Hess, *J. Chem. Phys.*, **99**, 1215-1223 (1993).

^cK. Tanaka, M. Sekiya, and M. Yoshimine, *J. Chem. Phys.*, **115**, 4558-4564 (2001).