

REACTION PRODUCTS OF LASER ABLATED EARLY TRANSITION METALS WITH BENZENE MOLECULES  
IN CONDENSING ARGON

JONATHAN T. LYON, LESTER ANDREWS, *Department of Chemistry, University of Virginia, Charlottesville, VA 22904-4319.*

Early transition metal atoms, produced by laser ablation, are reacted with benzene vapor diluted in argon during co-deposited onto a low temperature CsI window. The resulting reaction products are trapped and the M(C<sub>6</sub>H<sub>6</sub>) and M(C<sub>6</sub>H<sub>6</sub>)<sub>2</sub> complexes are identified by benzene isotopic studies (C<sub>6</sub>H<sub>6</sub>, <sup>13</sup>C<sub>6</sub>H<sub>6</sub>, C<sub>6</sub>D<sub>6</sub>). Density Functional Theory (DFT) frequency calculations are used to confirm the assignments. Calculated ground state energies predict these reaction products to be energetically favorable. The M(C<sub>6</sub>H<sub>6</sub>) and M(C<sub>6</sub>H<sub>6</sub>)<sub>2</sub> products have C<sub>6v</sub> and D<sub>6h</sub> symmetries respectively. Based on the observed aromatic C-C breathing mode frequency shifts, the metal-carbon bond strengths are shown to increase down a group (from vanadium to tantalum) and decrease from left to right along a period (from scandium to chromium). These trends indicate that back donation of electrons from the metal d orbitals to the unoccupied  $\pi^*$  orbitals of benzene is the primary electronic interaction responsible for the bonding in these molecules. Analogous experiments in progress with C<sub>60</sub> give similar product complexes.