

## VACUUM UV LIF MEASUREMENTS OF RO-VIBRATIONALLY EXCITED HYDROGEN MOLECULES IN A SUPERSONIC PLASMA EXPANSION

P. VANKAN, S. B. S. HEIL, D.C. SCHRAM, and R. ENGELN, *Department of Applied Physics, Centre for Plasma Physics and Radiation Technology (CPS), Eindhoven University of Technology, P.O. Box 513, NL-5600 MB Eindhoven, The Netherlands.*

Rotationally and vibrationally excited hydrogen molecules are measured using Laser Induced Fluorescence (LIF) spectroscopy in the vacuum-UV. The laser radiation is generated via the Stimulated Anti-Stokes Raman Scattering (SARS) process, and is tunable between 120 nm and 230 nm. Up to 10 Anti-Stokes beams are simultaneously directed into the hydrogen plasma expansion. This so-called multiplex excitation makes an accurate wavelength calibration of the pump laser imperative. This calibration is performed by simultaneously recording the VUV-LIF signal from the plasma expansion and the well-known two-photon laser induced fluorescence spectrum of nitric oxide in a gas cell<sup>a</sup>. The density of hydrogen molecules in the ro-vibrationally excited energy levels is calibrated on the basis of previously performed Coherent Anti-Stokes Raman Scattering measurements. Population distributions are measured in an expanding plasma to determine the production mechanism of the ro-vibrationally excited hydrogen molecules.

In this contribution we present results on the population distribution of  $\text{H}_2^{\nu}$  from  $\nu = 2$  to  $\nu = 6$  in the electronic ground state. The ro-vibrational population distribution in the expanding hydrogen plasma can be described with a two temperature Boltzmann distribution. The population distribution in the low rotational levels ( $J \leq 5$ ) in the measured vibrational states can be described with a temperature of 800 K, while the higher rotational levels are populated according to a *vibrational-like* temperature of 3600K. In the  $\nu = 2$  state rotational excitation is observed up to  $J = 19$ . A possible generation mechanism for the *super-rotational* excitation is the association of atomic hydrogen at the reactor wall.

---

<sup>a</sup>P. Vankan, S. B. S. Heil, S. Mazouffre, R. Engeln, D. C. Schram, H. F. Döbele, *accepted for publication in Rev. Sci. Instrum.*, issue April 2004