

SELF- AND H₂-BROADENING AND SHIFT COEFFICIENTS IN THE 2 ← 0 BAND OF ¹²C¹⁶O: REVISITED

V. MALATHY DEVI, D. CHRIS BENNER, *Department of Physics, The College of William and Mary, Williamsburg, VA 23187-8795*; A. PREDOI-CROSS, *Department of Physics, The University of Lethbridge, Lethbridge, AB T1K 3M4, Canada*; M. A. H. SMITH, C. P. RINSLAND, *Atmospheric Sciences, NASA Langley Research Center, Hampton, VA 23681-2199*; and A. W. MANTZ, *Department of Physics, Astronomy and Geophysics, Connecticut College, New London, CT 06320*.

Room temperature values for self-broadened and hydrogen-broadened Lorentz half width coefficients, and self and hydrogen pressure-induced shift coefficients have been measured for transitions with rotational quantum number index m ranging from -24 to $+24$ in the $2 \leftarrow 0$ band of ¹²C¹⁶O. The spectra were recorded with the McMath-Pierce Fourier transform spectrometer located at the National Solar Observatory on Kitt Peak. The analysis was performed using a multispectrum nonlinear least squares technique^a modified to constrain the Lorentz widths in the P and R branches to be identical for the same $|m|$ value. We have compared the present results with our previous measurements made with the same spectrometer^b and with other measurements published recently.

^aD. Chris Benner, C. P. Rinsland, V. Malathy Devi, M. A. H. Smith and D. Atkins, *JQSRT* **53**, 705-721 (1995).

^bV. Malathy Devi, D. Chris Benner, M. A. H. Smith, C. P. Rinsland and A. W. Mantz, *JQSRT* **75**, 455-471 (2002).