

INFRARED PHOTODISSOCIATION SPECTROSCOPY OF PROTONATED WATER CLUSTERS NEAR THE MAGIC NUMBER $H^+(H_2O)_{21}$

R. S. WALTERS, T. D. JAEGER, M. A. DUNCAN, *Department of Chemistry, University of Georgia, Athens, GA, 30602-2556*; J. W. SHIN, N. I. HAMMER, E. G. DIKEN, M. A. JOHNSON, *Sterling Chemistry Laboratory, Yale University, New Haven, CT, 06520*; R. A. CHRISTIE and K. D. JORDAN, *Department of Chemistry, University of Pittsburgh, Pittsburgh, PA, 15260*.

$H^+(H_2O)_n$ complexes in the intermediate size range ($n=10-30$) are studied by infrared photodissociation spectroscopy near the symmetric and asymmetric stretching vibrations of water (3657 cm^{-1} , 3756 cm^{-1}). The complexes fragment by the loss of intact, multiple water molecules and their infrared spectra are compared to the predictions of theory. The spectrum for the magic number $n=21$ cluster shows only one distinct feature in the O-H region indicating all the water molecules occupy similar binding sites. Surprisingly, a single band is also observed for $n=22$ and no photodissociation is observed for the calculated fundamental of the H_3O^+ ion core near 2500 cm^{-1} . Possible interpretations of the above results will be discussed.