

DIPOLE MOMENTS OF AMINE HYDROGEN HALIDE COMPLEXES

C. S. BRAUER, M. B. CRADDOCK, E. M. GRUMSTRUP, and K. R. LEOPOLD, *Department of Chemistry, University of Minnesota, Minneapolis, MN 55455*; M. C. ORILALL, *Department of Chemistry, Concordia University, Moorhead, MN 56562*.

Amine hydrogen halide complexes are prototypical systems in which to study the transfer of a proton across a hydrogen bond. We report the results of Stark effect experiments using a pulsed nozzle Fourier transform microwave spectrometer to determine the dipole moments of: $\text{H}_3^{15}\text{N-H}^{35}\text{Cl}$ (4.05865 ± 0.00095 D), $(\text{CH}_3)_3^{15}\text{N-H}^{35}\text{Cl}$ (7.129 ± 0.014 D), $\text{H}_3^{15}\text{N-H}^{79}\text{Br}$ (4.2577 ± 0.0022 D), and $(\text{CH}_3)_3^{15}\text{N-H}^{79}\text{Br}$ (8.396 ± 0.013 D). The results will be discussed in terms of the degree of proton transfer, as previously elucidated by Legon and coworkers^a and should be of interest in view of a large body of matrix isolation work in which the interaction between the complex and the host matrix has been investigated.

^aA. C. Legon *Chem. Soc. Rev.* **22**(3), 153, (1993).