

## INTRAMOLECULAR DYNAMICS IN $X_4$ RHOMBIC CLUSTERS

ROBERTO LINGUERRI, ISABELLE NAVIZET, and PAVEL ROSMUS, *Laboratoire de Chimie Théorique, Université de Marne la Vallée, F 77454 - Champs sur Marne, France*; STUART CARTER, *Department of Chemistry, University of Reading, Reading RG6 2AD, U.K.*; JOHN P. MAIER, *Department of Chemistry, University of Basel, Klingelbergstr. 80 CH-4056 Basel, Switzerland*.

Two independent approaches are applied to study the intra-molecular dynamics in  $B_4$  electronic ground state. First, a double minimum six-dimensional potential energy surface (PES) is determined in symmetry coordinates for the most stable rhombic ( $D_{2h}$ )  $B_4$  isomer in its  $^1A_g$  electronic ground state by fitting to energies calculated by the CCSD(T) approach. The PES exhibits a barrier to the  $D_{4h}$  square structure of  $255\text{ cm}^{-1}$ . The vibrational levels ( $J = 0$ ) are calculated variationally using the MULTIMODE<sup>a</sup> code, which involves the Watson kinetic energy operator expressed in normal coordinates. The pattern of about 65 vibrational levels up to  $1600\text{ cm}^{-1}$  for all stable isotopomers is analyzed. Analogous to the inversion in ammonia-like molecules, the rhombus rearrangements lead to splittings of the vibrational levels. In  $B_4$  it is the  $b_{1g}$  ( $D_{4h}$ ) mode which distorts the square molecule to its planar rhombic form. It is shown that the rhombic ground states of  $B_4^+$  and  $B_4^-$  consist of two vibronically coupled electronic states. The PES for  $B_4$  is used also in time-dependent molecular dynamics calculations using the multi-configurational time-dependent Hartree<sup>b</sup> (MCTDH) method. The 6-D wavepackets are analyzed and the results are compared with time-independent results.

<sup>a</sup>S. Carter, S. Culik and J. M. Bowman *J. Chem. Phys.* **107**, 10458 (1997).

<sup>b</sup>H.-D. Meyer, U. Manthe, and L. S. Cederbaum. *Chem. Phys. Lett.* **165**, 73 (1990).