

TIME-RESOLVED INFRARED DIODE LASER SPECTROSCOPY OF THE ν_1 BAND OF THE FeNO RADICAL PRODUCED BY THE ULTRAVIOLET LASER PHOTOLYSIS OF $\text{Fe}(\text{CO})_2(\text{NO})_2$

SEIKI IKEDA, MOTOKI NAKASHIMA and KEIICHI TANAKA, *Department of Chemistry, Faculty of Science, Kyushu University, Hakozaki, Higashiku, Fukuoka, 812-8581 JAPAN.*

Rovibrational transitions of the ν_1 band (N-O stretch) of the FeNO radical were observed in the 1750-1780 cm^{-1} region. The FeNO radical was produced by 193 nm excimer laser photolysis of $\text{Fe}(\text{CO})_2(\text{NO})_2$ and the transient absorption signal was detected by time-resolved infrared diode laser spectroscopy.

More than 40 lines were assigned to the ν_1 fundamental band of the $\Omega = 5/2$ spin component, together with ten Q-branch lines ($J = 2.5 - 11.5$) in the 1760 cm^{-1} region, to confirm the electronic ground state to be $X^2\Delta_i$. Effective molecular constants for the $\Omega = 5/2$ spin component, including the band origin ν_0 (1767.26093(38) cm^{-1}), the rotational constant B (4610.17754(93) MHz) and the centrifugal distortion constant D (1.17003(47) kHz), were derived from a least squares fitting of the observed transitions. The average bond length $r_{\text{Co-N}}$ between Co and N was calculated to be 1.621 Å from the rotational constant B_0 assuming $r_{\text{N-O}} = 1.186$ Å as given by *ab initio* calculation^a. The ν_1 hot band lines originated from the ν_2 (Fe-N-O bending; 308 cm^{-1} ^a) vibrationally excited state were also observed. Pure rotational lines of FeNO were also observed by the millimeter wave spectroscopy with the UV photolysis of $\text{Fe}(\text{CO})_2(\text{NO})_2$ ^b.

^aM. Zhou and L. Andrews, *J.Phys.Chem.A*, **104**, 3915 (2000)

^bMicrowave session in this symposium.