

FIR SPECTROSCOPY OF BARE CH_5^+ USING LASER INDUCED REACTIONS

OSKAR ASVANY, EDOUARD HUGO, *Leiden Observatory, P.O. Box 9513, 2300 RA Leiden, The Netherlands*; STEPHAN SCHLEMMER, *I. Physikalisches Institut, Universität zu Köln, Zùlpicher Straße 77, 50937 Köln, Germany*.

Laser Induced Reaction (LIR) is a novel technique for the spectroscopy of molecular ions as well as the study of state-specific rate coefficients and lifetimes of excited states. It is based on trapping mass-selected ions in the cold environment of a reactant gas and probing the laser-induced excitation of the ion by the outcome of the corresponding endothermic ion-molecule reaction. The feasibility of this method has been demonstrated in the past for the electronic spectroscopy of N_2^+ in the charge transfer reaction with Ar, and the spectroscopy of the IR-active modes of the acetylene cation in the abstraction reaction of C_2H_2^+ with hydrogen^{a,b}.

Recently, LIR has been applied to record the first far infrared spectra of bare CH_5^+ and some of its deuterated versions combining a low temperature 22-pole ion trap with the large tuning range of the free electron laser FELIX. To obtain the LIR spectrum, the product ions of the endothermic proton transfer reaction $\text{CH}_5^+ + \text{CO}_2 \rightarrow \text{CH}_4 + \text{OCOH}^+$ have been counted as a function of the FELIX wavelength. In addition to the earlier observed C–H-stretching feature^c at 3000 cm^{-1} , low-frequency extensions down to 2200 cm^{-1} become discernible, as well as a H–C–H bending feature at 1250 cm^{-1} . Comparison of the measured data to computed ones shows that CH_5^+ is fluxional at the temperature of the experiment, $T=110\text{K}$.

^aS. Schlemmer, E. Lescop, J. v. RIchthofen, and D. Gerlich, *J. Chem. Phys.* **117**, 2068–2075 (2002)

^bO. Asvany, T. Giesen, B. Redlich, and S. Schlemmer, *Phys. Rev. Lett.* **94**, 073001, (2005)

^cE. T. White, J. Tang, and T. Oka, *Science* **284**, 135–137 (1999)