

RESOLUTION OF A CONVERGENCE PROBLEM IN DIRECT-POTENTIAL-FIT DATA ANALYSES USING THE HERMAN-ASGHARIAN HAMILTONIAN

ROBERT J. LE ROY, *Department of Chemistry, University of Waterloo, Waterloo, Ontario N2L 3G1, Canada*; JOHN A. COXON, *Department of Chemistry, Dalhousie University, Halifax, Nova Scotia, B3H 4J3 Canada*; MICHAEL REY and VLADIMIR G. TYUTEREV, *Group de Spectrométrie Moléculaires et Atmosphérique, CNRS UMR 6089, BP 1039, F-51687, Reims Cedex 2, France*.

The effective radial Schrödinger equation based on the Herman-Asgharian^a Hamiltonian for a diatomic molecule in a $^1\Sigma$ state has the form

$$-\frac{\hbar^2}{2\mu} [1 + \beta(r)] \frac{d^2 \psi_{v,j}(r)}{dr^2} + \left\{ [V_{\text{CN}}(r) + \Delta V_{\text{ad}}(r)] + \frac{\hbar^2}{2\mu r^2} [1 + \alpha(r)] [J(J+1)] \right\} \psi_{v,j}(r) = E_{v,J} \psi_{v,j}(r) \quad (1)$$

in which $\beta(r)$ and $\alpha(r)$ represent the effects of non-adiabatic corrections to the radial and angular kinetic energy operators, respectively, and $\Delta V_{\text{ad}}(r)$ is the adiabatic correction to the “clamped nuclei” potential energy function $V_{\text{CN}}(r)$. An internal convergence problem encountered when utilizing wavefunction propagator methods for direct-potential-fit diatomic data analyses using this Hamiltonian is described and corrected. Improved Hamiltonian parameters for the ground states of GaH and ArH⁺ will be reported.

^a R. M. Herman and A. Asgharian, *J. Mol. Spectrosc.* **19**, 305 (1966).