

MAGNETIC HYPERFINE INTERACTIONS IN ^{171}YbF AND ^{173}YbF ^a.

TIMOTHY C. STEIMLE, TONGMEI MA, *Department of Chemistry and Biochemistry, Arizona State University, Tempe, AZ, USA 85287*; COLAN LINTON, *Physics Department, University of New Brunswick, Fredericton, NB, Canada E3B 5A3*.

The heavy polar molecule ytterbium monofluoride, YbF, has been used to set an upper limit on the electric-dipole moment of the electron, d_e ^b. The extraction of d_e from experimental data requires knowledge of the effective electric field in the region of the Yb nucleus, which can only be obtained from electron structure calculations^c. The most effective means of gauging the quality of the predicted wavefunction in the region of the nucleus is via a comparison with experimental measurements of the magnetic hyperfine interactions. Here we report on $^{171}\text{Yb}(I=1/2)$, $^{173}\text{Yb}(I=5/2)$, and $^{19}\text{F}(I=1/2)$ magnetic hyperfine interactions from the analysis of the $A^2\Pi_{1/2} - X^2\Sigma^+$ (0,0) band of ytterbium monofluoride, YbF, recorded at the near-natural linewidth limit. The results are compared with *ab initio* predictions^b, the microwave measurements for ^{174}YbF ^d, and the matrix-isolated ESR results for ^{171}YbF ^e.

^aSupported by NSF-Experimental Physical Chemistry

^bJ.J. Hudson, B.E. Sauer, M.R. Tarbutt, and E.A. Hinds, *Phys. Rev. Lett.* **98**, 023003-4 (2002).

^cA.V. Titov, N.S. Mosyagin, A.N. Petrov, and T.A. Iseav, *Int. J. Quantum Chem.* **104**, 223-239 (2005).

^dC.S. Dickinson, J.A. Coxon, N.R. Walker, and M.C.L. Gerry, *J. Chem. Phys.* **115**, 6979-6989 (2001).

^eR.J. Van Zee, M.L. Seely, T.C. DeVore, and W. Weltner, Jr. *J. Chem. Phys.* **82**, 1192-1194 (1978).