

## FOURIER-TRANSFORM SPECTROSCOPY OF $^{14}\text{NH}_3$ AND $^{15}\text{NH}_3$ IN THE NEAR-INFRARED

N. IBRAHIM, P. CHELIN, and J. ORPHAL, *Laboratoire Interuniversitaire des Systèmes Atmosphériques (LISA), CNRS and Universités Paris-7 et -12, Créteil, France*; C. E. FELLOWS, *Laboratório de Espectroscopia e Laser, Instituto de Física, Universidade Federal Fluminense, Niterói, Rio de Janeiro, Brazil*.

The near-infrared absorption spectrum of  $\text{NH}_3$  is an interesting opportunity for atmospheric *in-situ* measurements of this species using tuneable diode-lasers. For this purpose, accurate line positions and intensities are indispensable.

In the past, several studies of the absorption spectrum of  $^{14}\text{NH}_3$  in the near-infrared were carried out, using Fourier-transform<sup>1</sup> and tuneable diode-laser spectroscopy.<sup>2-4</sup> More recently, high-resolution diode-laser spectra of  $^{15}\text{NH}_3$  were analyzed for the first time.<sup>5</sup> The lines of  $^{14}\text{NH}_3$  in this region are also useful for wavenumber calibration of absorption spectra obtained with tuneable lasers.

In our group, we have recently investigated the possibility to detect atmospheric  $\text{NH}_3$  using photoacoustic laser spectroscopy in the 1.5  $\mu\text{m}$  region, and observed several discrepancies between individual line positions and intensities observed in our and previous studies<sup>1,4</sup>, as already noticed by other groups.<sup>2,5</sup> In order to solve this problem we have measured new absorption spectra of  $^{14}\text{NH}_3$  (and also of  $^{15}\text{NH}_3$ ) in the 6300-7500  $\text{cm}^{-1}$  region using a Bruker IFS 120-HR Fourier-transform spectrometer. The length of the absorption cell was 30 cm. The  $\text{NH}_3$  pressures employed were around 30 mbar, and the spectral resolution used was 0.02  $\text{cm}^{-1}$ , leading to linewidths (Full-Width at Half-Maximum) of about 0.03  $\text{cm}^{-1}$ . Based on these spectra, we have produced a list of individual line positions and line intensities at 296 K. Comparisons with the previous studies<sup>1-5</sup> will be presented.

---

[1] L. Lundsberg-Nielsen et al., *J. Mol. Spectrosc.*, **162**, 230-245 (1993).

[2] M. Webber et al., *Appl. Opt.*, **40**, 2031-2042 (2004).

[3] L.-H. Xu et al., *Infrared Phys. Technol.*, **45**, 31-45 (2004).

[4] J. S. Gibb et al., *Eur. Phys. J. D.*, **28**, 59-66 (2004).

[5] R. M. Lees et al., *J. Mol. Struct.*, in press (2006).