

## REINVESTIGATION OF THE GROUND AND FIRST TORSIONAL STATES OF METHYLFORMATE

M.. CARVAJAL, *Departamento de Física Aplicada, Facultad de Ciencias Experimentales, Universidad de Huelva, 21071 Huelva, Spain*; F. WILLAERT, J.. DEMAISON, *Laboratoire de Physique des Lasers, Atomes et Molécules, CNRS et Université de Lille I, F-59655, Villeneuve d'Ascq Cédex, France*; I. KLEINER, *Laboratoire Interuniversitaire des Systèmes Atmosphériques, CNRS et Universités Paris 7 et Paris 12, 61 av. Général de Gaulle, 94010, Créteil, France*.

We have reinvestigated the laboratory spectrum for the methylformate molecule involving both the ground and first torsional states. We have fitted within almost experimental accuracy a data set for HCOOCH<sub>3</sub> consisting of 3496  $v_t=0$  and 774  $v_t=1$  microwave lines, using the so-called "rho axis method" (RAM) and a model extended to include perturbation terms through eight order. The previously published  $v_t=0$  and  $v_t=1$  microwave lines covering J values up to 43 in the ground and up to 18 in the first excited states have been extended by new measurements from Lille with maximum values of J = 62. The final fit requires only 49 parameters to achieve a weighted standard deviation for the whole fit of 1.43 for a total of 4270 lines. This result represents an improvement over the previous fit which achieved a standard deviation of 1.96 for 3862 lines using 69 parameters. A calculation of the linestrengths of torsion-rotation transition up to J=60 needed for the astronomical survey is also presented.