

ANALYSIS OF ROTATIONAL STRUCTURE IN THREE C-TYPE BANDS IN THE HIGH-RESOLUTION INFRARED SPECTRUM OF *TRANS*-1,3,5-HEXATRIENE

NORMAN C. CRAIG, MATTHEW C. LEYDEN, and AMIE K. PATCHEN, *Department of Chemistry and Biochemistry, Oberlin College, Oberlin, OH 44074*; TONY MASIELLO, *Environmental and Molecular Sciences Laboratory, Pacific Northwest National Laboratory, P.O. Box 999, Mail Stop K8-88, Richland, WA 99352*.

We seek a semi-experimental equilibrium structure for *trans*-1,3,5-hexatriene, which should show structural evidence for increased pi-electron delocalization in comparison with butadiene. Despite the marginal resolution in the spectrum of *trans*-hexatriene, rotational structure of three C-type bands of *trans*-hexatriene in the high-resolution infrared spectrum (0.0015 cm^{-1}) has been analyzed as a first step. Strong bands are at 1010.95 and 901.910 cm^{-1} ; a weaker band is at 683.45 cm^{-1} . The most extensive analysis was for the band at 901.910 cm^{-1} . The composite fit of 1628 ground state combination differences from the three bands gave the ground state rotational constants of $A = 0.8742480(7)$, $B = 0.0446600(10)$, and $C = 0.0425099(8)\text{ cm}^{-1}$. For this near-symmetric top, $\kappa = -0.99483$. Procedures for synthesizing needed isotopomers are being explored.