

IR SPECTROSCOPY OF LONG CARBON CHAINS AND THEIR DERIVATIVES IN CRYOGENIC MATRICES:
SPECIES IDENTIFICATION BY $^{13}\text{C}_3 \rightarrow ^{12}\text{C}_3$ ISOTOPIC SUBSTITUTION.

DMITRY STRELNIKOV and WOLFGANG KRÄTSCHMER, *Max-Planck-Institut für Kernphysik, D-69029
Heidelberg, PO Box 103980, Germany.*

Along with matrix gas, we co-deposited carbon vapor from two separated carbon evaporation sources, namely a ^{12}C -source and a ^{13}C -source. Under the applied conditions the C_3 molecules are the most abundant species in both carbon sources. We observed that in the formation process of long carbon chains the C_3 units play a major role. This finding was used to establish a new species identification method based on isotopic replacement. The old technique of atomic $^{13}\text{C} \rightarrow ^{12}\text{C}$ substitution can be reasonably applied only for molecules with a small number of carbon atoms ($n \leq 9$). Using the new method of molecular $^{13}\text{C}_3 \rightarrow ^{12}\text{C}_3$ substitution, quite simple distributions having small number of isotopomeric IR absorptions were obtained also for larger species. We present data for C_n and $\text{C}_n\text{O}_{1,2}$ ($n \leq 6$).