

VIBRATIONAL SPECTRA AND *AB-INITIO* CALCULATIONS FOR THE STUDY OF INTRAMOLECULAR VIBRATIONAL REDISTRIBUTION IN THE CH-CHROMOPHORE IN CHD₂I

VERONIKA HORKÁ, MARTIN QUACK, MARTIN WILLEKE, *Laboratorium für Physikalische Chemie, ETH Zürich, CH-8093, Zürich, Switzerland.*

Intramolecular vibrational energy redistribution (IVR) on the femto- to picosecond time scale is a central primary process in unimolecular chemical reaction dynamics. There are two main approaches towards its understanding. One is the analysis of the highly resolved vibrational spectra of polyatomic molecules^{a,b} combined with *ab initio* calculations. Another approach consists in time resolved femtosecond (fs) pump-probe experiments, which we have recently carried out for a series of alkyl iodides^c including CHD₂I. For the latter molecule we present here results for IVR processes within the CH-chromophore. The infrared absorption spectrum of CHD₂I from 500 to 12000 cm⁻¹ has been recorded with a Bomem Fourier transform infrared spectrometer equipped with a White type multi reflection cell and resolutions of 0.10 and 0.15 cm⁻¹. The CH-chromophore bands dominate the vibrational overtone spectrum of CHD₂I and we have assigned 24 of them. Vibrational variational calculations were carried out for the overtone spectrum of the CH-chromophore based on a 3-dimensional *ab initio* potential energy and electric dipole moment surface in normal coordinates. The anharmonic resonance structure between the CH-stretching and bending modes was investigated via an effective Hamiltonian for both experimental and theoretical spectra. Finally, we have calculated the time dependent dynamics on the fs time scale^d.

^aM. Quack, *Annu. Rev. Phys. Chem.*, 41, 839 (1990); M. Quack, Chapter 27 in *Femtosecond Chemistry* (ed. by J. Manz and L. Woeste), Verlag Chemie, Weinheim, 781 (1995); M. Quack, *Chimia*, 55 (10), 753 (2001) and references therein.

^bJ. Pochert, M. Quack, J. Stohner, and M. Willeke, *J. Chem Phys.*, 113, 2719 (2000).

^cV.N. Krylov, M.V. Nikitchenko, M. Quack, and G. Seyfang, *Proc. SPIE* 5337, 178 (2004); V.N Krylov, A. Kushnarenko, E. Miloglyadov, M. Quack, and G. Seyfang, *Proc. SPIE*, (2007) in press.

^dM. Quack, and J. Stohner, *J. Phys. Chem.*, 97(48), 12574 (1993).