

ROTATIONAL SPECTROSCOPY AND MOLECULAR PROPERTIES OF TITANIUM DIOXIDE, TiO₂

HOLGER S. P. MÜLLER, *I. Physikalisches Institut, Universität zu Köln, 50937 Köln; and Max-Planck-Institut für Radioastronomie, 53121 Bonn, Germany*; SANDRA BRÜNKEN, MICHAEL C. McCARTHY, PATRICK THADDEUS, *Harvard-Smithsonian Center for Astrophysics and Division of Engineering and Applied Sciences, Harvard University, Cambridge, MA 02138, USA*.

Titanium is one of the more abundant transition metals in space, and its dioxide is thought to be an important constituent of grains generated in the vicinity of oxygen-rich stars. Therefore, we have investigated the rotational spectrum of TiO₂ by laser-ablation molecular-beam Fourier transform microwave (LAMB-FTMW) spectroscopy between 7 and 42 GHz. Five isotopic species containing ^{46–50}Ti were studied in natural isotopic composition and two, ⁴⁸Ti¹⁶O¹⁸O and ⁴⁸Ti¹⁸O₂ were obtained through the use of ¹⁸O₂. Despite the comparatively large rotational constants and the absence of half of the rotational levels for all but the mixed ¹⁶O¹⁸O isotopic species because of spin-statistics, up to 13 rotational transitions with energies up to almost 40 cm⁻¹ could be recorded, permitting rotational and quartic centrifugal distortion parameters to be determined along with hyperfine structure parameters for the species containing ⁴⁷Ti or ⁴⁹Ti. The derived molecular structure and the harmonic force field as well as the hyperfine structure parameters will be compared with those of related molecules. Preliminary results of a search toward several oxygen-rich late-type stars carried out with the IRAM 30 m telescope will also be presented.