

THE ROTATIONAL SPECTRUM OF CN^-

C. A. GOTTLIEB, S. BRÜNKEN, M. C. McCARTHY, and P. THADDEUS, *Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, and Division of Engineering and Applied Sciences, Harvard University, Cambridge, MA 02138.*

CN^- is a unique molecular anion because neutral CN has one of the highest known electron affinities, is observed under a wide range of terrestrial conditions, and because this closed-shell anion is structurally and spectroscopically quite simple, making it amenable to both laboratory detection and high-level quantum chemical calculations. Although CN^- has been studied by photoelectron spectroscopy,^a no high-resolution spectra have been reported. Here we report the laboratory rotational spectrum of CN^- observed in a low pressure glow discharge through C_2N_2 and N_2 . Determination of the rotational and centrifugal distortion constants and the nitrogen quadrupole hyperfine coupling constant allow line frequencies to be calculated well into the far IR. CN^- is an excellent candidate for astronomical detection, because the CN radical is observed in many galactic molecular sources, and calculations^b indicate CN^- should be detectable in IRC+10216 — the carbon star where C_6H^- has recently been observed. The fairly high concentration of CN^- in our discharge implies that other molecular anions containing the nitrile group may be within reach.

^aS. E. Bradforth, E. H. Kim, D. W. Arnold, and D. M. Neumark, *J. Chem. Phys.* **98**, 800 (1993).

^bS. Petrie, *Mon. Not. R. Astron. Soc.* **281**, 137 (1996).