

## $\text{CH}_3^+$ : A NEW TOOL FOR STUDYING THE ENIGMA OF INTERSTELLAR $\text{CH}^+$ ?

BENJAMIN J. McCALL, *Departments of Chemistry and Astronomy, University of Illinois at Urbana-Champaign, Urbana, IL 61801*; THOMAS R. GEBALLE, *Gemini Observatory, Hilo, HI 96720*; TAKESHI OKA, *Department of Astronomy & Astrophysics and Department of Chemistry, University of Chicago, Chicago, IL 60637*.

Although  $\text{CH}^+$  was one of the very first interstellar molecules to be identified (in the early 1940s), its ubiquitous presence in the diffuse interstellar medium remains an enigma.  $\text{CH}^+$  is thought to be rapidly destroyed by the abundant species H,  $\text{H}_2$ , and  $e^-$ , but there is no known efficient formation mechanism at low temperatures. In steady state, one would therefore expect a very low abundance of  $\text{CH}^+$ , but yet strong  $\text{CH}^+$  lines are seen in nearly all diffuse cloud sightlines. Various solutions to this paradox have been proposed, and most of them invoke shocks (especially magnetohydrodynamic shocks) to drive the endothermic reaction  $\text{C}^+ + \text{H}_2 \rightarrow \text{CH}^+ + \text{H}$ .

Regardless of the mechanism by which it is produced,  $\text{CH}^+$  will undergo fast hydrogen abstraction reactions to form  $\text{CH}_2^+$  and then  $\text{CH}_3^+$ , if it is in an environment with a large fraction of the hydrogen in molecular form. Since  $\text{CH}_3^+$  reacts only very slowly with  $\text{H}_2$ , by radiative association to form  $\text{CH}_5^+$ , it is destroyed primarily by dissociative recombination with electrons. A simple steady state analysis suggests that  $\text{CH}_3^+$  may be over 10 times more abundant than  $\text{CH}^+$ , although this factor depends on the molecular fraction and the electron temperature.

In order to provide observational constraints on the environment in which  $\text{CH}^+$  is observed, we have searched for the infrared  $\nu_3$  band of  $\text{CH}_3^+$  at  $3.2 \mu\text{m}$  in diffuse cloud sources. We will present our upper limits on the equivalent width of the  $\text{CH}_3^+$  lines, discuss the inferred limits on the  $\text{CH}_3^+$  column density, and offer some speculations about the conditions in the regions where  $\text{CH}^+$  exists.