

## STUDY OF STATE DENSITY EFFECTS BY IMPULSIVE COLLISIONS OF HIGHLY VIBRATIONALLY EXCITED MOLECULES AND H<sub>2</sub>O

Q. LIU, D. K. HAVEY and A. S. MULLIN, *Department of Chemistry and Biochemistry, University of Maryland, College Park, MD.*

We investigated how donor molecules' state density affect impulsive collisions by studying collisional energy transfer of vibrationally excited pyridine-d<sub>5</sub> (donor) and H<sub>2</sub>O (bath), and the result was compared with experiments of pyridine:H<sub>2</sub>O, picoline:H<sub>2</sub>O and lutidine:H<sub>2</sub>O. Vibrationally excited pyridine-d<sub>5</sub> was prepared by absorbing pulsed 266nm UV light from fourth harmonic generator of a Nd:YAG laser. A tunable F-center laser with 0.0003 cm<sup>-1</sup> resolution was used to probe individual H<sub>2</sub>O (000 to 001) rotational states. The population of each H<sub>2</sub>O rotational state was measured by time dependent transient IR absorption and the rate of single collision between pyridine-d<sub>5</sub> and H<sub>2</sub>O was determined afterwards. The translational energy gain in H<sub>2</sub>O was obtained by Doppler-broadened linewidths for each rotational state and the rotational energy gain was measured by a population distribution of H<sub>2</sub>O rotational states with  $E_{rot}$  between 1000 and 2000 cm<sup>-1</sup>. The probability distributions for the large energy transfer of pyridine-d<sub>5</sub> and H<sub>2</sub>O was calculated and the result was compared with other donors and H<sub>2</sub>O.