

PARTIAL AND TOTAL COLUMN SFIT2 RETRIEVALS FROM DA8 AND PARIS-IR FOURIER TRANSFORM INFRARED SPECTRA RECORDED OVER CANADIAN ARCTIC IN THE SPRING OF 2004-5, INCLUDING COMPARISONS WITH ACE SATELLITE MEASUREMENTS

KEEYOON SUNG, KALEY A. WALKER, JAMES R. DRUMMOND, *Dept. of Physics, University of Toronto, 60 St. George St., Toronto, ON., M5S 1A7, CANADA*; DEJIAN FU, PETER F. BERNATH, *Dept. of Chemistry, University of Waterloo, 200 University Ave. W., ON., N2L 3G1, CANADA*; RICHARD L. MITTERMEIER, HANS FAST, *ARQX, Environment Canada, 4905 Dufferin Street, DOWNSVIEW, ON., M3H 5T4, CANADA*; and KIMBERLY STRONG, *Dept. of Physics, University of Toronto, 60 St. George St., Toronto, ON., M5S 1A7, CANADA*.

In the springs of 2004 and 2005, two Fourier transform spectrometers (FTSs) were operated at Eureka, Nunavut, Canada (80.05°N, 86.42°W) as part of Arctic validation campaigns for the Atmospheric Chemistry Experiment (ACE). Using the SFIT2 profile retrieval program, we have retrieved total and partial column densities from the resulting solar absorption spectra for seven stratospheric species (O<sub>3</sub>, HF, HCl, NO, NO<sub>2</sub>, HNO<sub>3</sub>, ClONO<sub>2</sub>) and two tropospheric species (CH<sub>4</sub>, N<sub>2</sub>O) out of the 14 ACE baseline molecules. A complete set of a priori trace gas volume mixing ratio (VMR) profiles and their covariances has been constructed by using zonal mean profiles for latitudes higher than 65°N from HALOE (v.19, 1992-2005), SPARC 2000, MIPAS (v.3, 2002-2004), and the Eureka ozonesonde archive (1993-2005), and relative humidity from daily radiosonde measurements. Through investigation of column averaging kernels and degrees of freedom for signal, the ground-based retrievals have been well characterized for total and partial columns. For ozone and HCl, total error budgets for the partial and total column retrievals have been made. The DA8 measurements show good agreement with those from PARIS-IR, ACE-FTS, and ozonesondes, except those of NO<sub>2</sub>. In addition, the DA8 measurements fall on the time series of PARIS-IR measurements for a given day for the total columns of ozone, HCl, and HNO<sub>3</sub>. Differences in the chemical composition between the two consecutive years have also been analyzed using column ratios of O<sub>3</sub>/HF, HNO<sub>3</sub>/HF, HCl/HF, and ClONO<sub>2</sub>/HF from the three Fourier transform spectrometers.