

NEAR-INFRARED DOUBLE RESONANCE SPECTRUM OF THE CH₂ $\tilde{b} \leftarrow \tilde{a}$ ORIGIN BAND

ZHONG WANG, *Department of Chemistry, The Stony Brook University, Stony Brook, New York 11794*;
BOR-CHEN CHANG, *Department of Chemistry, National Central University, Jhongli 32001, Taiwan*;
TREVOR J. SEARS, *Department of Chemistry, Brookhaven National Laboratory, Upton, New York 11973*
and *Department of Chemistry, The Stony Brook University, Stony Brook, New York 11794*.

Since Herzberg and Johns reported the first observation of the CH₂ $\tilde{b} \leftarrow \tilde{a}$ singlet-singlet transition, there have been numerous experimental and theoretical investigations of this system. Although the spectrum of the CH₂ $\tilde{b} \leftarrow \tilde{a}$ transition has now been studied from the ultraviolet to the near-infrared (NIR), the origin band has not previously been observed due to a poor Franck-Condon factor as well as the lack of a convenient high resolution light source at approximately 1.2 μm . Here we report the successful observation of the origin band of the CH₂ $\tilde{b} \leftarrow \tilde{a}$ system using a NIR-NIR double resonance technique. Based upon our results, effective rotational constants for the \tilde{b} (0,0,0) level can be determined. The level positions also provide information that can be used to improve the potential energy surface for the \tilde{b} state. Singlet CH₂ is one of very few polyatomic molecules for which continuous high resolution data exist for rovibronic levels from the zero-point level of the \tilde{a} state through to predissociation.

Acknowledgments: This work at Brookhaven National Laboratory was carried out under Contract No.DE-AC02-98CH10886 with the U.S. Department of Energy, and BCC is supported by National Science Council, Taiwan.