

CONSTRAINING THE FLUX OF LOW-ENERGY COSMIC RAYS ACCELERATED BY THE SUPERNOVA REMNANT IC 443

NICK INDRIOLO, *Department of Astronomy, University of Illinois at Urbana-Champaign, Urbana, IL 61801*; GEOFFREY A. BLAKE, *Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA 91125*; MIWA GOTO, *Max Planck Institute for Astronomy, Königstuhl 17, Heidelberg D-69117, Germany*; TOMONORI USUDA, *Subaru Telescope, Hilo, HI 96720*; THOMAS R. GEBALLE, *Gemini Observatory, Hilo, HI 96720*; TAKESHI OKA, *Department of Astronomy & Astrophysics and Department of Chemistry, University of Chicago, Chicago, IL 60637*; BENJAMIN J. McCALL, *Departments of Chemistry and Astronomy, University of Illinois at Urbana-Champaign, Urbana, IL 61801*.

It has long been theorized that supernova remnants (SNR) accelerate the majority of Galactic cosmic rays. Observations in the  $\gamma$ -ray, X-ray, and radio regimes support this theory, at least for cosmic rays with energies above a few GeV. However, there is no direct evidence that SNRs accelerate cosmic rays in the MeV–GeV range. These low-energy cosmic rays are of great importance, as they are the primary means by which  $\text{H}_2$  is ionized in the interstellar medium. Collisions between  $\text{H}_2^+$  and  $\text{H}_2$  will rapidly form  $\text{H}_3^+$ , a molecule which can then be observed to infer the ionization rate of  $\text{H}_2$ . Using the Subaru and Keck telescopes, we have searched for  $\text{H}_3^+$  absorption in sight lines which probe molecular material known to be interacting with the SNR IC 443. By computing the ionization rate of  $\text{H}_2$  in these sight lines, we constrain the flux of low-energy cosmic rays generated by this particular supernova remnant.