

## LONG-WAVELENGTH PHOTOCHEMISTRY OF MATRIX-ISOLATED BIACETYL

NATHAN G. KUCHMAS and C. A. BAUMANN, *Department of Chemistry, The University of Scranton, Scranton, PA 18510-4626.*

Visible irradiation ( $520 \text{ nm} < \lambda < 485 \text{ nm}$ ) of matrix-isolated biacetyl ( $\text{C}_4\text{H}_6\text{O}_2$ ) results in the formation of a complex of *cis*-methylhydroxycarbene ( $\text{CH}_3\text{COH}$ ) and ketene ( $\text{CH}_2\text{CO}$ ), as well as other products. The wavelengths used in this study are longer than those necessary for the  $S_0$ - $S_1$  ( $\tilde{X}^1A_g$  -  $\tilde{A}^1A_u$ ) transition, indicating that the photolysis is the result of a multiphoton process. One such process is sequential, where the spin-forbidden  $S_0$ - $T_1$  ( $\tilde{X}^1A_g$  -  $\tilde{a}^3A_u$ ) transition is followed by the allowed  $T_1$ - $T_2$  ( $\tilde{a}^3A_u$  -  $\tilde{b}^3B_g$ ) transition, with photoproducts emerging from the higher triplet (or following a radiationless transition to another state). A simultaneous two-photon process is also possible through the symmetry-forbidden  $S_0$ - $S_2$  ( $\tilde{X}^1A_g$  -  $\tilde{B}^1B_g$ ) transition. Photoproduct formation as a function of irradiation flux and wavelength was used to sort out the relative contributions of each of these processes.