

## THE PHOTOELECTRON ANGULAR DISTRIBUTION AS A PROBE OF ENERGETICALLY INDISTINGUISABLE CHANNELS IN PHOTODETACHMENT

MATTHEW VAN DUZOR, FOSTER MBAIWA, JIE WIE and RICHARD MABBS, *Department of Chemistry, Washington University, One Brookings Dr., Campus Box 1134 Saint Louis, Missouri 63130, USA.*

Photoelectron imaging measurements of  $\text{I}^- \cdot \text{CH}_3\text{CN}$  are presented from the lowest energy  $\text{I}(^2\text{P}_{3/2}) \cdot \text{CH}_3\text{CN}$  channel threshold to 0.3 eV above the  $\text{I}(^2\text{P}_{1/2}) \cdot \text{CH}_3\text{CN}$  threshold. Excitation of the cluster just below the latter threshold leads to competition between direct detachment and the production of a dipole-bound state  $[\text{I}(^2\text{P}_{1/2}) \cdot \text{CH}_3\text{CN}]^-$ . Subsequent relaxation of the I moiety ( $^2\text{P}_{1/2} \rightarrow ^2\text{P}_{3/2}$ ) occurs via autodetachment of the dipole bound electron. However, the autodetached electrons are energetically equivalent to those directly detached via the  $^2\text{P}_{3/2}$  channel. While the photoelectron spectrum is insensitive to this phenomenon, the kinetic energy dependence of the photoelectron angular distribution (PAD) of the  $^2\text{P}_{3/2}$  channel reflects these dynamics. Dramatic changes are observed within a 100 meV window of the  $\text{I}(^2\text{P}_{1/2}) \cdot \text{CH}_3\text{CN}$  threshold. Although previously unreported, further results are presented that show sharp changes in the PAD are relatively common in  $\text{I}^- \cdot \text{X}$  cluster anions ( $\text{X}=\text{polar molecule}$ ).