

AIR-BROADENED LINE PARAMETERS FOR THE $2\leftarrow 0$ BANDS OF $^{13}\text{C}^{16}\text{O}$ AND $^{12}\text{C}^{18}\text{O}$ AT $2.3\ \mu\text{m}$

V. MALATHY DEVI, D. CHRIS BENNER, *The College of William and Mary, Williamsburg, VA 23187*; M. A. H. SMITH, *Science Directorate, NASA Langley Research Center, Hampton, VA 23681*; A. W. MANTZ, *Dept. of Physics, Astronomy and Geophysics, Connecticut College, New London, CT 06320*; K. SUNG and L. R. BROWN, *Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109*.

Air-broadened line shape parameters were determined for the first time in the $2\leftarrow 0$ bands of $^{13}\text{C}^{16}\text{O}$ near $4166.8\ \text{cm}^{-1}$ and $^{12}\text{C}^{18}\text{O}$ near $4159.0\ \text{cm}^{-1}$. Spectra were recorded at $0.005\ \text{cm}^{-1}$ resolution using a coolable absorption cell^a in the sample compartment of the Bruker IFS 125HR Fourier transform spectrometer at Jet Propulsion Laboratory. Gas temperatures and pressures ranged from 150 to 298 K and 20 to 700 Torr, respectively. Line parameters were determined by broad-band multispectrum least-squares fitting^b of the $4000\text{--}4360\ \text{cm}^{-1}$ region in 16 spectra simultaneously; each set included 4 isotope-enriched pure sample scans and 12 air+CO samples (^{13}CO or C^{18}O , as appropriate). The air-broadened parameters measured were Lorentz half-width coefficients, their temperature dependence exponents; pressure-induced shift coefficients, their temperature dependences; and off-diagonal relaxation matrix elements. Speed dependence parameters were included to minimize the fit residuals. For both isotopologues the individual line positions and intensities were constrained to their theoretical relationships in order to obtain the rovibrational (G, B, D, and H) and band intensity parameters, including Herman-Wallis coefficients^c. The results for $^{13}\text{C}^{16}\text{O}$ and $^{12}\text{C}^{18}\text{O}$ are compared with those for the $^{12}\text{C}^{16}\text{O}\ 2\leftarrow 0$ band^d and discussed.^e

^aK. Sung, A. W. Mantz, M. A. H. Smith, et al., *JMS* **262** (2010) 122-134.

^bD. C. Benner, C. P. Rinsland, V. Malathy Devi, M. A. H. Smith and D. A. Atkins, *JQSRT* **53** (1995) 705-721.

^cV. Malathy Devi, D. C. Benner, L. R. Brown, C. E. Miller and R. A. Toth, *JMS* **242** (2007) 90-117.

^dV. Malathy Devi, D. C. Benner, M. A. H. Smith, et al., *JQSRT* (2012) in press.

^eResearch described in this paper was performed at Connecticut College, the College of William and Mary, NASA Langley Research Center and the Jet Propulsion Laboratory, California Institute of Technology, under contracts and cooperative agreements with the National Aeronautics and Space Administration.