HELIUM NANODROPLET ISOLATION OF IONIC LIQUID VAPOR: INRARED LASER SPECTROSCOPY OF [EMIM][Tf_2N]

STEVEN D. FLYNN, GARY E.DOUBERLY, *Department of Chemistry, University of Georgia, Athens, Georgia, USA 30602.*

The Infrared spectrum of the vapor produced upon thermal vaporization of the [emim][Tf_2N][a] ionic liquid has been obtained using the helium nanodroplet isolation method. Despite the low vapor pressure of [emim][Tf_2N], sufficient gas phase densities are produced, allowing for efficient helium nanodroplet pick-up. The mass spectrum of the emim[Tf_2N] doped droplet beam shows signatures[b] that have been attributed in gas phase measurements to the presence of isolated, intact [emim][Tf_2N] ion-pairs. Furthermore, the mass spectrometry results indicate that emim[Tf_2N] does not undergo thermal decomposition at 410 K. Comparisons are made between the experimental measurements and *ab initio* calculations (mp2/6-311++g(d,p)) of the CH stretch vibrational bands and permanent electric dipole moments for several [emim][Tf_2N] low energy isomers. The helium nanodroplet infrared spectrum of this species provides rather definitive support to the previously suggested vaporization mechanism of ionic liquids.[c]

[a] [emim][Tf_2N] is defined as 1-ethyl-3-methylimidazolium[bis(trifluoromethylsulfonyl)imide]

[b] Armstrong, J.P.; Hurst, C.; Jones, R. G.; Licence, P.; Lovelock, K. R. J.; Satterley, C. J.; Villar-Garcia, I. J. *Physical Chemistry Chemical Physics* **2007**, *9*, 982.

Strasser, D.; Goulay, F.; Belau, L.; Kostko, O.; Koh, C.; Chambreau, S. D.; Vaghjiani, G. L.; Ahmed, M.; Leone, S. R. textitJournal of Physical Chemistry A **2010**, textit114, 879.

[c] Strasser, D.; Goulay, F.; Kelkar, M. S.; Maginn, E. J.; Leone, S. R. *Journal of Physical Chemistry A* **2007**, *111*, 3191.

Chambreau, S. D.; Vaghjiani, G. L.; To, A.; Koh, C.; Strasser, D.; Kostko, O.; Leone, S. R. *Journal of Physical Chemistry B* **2010**, 114, 1361.

Maginn, E. J.; Kelkar, M. S. *Journal of Physical Chemistry B* **2007**, *111*, 9424.