

## Kronecker Delta Function $\delta_{ij}$ and Levi-Civita (Epsilon) Symbol $\varepsilon_{ijk}$

### 1. Definitions

$$\delta_{ij} = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{otherwise} \end{cases} \quad \varepsilon_{ijk} = \begin{cases} +1 & \text{if } \{ijk\} = 123, 312, \text{ or } 231 \\ -1 & \text{if } \{ijk\} = 213, 321, \text{ or } 132 \\ 0 & \text{all other cases (i.e., any two equal)} \end{cases}$$

- So, for example,  $\varepsilon_{112} = \varepsilon_{313} = \varepsilon_{222} = 0$ .
- The +1 (or *even*) permutations are related by rotating the numbers around; think of starting with 123 and moving (in your mind) the 3 to the front of the line, to get 312. Do it again with the 2 and you get 231. The -1 (or *odd*) permutations starting with 213 are related to each other the same way; they are related to 123 by interchanging just two of the numbers (e.g., switch the 1 and 3 to get 321).

### 2. Applying $\delta_{ij}$ and $\varepsilon_{ijk}$ to Vectors in Cartesian coordinates

- Instead of using  $x$ ,  $y$ , and  $z$  to label the components of a vector, we use 1, 2, 3.
- Then the letters  $i$ ,  $j$ ,  $k$ , ... can be used as (dummy) summation variables, running from 1 to 3. (We could use any other letters, like  $a$ ,  $b$ , ...; it is merely a convention.)
- Don't confuse the use of the dummy summation variables  $i$ ,  $j$ ,  $k$ , each of which can be 1, 2, or 3, with the unit vectors  $\hat{\mathbf{i}}$ ,  $\hat{\mathbf{j}}$ ,  $\hat{\mathbf{k}}$ . These are two independent notations!
- The dot product of two vectors  $\mathbf{A} \cdot \mathbf{B} = A_x B_x + A_y B_y + A_z B_z$  in this notation is

$$\mathbf{A} \cdot \mathbf{B} = A_1 B_1 + A_2 B_2 + A_3 B_3 = \sum_{i=1}^3 A_i B_i = \sum_{i=1}^3 \sum_{j=1}^3 A_i B_j \delta_{ij} = \sum_{i=1}^3 \sum_{j=1}^3 A_i \delta_{ij} B_j .$$

Note that there are nine terms in the final sums, but only three terms are non-zero, and that the final sum is in the form of a matrix multiplication.

- The  $i^{\text{th}}$  component of the cross product of two vectors  $\mathbf{A} \times \mathbf{B}$  becomes

$$(\mathbf{A} \times \mathbf{B})_i = \sum_{j=1}^3 \sum_{k=1}^3 \varepsilon_{ijk} A_j B_k .$$

Again, there are nine terms in the sum, but this time only two of them are non-zero. Note also that this expression summarizes three equations, namely for  $i = 1, 2, 3$ .

### 3. Einstein Summation Convention

- We might notice that the summations in the expressions for  $\mathbf{A} \cdot \mathbf{B}$  and  $\mathbf{A} \times \mathbf{B}$  are redundant, because they only appear when an index like  $i$  or  $j$  appears twice on one side of an equation. So we can omit them. Thus

$$\sum_{i=1}^3 \sum_{j=1}^3 A_i B_j \delta_{ij} \longrightarrow A_i B_j \delta_{ij} = A_i B_i \quad \text{and} \quad \sum_{j=1}^3 \sum_{k=1}^3 \varepsilon_{ijk} A_j B_k \longrightarrow \varepsilon_{ijk} A_j B_k .$$

- Rules: If an index appears (exactly) twice, then it is summed over and appears only on one side of an equation. A single index (called a *free index*) appears once on *each side* of the equation. So

$$\text{Valid: } A_i = A_j \delta_{ij}, \quad B_k = \varepsilon_{ikl} A_i C_l \qquad \text{Invalid: } A_i = B_i C_i, \quad A_i = \varepsilon_{ijk} B_i C_j .$$

- When you have a Kronecker delta  $\delta_{ij}$  and one of the indices is repeated (say  $i$ ), then you simplify by replacing the other  $i$  index on that side of the equation by  $j$  and removing the  $\delta_{ij}$ . For example:

$$A_j \delta_{ij} = A_i, \quad B_{ij} C_{jk} \delta_{ik} = B_{kj} C_{jk} = B_{ij} C_{ji}$$

Note that in the second case we had two choices of how to simplify the equation; use either one!

- The triple or box product  $\mathbf{A} \cdot (\mathbf{B} \times \mathbf{C})$  can be written

$$\mathbf{A} \cdot (\mathbf{B} \times \mathbf{C}) = \varepsilon_{ijk} A_i B_j C_k = \varepsilon_{kij} A_i B_j C_k = \varepsilon_{kij} C_k A_i B_j = \mathbf{C} \cdot (\mathbf{A} \times \mathbf{B}) ,$$

where we've used the properties of  $\varepsilon_{ijk}$  in the middle equations to prove a relation among triple products with the vectors in a different order.

- A very useful identity (if the repeated index is not first in both  $\varepsilon$ 's, permute until it is):

$$\varepsilon_{ijk} \varepsilon_{ilm} = \delta_{jl} \delta_{km} - \delta_{jm} \delta_{kl} .$$

#### 4. Example: Proving a Vector Identity

- We'll write the  $i^{\text{th}}$  Cartesian component of the gradient operator  $\nabla$  as  $\partial_i$  (cf.  $\frac{\partial}{\partial x_i}$ ).
- Let's simplify  $\nabla \times (\nabla \times \mathbf{A}(\mathbf{x}))$ . We start by considering the  $i^{\text{th}}$  component and then we use our expression for the cross product (working from the outside in):

$$(\nabla \times (\nabla \times \mathbf{A}))_i = \varepsilon_{ijk} \partial_j (\nabla \times \mathbf{A})_k .$$

Next we replace the remaining cross product, making sure to introduce new dummy summation variables  $l$  and  $m$ :

$$(\nabla \times (\nabla \times \mathbf{A}))_i = \varepsilon_{ijk} \partial_j \varepsilon_{klm} \partial_l A_m = \varepsilon_{kij} \varepsilon_{klm} \partial_j \partial_l A_m .$$

(The partial derivatives act only on the components of  $\mathbf{A}$ , so we can pull out the  $\varepsilon$ 's.) We rotated the indices in one of the  $\varepsilon$ 's in the last step so that we can now directly apply our very useful identity (and simplify):

$$(\nabla \times (\nabla \times \mathbf{A}))_i = (\delta_{il} \delta_{jm} - \delta_{im} \delta_{lj}) \partial_j \partial_l A_m = \partial_m \partial_i A_m - \partial_l \partial_l A_i = \partial_i (\partial_m A_m) - (\partial_l \partial_l A)_i$$

or, finally,

$$\nabla \times (\nabla \times \mathbf{A}) = \nabla (\nabla \cdot \mathbf{A}) - \nabla^2 \mathbf{A} .$$