



# Social distancing decreases an individual's likelihood of contracting COVID-19

Russell H. Fazio<sup>a,1</sup>, Benjamin C. Ruisch<sup>b</sup>, Courtney A. Moore<sup>a</sup>, Javier A. Granados Samayoa<sup>a</sup>, Shelby T. Boggs<sup>a</sup>, and Jesse T. Ladanyi<sup>a</sup>

<sup>a</sup>Department of Psychology, The Ohio State University, Columbus, OH 43210; and <sup>b</sup>Institute of Psychology, Leiden University, 2300 RB Leiden, The Netherlands

Edited by Douglas S. Massey, Princeton University, Princeton, NJ, and approved January 8, 2021 (received for review November 5, 2020)

Past research has established the value of social distancing as a means of deterring the spread of COVID-19 largely by examining aggregate level data. Locales in which efforts were undertaken to encourage distancing experienced reductions in their rate of transmission. However, these aggregate results tell us little about the effectiveness of social distancing at the level of the individual, which is the question addressed by the current research. Four months after participating in a study assessing their social distancing behavior, 2,120 participants indicated whether they had contracted COVID-19. Importantly, the assessment of social distancing involved not only a self-report measure of how strictly participants had followed social distancing recommendations but also a series of virtual behavior measures of social distancing. These simulations presented participants with graphical depictions mirroring specific real-world scenarios, asking them to position themselves in relation to others in the scene. Individuals' social distancing behavior, particularly as assessed by the virtual behavior measure, predicted whether they contracted COVID-19 during the intervening 4 mo. This was true when considering only participants who reported having tested positively for the virus and when considering additional participants who, although untested, believed that they had contracted the virus. The findings offer a unique form of additional evidence as to why individuals should practice social distancing. What the individual does matters, not only for the health of the collective, but also for the specific individual.

contracting COVID-19 | social distancing | virtual behavior

Since its emergence in late 2019, the COVID-19 pandemic has had a devastating impact on nations around the world. The negative effects are evident with respect to the shocking number of people who have been infected with this novel coronavirus and the sheer number of deaths, as well as the impact upon nations' economies. Given the lack of a widely available vaccine, minimizing the spread of COVID-19 has required that people change their behavior. Government officials and public health experts worldwide have urged people to engage in various preventive actions: Wash your hands frequently for at least 20 s, use hand sanitizer when handwashing is not feasible, avoid touching your face, disinfect surfaces that you touch, and wear masks. Perhaps most importantly, they have pleaded that their citizens engage in social distancing—avoid close contact with other people. The mantra “six feet” has been repeated regularly and now appears on signs and floor decals throughout society.

Although knowledge regarding SARS-CoV-2 transmission is constantly expanding, much has been learned about the dynamics responsible for its spread (1). Considerable evidence stemming from the investigation of such viruses as SARS, MERS, and SARS-CoV-2 highlights that physical distancing reduces their aerial transmission (e.g., ref. 2). Mathematical models of the spread of COVID-19 in particular (e.g., refs. 3–5) have illustrated the efficacy of social distancing as a means of mitigating its spread. Moreover, recent epidemiological evidence also documents the effectiveness of social distancing at the societal level. For example, a recent analysis involving both the US

states and 134 nations revealed that the implementation of social distancing policies was associated with a reduction in COVID-19 spread rates within those locales (6). Similarly, an examination of the daily growth rate in COVID-19 cases across US counties during March and April 2020 found that the imposition of shelter-in-place orders and closures of restaurants, bars, and entertainment-related businesses slowed the spread of COVID-19 (7). Other analyses have found that greater social distancing at the county level, as estimated by smartphone GPS data, is associated with reduced spread of COVID-19 and fewer deaths (8).

Yet what remains unknown is the extent to which an individual's social distancing behavior predicts their personal likelihood of contracting COVID-19. This is a critical theoretical and empirical gap, particularly given that the demonstrated effectiveness of social distancing at the aggregate level does not necessarily mean that individual differences in social distancing behavior will predict whether a given individual contracts the disease. For example, it may be that a “critical mass” of social distancing behavior in a given locale is necessary for any individual-level benefits to emerge. Until that critical level is reached, it is possible that individual differences may matter little for disease transmission.

Furthermore, beyond the question of whether individual differences in social distancing behavior predict disease

## Significance

Previous work establishes the effectiveness of social distancing for reducing COVID-19 transmission at the aggregate level: Locales in which restrictions were imposed experienced a reduction in spread of the virus. However, we know little about the effectiveness of social distancing at the level of the individual. Do individuals who engage in social distancing reduce their personal likelihood of contracting COVID-19? Or are these effects only evident in the aggregate? A longitudinal investigation involving 2,120 online participants demonstrated that individual differences matter. Participants who exhibited greater social distancing on a virtual behavior measure—simulations presenting graphical depictions of specific real-world scenarios, asking them to position themselves relative to others in the scene—were less likely to contract COVID-19 subsequently.

Author contributions: R.H.F., B.C.R., C.A.M., J.A.G.S., S.T.B., and J.T.L. designed research; R.H.F., B.C.R., C.A.M., J.A.G.S., S.T.B., and J.T.L. performed research; B.C.R. and C.A.M. contributed new reagents/analytic tools; R.H.F. analyzed data; and R.H.F., B.C.R., C.A.M., J.A.G.S., S.T.B., and J.T.L. wrote the paper.

The authors declare no competing interest.

This article is a PNAS Direct Submission.

Published under the PNAS license.

<sup>1</sup>To whom correspondence may be addressed. Email: fazio.11@osu.edu.

This article contains supporting information online at <https://www.pnas.org/lookup/suppl/doi:10.1073/pnas.2023131118/-DCSupplemental>.

Published February 4, 2021.

transmission, there is also the critical question of the degree to which these individual differences matter. In other words, what is the size of this effect? Do those who practice social distancing enjoy a notably reduced likelihood of contracting the disease? Or are these effects relatively modest in size? Aggregate data offer little insight into these questions. Indeed, aggregation over a large number of instances may be necessary to observe what might be small effects at the level of the individual.

To address this theoretical and empirical gap, the current research pursues a different approach to testing the effectiveness of social distancing—one that focuses not on aggregates but on the level of the individual. Are individuals who engage in social distancing behaviors less likely to contract COVID-19? If so, to what degree? To examine this question, we assessed participants' (US residents) social distancing behavior at the height of the first wave of the pandemic and then 4 mo later asked them to report whether they had contracted the virus.

The assessment of the extent to which individuals practiced social distancing was accomplished by two very different measurement techniques. The standard approach generally employed in social science research, and certainly true of hundreds of recent studies concerning the pandemic (9), is simply to prepare survey questions asking people to report the extent to which they personally practice social distancing. We did precisely the same in the current research.

However, what makes our research all the more unique is that we supplemented the self-report measure with a more innovative, behaviorally oriented approach to the assessment of social distancing. We simulated social distancing behavior by presenting participants with graphical depictions that mirrored different real-world scenarios and asking them to position themselves as they personally would in relation to others in the scene. These virtual social distancing scenarios required a concrete, "in-the-moment" behavioral decision, which could vary in the degree to which participants did or did not distance themselves from others. For example, in one scenario participants chose whether to cross a park via a circuitous but isolated path versus a more direct but crowded route. In another, they were presented an aerial image of a crowded beach and asked to click on the spot where they personally would lay down their towel. Yet another presented an interactive image of two people approaching each other in a crosswalk for which participants were asked to move a slider that shifted the walkers from the center of the crosswalk to the distance that they personally would leave between themselves and the other individual. The 10 behavioral scenarios can be viewed at our demonstration website, <http://psychvault.org/social-distancing/>.

We included this virtual behavior distancing measure out of concern for the potential problems that have been documented to be associated with self-reports of behavior. Individuals may overreport their social distancing to convey a socially desirable impression to others and themselves (10–13). Moreover, self-reports may be all the more problematic to the extent that they rely on retrospective memory regarding past behavior (14, 15). By simulating concrete real-world situations that require an immediate decision, the graphical scenarios offer a means, in addition to a self-report, of indexing the extent to which individuals behave in a manner that accords with the principle of social distancing.

The present study examines whether both the self-report measure and the virtual behavior measure of social distancing prospectively predict, 4 mo later, whether an individual contracted COVID-19. The data are available on the Open Science Framework: <https://osf.io/x79ak/>.

## Method

We recruited our participant samples from Amazon's Mechanical Turk participants (16). Although not representative of the US population, MTurk

samples tend to be more demographically, politically, and geographically diverse than the samples typically used in psychological research (17). They also perform similarly to non-MTurk samples across many tasks and measures (18, 19), including surveys on political attitudes (20). Most importantly, however, our aim is not to make claims regarding the absolute frequency of social distancing behaviors or COVID-19 illness. Instead, we seek to understand whether the extent to which individuals engage in social distancing, as measured at one point in time, relates to their having contracted the coronavirus, as reported at a later point in time. Given this covariation aim, MTurk participants offer the opportunity for an appropriate test of the hypothesis that social distancing behavior is prospectively predictive of individuals' reports of having become ill with COVID-19. Ohio State University's Institutional Review Board approved the study procedures (IRB: 2020B0129).

**Participants.** The sample consisted of MTurkers who had participated in one of two earlier studies conducted in spring 2020. All participants who had granted permission to recontact them were invited to complete a brief survey approximately 4 mo after their initial study for a payment of \$1. A total of 2,120 individuals, all US residents, did so (1,031 women, 1,074 men, 15 no response;  $M_{age} = 40.39$ ,  $SD_{age} = 15.34$ ). Study 1 had been completed on May 7–8 ( $n = 1,281$ ) and study 2 on June 9 ( $n = 839$ ).\*

**Measures.** Each of the initial studies concerned reactions to the pandemic and relevant individual difference measures. The variables of interest for the current research concern social distancing behavior and two variables identified on an a priori basis as likely predictors of contracting COVID-19: preexisting conditions and working outside the home. The former was assessed by asking participants to consider their "personal health prior to the outbreak of the COVID-19 virus" and to then indicate whether they would have described themselves "as having pre-existing medical conditions that left you more vulnerable to the virus than the average person" by selecting one of five response options: Definitely not/Probably not/Maybe/Probably yes/Definitely yes (coded as 1–5, respectively). Participants were also asked about their employment status. Our interest was in whether they selected the option "My job requires that I leave my home" (coded as 1) as opposed to other options (coded as 0) concerning working from home, having lost their job due to COVID-19, or having been unemployed before the pandemic began. Our reasoning was that individuals whose job required leaving home would be at greater risk for contracting the virus.

**Self-reported social distancing behavior.** Participants also were asked to report on the extent to which they were practicing social distancing. For study 1, which was conducted when only a few states had even begun the process of relaxing stay-at-home orders and reopening their economies, a single relevant question was employed. Participants were asked "Generally speaking, how strictly have you personally been following the 'social distancing' recommendations?" and responded using a seven-point scale ranging from "not at all" to "very strictly." By the time of study 2, most states had reopened, at least partially. Hence, we included two relevant questions involving the same seven-point scale. One inquired specifically about behavior during the lockdown: "Up to a few weeks ago, most people were under 'shelter-in-place' orders (for example, staying home except for absolute necessities, having no contact with people outside your household, etc.). During that time, how strictly did you follow these recommendations?" A second question asked specifically about maintaining safe distances: "Generally speaking, how strictly have you personally been following the 'social distancing' recommendations of the government and CDC to maintain a distance of six feet or more from others?" Responses to these two questions correlated highly ( $r = 0.73$ ) and, hence, were averaged to form a single composite of self-reported social distancing for the participants in this later study.

**Virtual social distancing behavior.** Following the provision of informed consent, participants in each of the initial studies read that the researchers were "interested in people's behavior in common everyday situations, and how this behavior may have changed in light of the current COVID-19/coronavirus pandemic." They would "view images depicting various social situations" and would "be asked to indicate how **you personally** (emphasis in original) would behave." What followed was the presentation of graphical scenarios that asked participants to position themselves relative to others in a specific real-world situation. A total of 10 behavioral scenarios, all similar to the examples described earlier, were presented. The scenarios are described in *SI Appendix* and, as noted earlier, can be viewed at our demonstration

\*See Fazio et al. (21) for a detailed report of study 1.

**Table 1. Binary logistic regression predicting COVID-19 status\***

Variable*	B	SE	Wald	p	Odds ratio
Preexisting conditions <sup>†</sup>	0.395	0.072	30.326	0.000	1.484
Leave home for work*	0.350	0.183	3.684	0.055	1.420
Self-reported behavior <sup>†</sup>	-0.031	0.085	0.131	0.717	0.970
Virtual behavior <sup>†</sup>	-0.202	0.087	5.459	0.019	0.817
Constant	-2.286	0.091	624.055	0.000	NA

NA, not applicable.

\*Coded as 0 = No; 1 = Yes.

<sup>†</sup>Standardized.

website, <http://psychvault.org/social-distancing/>. After standardizing scores from each scenario within each study, we computed the average as our index of virtual social distancing behavior ( $\alpha = 0.83$ ).

**Follow-up survey.** Four months after the initial study, the participants completed a brief survey about whether they had or had not contracted the coronavirus. They first were asked whether they had been tested for COVID-19. If so, they indicated whether the test showed that they had COVID-19. If they had not been tested, they were asked "Even though you may not have been tested, do you believe that you have ever had COVID-19/the coronavirus?" to which they responded yes or no. (These very same questions were included in the initial survey that participants completed.) Participants who reported either testing positively or believing they had COVID-19 were then asked to select from a list of possibilities how they thought they might have contracted the virus.

## Results

**Descriptive Data.** Of the 2,120 participants, 516 (24.3%) reported having been tested for COVID-19, with 116 (5.5% of the total sample) reporting a positive result. Another 232 participants (10.9% of the total sample) reported that, although they had not been tested, they believed that they had indeed contracted the coronavirus. In sum, at the time of the follow-up survey, a total of 348 participants (16.42%) reported having experienced COVID-19 illness. When asked to indicate how they might have contracted the virus, 23.3% of these 348 participants selected "from a member of my household," 21.9% selected "at work," and 17.6% selected "at an indoor setting (a restaurant, bar, store, church or other indoor venue)." The other possibilities that were selected with some frequency were: "from a friend or relative who visited me at home" (11.2%), "at an outdoor setting (a public event, park, restaurant patio, outdoor party, etc.," (6.9%), and "while traveling on a trip out of town" (4.9%).

**Predicting COVID-19 illness at Follow-Up.** Our major interest is to examine whether social distancing behavior prospectively predicts subsequent illness. Hence, we need to exclude from the analyses any participants who reported having had COVID-19 at the time of their initial study. Of the 2,120 follow-up participants, 235 reported a positive test result at time 1 or the untested belief that they had COVID-19, resulting in a sample of 1,885 participants for our first set of analyses. Of these, 199 subsequently reported a positive test result or the belief that they had contracted COVID-19 (coded as 1) and 1,686 who reported either a negative test result or the belief that they had not contracted the virus<sup>†</sup> (coded as 0).

A binary logistic regression was conducted examining the dichotomous COVID-19 status variable at follow-up as a function of the predictor variables of interest. To ease interpretation, all continuous predictor variables were standardized. We first considered the two a priori risk variables of preexisting conditions and leaving home for work (coded 0/1 for no/yes). As anticipated, each accounted for significant unique variance

(preexisting conditions:  $B = 0.36$ , Wald = 27.03,  $P < 0.001$ , odds ratio = 1.44; leave home for work:  $B = 0.41$ , Wald = 5.48,  $P = 0.019$ , odds ratio = 1.51). We next considered the virtual behavioral and self-report measures of social distancing, which correlated at 0.50,  $P < 0.001$ . When the behavioral distancing score was added as the next variable in the equation, it too revealed a statistically significant effect ( $B = -0.22$ , Wald = 8.25,  $P = 0.004$  odds ratio = 0.80); less social distancing was associated with a greater likelihood of contracting COVID-19. A similar effect, albeit weaker, was observed when the self-report measure of social distancing was added as the third variable in the equation, instead of the virtual behavior measure ( $B = -0.13$ , Wald = 3.04,  $P = 0.081$ , odds ratio = 0.88). Table 1 presents the relevant statistics when all four variables are considered simultaneously. The virtual behavior measure accounted for significant unique variance, whereas the self-report measure did not.

Parallel analyses were conducted predicting whether participants reported having tested positively for COVID-19 at follow-up. In other words, the focus here is solely on positive test results as an indicator of COVID-19 illness, in contrast to the earlier approach in which participants who had not been tested but believed they had contracted the virus also were considered within the COVID-19 classification. These analyses excluded only participants who had reported a positive test result in their initial survey. They compare the 85 participants who subsequently tested positively to the 1,993 who reported either a negative test or not having been tested at all.<sup>‡</sup> As before, all continuous predictor variables were standardized, and we first considered the two a priori risk variables of preexisting conditions and leaving home for work. The former accounted for substantial unique variance ( $B = 0.69$ , Wald = 44.14,  $P < 0.001$ , odds ratio = 1.98). Whether participants worked away from home achieved a more marginal level of statistical significance ( $B = 0.43$ , Wald = 2.64,  $P = 0.104$ , odds ratio = 1.53). When entered next, the virtual behavior measure was, as before, statistically significant ( $B = -0.22$ , Wald = 3.95,  $P = 0.047$ , odds ratio = 0.80), with less social distancing being associated with a greater likelihood of testing positively for COVID-19. This was not true when the self-report measure of social distancing was instead entered as the third variable in the equation ( $B = -0.09$ , Wald = 0.53,  $P = 0.466$ , odds ratio = 0.92). Table 2 presents the relevant statistics when all four variables are considered simultaneously. Once again, the virtual behavior measure was the superior predictor.

## Discussion

The findings offer clear support regarding the efficacy of practicing social distancing. The more participants practiced social distancing, the less likely they were to have contracted COVID-19 over the next 4 mo. In fact, an increase of one SD on the virtual behavior measure of social distancing was associated with roughly a 20% reduction in the odds of contracting COVID-19. This was true both when considering only participants who reported having tested positively for the virus and when considering additional participants who, although untested, believed that they had contracted the virus.

Importantly, this prospective relation was observed at the level of the individual. The relation was apparent without aggregation across large numbers of people residing in a given locale—a characteristic of much of the research that has examined social distancing as a mitigating factor in the spread of the virus. In this respect, the current findings offer a unique form of additional evidence as to why individuals should indeed practice social distancing. What the individual does matters. It surely matters at

<sup>†</sup>Of these 1,686 cases, 5 involved missing values on one of the predictor variables. As a result, the analysis compared 199 cases of virus to 1,681 cases.

<sup>‡</sup>Once again, 5 of these latter cases involved missing values on one of the predictor variables. As a result, the analysis compared 85 positive test cases to 1,988 cases.

**Table 2. Binary logistic regression predicting having tested positively\***

Variable*	B	SE	Wald	P	Odds ratio
Preexisting conditions <sup>†</sup>	0.718	0.106	45.635	0.000	2.051
Leave home for work*	0.395	0.270	2.137	0.144	1.484
Self-reported behavior <sup>†</sup>	0.029	0.134	0.047	0.829	1.029
Virtual behavior <sup>†</sup>	-0.237	0.127	3.464	0.063	0.789
Constant	-3.501	0.154	519.935	0.000	NA

NA, not applicable.

\*Coded as 0 = No; 1 = Yes.

<sup>†</sup>Standardized.

the “collective” level, reducing transmission within the community—the very aim of officials’ social distancing directives and pleas. However, engaging in social distancing also carries with it clear and relatively immediate benefits for those who comply, leading to a reduced likelihood of contracting the disease for that specific individual.

It is important to acknowledge that our outcome variables regarding having contracted COVID-19 necessarily relied on participants’ reports. One might argue that individuals who earlier exhibited evidence of practicing social distancing were for some reason less likely to interpret any ambiguous physical symptoms as an indication of having the virus. Importantly, however, parallel results were observed regarding participants’ reports of having tested positively for COVID-19—a clear and salient outcome that seems considerably less likely to be misconstrued or misreported. Moreover, any biased interpretations and reports of illness would arguably be better predicted by the self-report measure of social distancing than the virtual behavior measure, precisely the opposite of what was observed. The specific combination of the virtual distancing measure as the predictor and positive test results as the outcome variable offers the strongest evidence for the efficacy of social distancing behavior as a means of reducing an individual’s likelihood of contracting COVID-19.

Interestingly, our results also offered a second critical contribution of the research. As noted, the virtual behavior measure of social distancing proved to be a better predictor of subsequent illness than did the self-report measure. The two measures were associated, but the correlation of 0.50 did not rise to a level that suggested these were essentially equivalent measures of the same construct, and differences in their predictive utility were evident. A significant relation was observed with the virtual measure both when predicting reports of believing one had contracted the virus and when restricting the sample to those who reported having tested positively. In contrast, the self-report measure revealed only a weak, marginally significant relation when predicting reports of illness and a null relation when predicting positive test outcomes. Moreover, when considered as simultaneous predictors, only the virtual distancing measure was associated with COVID-19 illness. Thus, the research findings also offer an important methodological advance regarding the value of behavioral measures that simulate concrete, real-world decisions.

The findings suggest a need for caution regarding the inferences that can be drawn from commonly employed self-report measures of social distancing. If we had included only the self-report measure, the current study would have seriously underestimated the effectiveness of social distancing. Of course, the self-report items that we employed were very general, asking participants to provide a relatively abstract summary assessment of their behavior. Questions involving greater specificity—for example, asking about adherence across an array of stipulated circumstances—may have proved more predictive of contracting COVID-19. Nevertheless, it is worth noting that the measures we

employed are very similar to those used in other recent studies examining compliance with pandemic-related recommendations. Many have relied on similarly general scales consisting of one or two items (e.g., refs. 22–24).

As noted earlier, self-report measures of behavior, especially general ones, are open to a number of potential pitfalls. Respondents must reconstruct their past, a memory process that may itself be influenced by individuals’ beliefs about how they prefer to view themselves. Whatever is implied by this reconstructive memory process must then be aligned with the response options offered by the questionnaire item. For example, just what does “following” social distancing recommendations “very strictly” mean? To which subjective label or scale point does one’s assessment of the past correspond? Again, how individuals wish to view themselves or portray themselves to others is likely to affect how the scale points are disambiguated. These potential sources of bias are far less characteristic of the behavioral decisions required by the graphical scenarios that comprise our virtual behavior measure. Participants made concrete, “in-the-moment” decisions about alternative courses of action differing in the extent to which they allowed for social distancing. They interactively separated themselves visually from oncoming passersby; they selected a position on a crowded beach; they decided whether to walk along a route that required encountering other people. Participants made choices, much as they would in the real world.

The data appear to suggest that a sizeable number of participants offered self-reports that were overestimates of their actual social distancing behavior. Whereas the virtual behavior data displayed a normal distribution, the distribution of scores on the self-report measure was skewed with a substantial majority responding at or near the positive endpoint of the scale ( $M = 5.98$  on a seven-point scale,  $SD = 1.18$ ). Such overestimation is to be expected to the extent that participants wished to believe themselves as having acted in manners that avoided placing their health, or that of others, at risk. Such self-beliefs have been shown to influence retrospective memory processes (e.g., refs. 10, 14, and 15).

Might the virtual behavior items have served not solely as a means of estimating the participants’ real-world social distancing behavior but also as a force that enhanced the salience of social distancing norms and motivated stronger subsequent adherence to the norm? In other words, might the simulations have acted more as an intervention that changed some participants’ responses to the graphical scenarios and, more importantly, their subsequent real-world behavior? Although any such possibility may have contributed to the effect and surely does not detract from the significance of the prospective findings regarding illness, we believe it is more likely that the virtual behavior measure primarily captured individual differences in commitment to the practice of social distancing that existed at the time of measurement. In an earlier study (21), greater social distancing on the virtual behavior measure related strongly to a number of individual difference variables, such as greater trust in scientists, lower confidence in President Trump, lower conspiratorial ideation tendencies, greater interpersonal compassion, greater scientific literacy, less exposure to FOX News, and more accurate knowledge regarding COVID-19. Moreover, these individual difference variables and virtual social distancing behavior were themselves associated with expressed concerns about the pandemic and the severity of the threat it posed. The conceptual coherence of this diverse array of correlations leads us to the view that the associations emerged because our virtual behavior measure did in fact capture participants’ current social distancing behavior.

Whether these multiple associations also may reflect the identification of individuals who, as a consequence of responding to the virtual behavior items, experienced an enhanced and sustained motivation to subsequently adhere to the social

distancing norm is an open question. However, the two processes are not mutually exclusive, and it is possible that increased salience and/or motivation to adhere to the social distancing guidelines contributed to the longitudinal effects that we observed. Future researchers may wish to experimentally manipulate the presence or absence of the virtual behavior measure to more decisively test whether these measures may also act as an intervention that decreases the likelihood that an individual will contract the illness subsequently.

Irrespective of the processes that may have contributed to the effects we observed, the current findings clearly illustrate the value of social distancing as a means of mitigating the likelihood

of contracting COVID-19. The study offers prospective evidence to that effect. In contrast to past research, we show that social distancing matters not only at the aggregate level but also critically at the level of the individual. Decisions that reflect the sound practice of social distancing reduce individuals' risk of contracting COVID-19.

**Data Availability.** The SPSS data file has been deposited in the Open Science Framework (<https://osf.io/x79ak/>).

**ACKNOWLEDGMENTS.** This research was supported by a grant from the National Science Foundation (Grant 2031097).

1. E. C. Lee, N. I. Wada, M. K. Grabowski, E. S. Gurley, J. Lessler, The engines of SARS-CoV-2 spread. *Science* **370**, 406–407 (2020).
2. D. K. Chu *et al.*, Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and COVID-19: A systematic review and meta-analysis. *Lancet* **395**, 1973–1987 (2020).
3. N. M. Ferguson *et al.*, Impact of non-pharmaceutical interventions (NPIs) to reduce COVID-19 mortality and healthcare demand. <https://www.doi.org/10.25561/77482>. Accessed 27 January 2021.
4. M. Greenstone, V. Nigam, Does social distancing matter? *SSRN* [Preprint] (2020). <https://doi.org/10.2139/ssrn.3561244> (Accessed 27 January 2021).
5. S. M. Kissler, C. Tedijanto, M. Lipsitch, Y. Grad, Social distancing strategies for curbing the COVID-19 epidemic. *medRxiv* [Preprint] (2020). <https://doi.org/10.1101/2020.03.22.20041079> (Accessed 27 January 2021).
6. D. J. McGraill, J. Dai, K. M. McAndrews, R. Kalluri, Enacting national social distancing policies corresponds with dramatic reduction in COVID19 infection rates. *PLoS One* **15**, e0236619 (2020).
7. C. Courtemanche, J. Garuccio, A. Le, J. Pinkston, A. Yelowitz, Strong social distancing measures in the United States reduced the covid-19 growth rate. *Health Aff. (Millwood)* **39**, 1237–1246 (2020).
8. T. VoPham *et al.*, Effect of social distancing on COVID-19 incidence and mortality in the US. *medRxiv* [Preprint] (2020). <https://doi.org/10.1101/2020.06.10.20127589> (Accessed 27 January 2021).
9. A. Gollwitzer, C. Martel, J. Marshall, J. M. Höhs, J. A. Bargh, Connecting self-reported social distancing to real-world behavior at the individual and U.S. state level. *PsyArXiv* [Preprint] (2020). <https://doi.org/10.31234/osf.io/kvnwp> (Accessed 27 January 2021).
10. E. Balcetis, Where the motivation resides and self-deception hides: How motivated cognition accomplishes self-deception. *Soc. Personal. Psychol. Compass* **2**, 361–381 (2008).
11. R. J. Fisher, Social desirability bias and the validity of indirect questioning. *J. Consum. Res.* **20**, 303–315 (1993).
12. R. C. Gur, H. A. Sackeim, Self-deception: A concept in search of a phenomenon. *J. Pers. Soc. Psychol.* **37**, 147–169 (1979).
13. M. R. Leary, R. M. Kowalski, Impression management: A literature review and two component model. *Psychol. Bull.* **107**, 34–47 (1990).
14. M. Kouchaki, F. Gino, Memories of unethical actions become obfuscated over time. *Proc. Natl. Acad. Sci. U.S.A.* **113**, 6166–6171 (2016).
15. M. Ross, Relation of implicit theories to the construction of personal histories. *Psychol. Rev.* **96**, 341–357 (1989).
16. M. Buhrmester, T. Kwang, S. D. Gosling, Amazon's mechanical turk: A new source of inexpensive, yet high-quality, data? *Perspect. Psychol. Sci.* **6**, 3–5 (2011).
17. G. Paolacci, J. Chandler, Inside the Turk: Understanding Mechanical Turk as a participant pool. *Curr. Dir. Psychol. Sci.* **23**, 184–188 (2014).
18. A. J. Berinsky, G. A. Huber, G. S. Lenz, Evaluating online labor markets for experimental research: Amazon.com's mechanical turk. *Polit. Anal.* **20**, 351–368 (2012).
19. D. Hauser, G. Paolacci, J. Chandler, "Common concerns with MTurk as a participant pool evidence and solutions" in *Handbook of Research Methods in Consumer Psychology*, F. R. Kardes, P. M. Herr, N. Schwarz, Eds. (Routledge, New York, NY, 2019), pp. 319–336.
20. S. Clifford, R. M. Jewell, P. D. Waggoner, Are samples drawn from mechanical turk valid for research on political ideology? *Research & Politics*, 10.1177/2053168015622072 (2015).
21. R. H. Fazio *et al.*, Who is (not) complying with the social distancing directive and why? Testing a general framework of compliance with a behavioral measure of social distancing. *medRxiv* [Preprint] (2020). <https://doi.org/10.1101/2020.10.26.20219634> (Accessed 27 January 2021).
22. M. S. Hagger, S. R. Smith, J. J. Keech, S. A. Moyers, K. Hamilton, Predicting social distancing intention and behavior during the COVID-19 pandemic: An integrated social cognition model. *Ann. Behav. Med.* **54**, 713–727 (2020).
23. F. J. Kachanoff, Y. E. Bigman, K. Kapsaskis, K. Gray, Measuring realistic and symbolic threats of COVID-19 and their unique impacts on well-being and adherence to public health behaviors. *Soc. Psychol. Personal. Sci.*, 10.1177/1948550620931634 (2020).
24. M. Zajenkowski, P. K. Jonason, M. Leniarska, Z. Kozakiewicz, Who complies with the restrictions to reduce the spread of COVID-19?: Personality and perceptions of the COVID-19 situation. *Pers. Individ. Dif.* **166**, 110199 (2020).