

# Grand Unified Theories

Stuart Raby

*Department of Physics, The Ohio State University, 191 W. Woodruff Ave.  
Columbus, OH 43210*

## 1 Grand Unification

### 1.1 Standard Model : An Introduction

In spite of all the successes of the Standard Model [SM] it is unlikely to be the final theory. It leaves many unanswered questions. Why the local gauge interactions  $SU(3)_C \times SU(2)_L \times U(1)_Y$  and why 3 families of quarks and leptons? Moreover why does one family consist of the states  $[Q, u^c, d^c; L, e^c]$  transforming as  $[(3, 2, 1/3), (\bar{3}, 1, -4/3), (\bar{3}, 1, 2/3); (1, 2, -1), (1, 1, 2)]$ , where  $Q = (u, d)$  and  $L = (\nu, e)$  are  $SU(2)_L$  doublets and  $u^c, d^c, e^c$  are charge conjugate  $SU(2)_L$  singlet fields with the  $U(1)_Y$  quantum numbers given? [We use the convention that electric charge  $Q_{EM} = T_{3L} + Y/2$  and all fields are left handed.] Note the SM gauge interactions of quarks and leptons are completely fixed by their gauge charges. Thus if we understood the origin of this charge quantization, we would also understand why there are no fractionally charged hadrons. Finally, what is the origin of quark and lepton masses or the apparent hierarchy of family masses and quark mixing angles? Perhaps if we understood this, we would also know the origin of CP violation, the solution to the strong CP problem, the origin of the cosmological matter - antimatter asymmetry or the nature of dark matter.

The SM has 19 arbitrary parameters; their values are chosen to fit the data. Three arbitrary gauge couplings:  $g_3, g, g'$  (where  $g, g'$  are the  $SU(2)_L, U(1)_Y$  couplings, respectively) or equivalently  $\alpha_s = (g_3^2/4\pi), \alpha_{EM} = (e^2/4\pi)$  ( $e = g \sin \theta_W$ ) and  $\sin^2 \theta_W = (g')^2/(g^2 + (g')^2)$ . In addition there are 13 parameters associated with the 9 charged fermion masses and the four mixing angles in the CKM matrix. The remaining 3 parameters are  $v, \lambda$  [the Higgs VEV and quartic coupling] (or equivalently  $M_Z, m_h^0$ ) and the QCD  $\theta$  parameter. In addition, data from neutrino oscillation experiments provide convincing evidence for neutrino masses. With 3 light Majorana neutrinos there are at least 9 additional parameters in the neutrino sector; 3 masses and 6 mixing angles and phases. In summary, the SM has too many arbitrary parameters and leaves open too many unresolved questions to be considered complete. These are the problems which grand unified theories hope to address.

### 1.2 Charge Quantization

In the Standard Model, quarks and leptons are on an equal footing; both fundamental particles without substructure. It is now clear that they may be two faces of the same coin; unified, for example, by extending QCD (or  $SU(3)_C$ ) to include leptons

as the fourth color,  $SU(4)_C$  [1]. The complete Pati-Salam gauge group is  $SU(4)_C \times SU(2)_L \times SU(2)_R$  with the states of one family  $[(Q, L), (Q^c, L^c)]$  transforming as  $[(4, 2, 1), (\bar{4}, 1, 2)]$  where  $Q^c = (d^c, u^c)$ ,  $L^c = (e^c, \nu^c)$  are doublets under  $SU(2)_R$ . Electric charge is now given by the relation  $Q_{EM} = T_{3L} + T_{3R} + 1/2(B - L)$  and  $SU(4)_C$  contains the subgroup  $SU(3)_C \times (B - L)$  where  $B$  ( $L$ ) is baryon (lepton) number. Note  $\nu^c$  has no SM quantum numbers and is thus completely “sterile”. It is introduced to complete the  $SU(2)_R$  lepton doublet. This additional state is desirable when considering neutrino masses.

Although quarks and leptons are unified with the states of one family forming two irreducible representations of the gauge group; there are still 3 independent gauge couplings (two if one also imposes parity, i.e.  $L \leftrightarrow R$  symmetry). As a result the three low energy gauge couplings are still independent arbitrary parameters. This difficulty is resolved by embedding the SM gauge group into the simple unified gauge group, Georgi-Glashow  $SU(5)$ , with one universal gauge coupling  $\alpha_G$  defined at the grand unification scale  $M_G$  [2]. Quarks and leptons still sit in two irreducible representations, as before, with a  $\mathbf{10} = [Q, u^c, e^c]$  and  $\bar{\mathbf{5}} = [d^c, L]$ . Nevertheless, the three low energy gauge couplings are now determined in terms of two independent parameters :  $\alpha_G$  and  $M_G$ . Hence there is one prediction.

In order to break the electroweak symmetry at the weak scale and give mass to quarks and leptons, Higgs doublets are needed which can sit in either a  $\mathbf{5}_H$  or  $\bar{\mathbf{5}}_H$ . The additional 3 states are color triplet Higgs scalars. The couplings of these color triplets violate baryon and lepton number and nucleons decay via the exchange of a single color triplet Higgs scalar. Hence in order not to violently disagree with the non-observation of nucleon decay, their mass must be greater than  $\sim 10^{10-11}$  GeV. Moreover, in supersymmetric GUTs, in order to cancel anomalies as well as give mass to both up and down quarks, both Higgs multiplets  $\mathbf{5}_H$ ,  $\bar{\mathbf{5}}_H$  are required. As we shall discuss later, nucleon decay now constrains the color triplet Higgs states in a SUSY GUT to have mass significantly greater than  $M_G$ .

Complete unification is possible with the symmetry group  $SO(10)$  with one universal gauge coupling  $\alpha_G$  and one family of quarks and leptons sitting in the 16 dimensional spinor representation  $\mathbf{16} = [\mathbf{10} + \bar{\mathbf{5}} + \mathbf{1}]$  [3]. The  $SU(5)$  singlet  $\mathbf{1}$  is identified with  $\nu^c$ . In Table 1 we present the states of one family of quarks and leptons, as they appear in the  $\mathbf{16}$ . It is an amazing and perhaps even profound fact that all the states of a single family of quarks and leptons can be represented digitally as a set of 5 zeros and/or ones or equivalently as the tensor product of 5 “spin” 1/2 states with  $\pm = |\pm \frac{1}{2}\rangle$  and with the condition that we have an even number of  $|-\rangle$  spins. The first three “spins” correspond to  $SU(3)_C$  color quantum numbers, while the last two are  $SU(2)_L$  weak quantum numbers. In fact an  $SU(3)_C$  rotation just raises one color index and lowers another, thereby changing colors  $\{r, b, y\}$ . Similarly an  $SU(2)_L$  rotation raises one weak index and lowers another, thereby flipping the weak isospin from up to down or vice versa. In this representation weak hypercharge  $Y$  is given by the simple relation  $Y = 2/3(\sum \text{color spins}) - (\sum \text{weak spins})$ .  $SU(5)$  rotations then raise (or lower) a color index, while at the same time lowering (or raising) a weak index. It is easy to see that such rotations can mix the states  $\{Q, u^c, e^c\}$  and  $\{d^c, L\}$  among themselves and  $\nu^c$  is a singlet. The new  $SO(10)$  rotations [not in  $SU(5)$ ] are then given by either raising or lowering any two spins. For example, by lowering the two weak indices  $\nu^c$  rotates into  $e^c$ , etc.

Table 1: The quantum numbers of the **16** dimensional representation of  $SO(10)$ .

State	$Y$	Color	Weak
$\nu^c$	0	+ + +	++
$e^c$	2	+ + +	--
$u_r$	1/3	- + +	+-
$d_r$	1/3	- + +	-+
$u_b$	1/3	+ - +	+-
$d_b$	1/3	+ - +	-+
$u_y$	1/3	+ + -	+-
$d_y$	1/3	+ + -	-+
$u_r^c$	-4/3	+ - -	++
$u_b^c$	-4/3	- + -	++
$u_y^c$	-4/3	- - +	++
$d_r^c$	2/3	+ - -	--
$d_b^c$	2/3	- + -	--
$d_y^c$	2/3	- - +	--
$\nu$	-1	- - -	+-
$e$	-1	- - -	-+

$SO(10)$  has two inequivalent maximal breaking patterns.  $SO(10) \rightarrow SU(5) \times U(1)_X$  and  $SO(10) \rightarrow SU(4)_C \times SU(2)_L \times SU(2)_R$ . In the first case we obtain Georgi-Glashow  $SU(5)$  if  $Q_{EM}$  is given in terms of  $SU(5)$  generators alone or so-called flipped  $SU(5)$  [4] if  $Q_{EM}$  is partly in  $U(1)_X$ . In the latter case we have the Pati-Salam symmetry. If  $SO(10)$  breaks directly to the SM at  $M_G$ , then we retain the prediction for gauge coupling unification. However more possibilities for breaking (hence more breaking scales and more parameters) are available in  $SO(10)$ . Nevertheless with one breaking pattern  $SO(10) \rightarrow SU(5) \rightarrow$  SM, where the last breaking scale is  $M_G$ , the predictions from gauge coupling unification are preserved. The Higgs multiplets in minimal  $SO(10)$  are contained in the fundamental  $\mathbf{10}_H = [\mathbf{5}_H, \bar{\mathbf{5}}_H]$  representation. Note, only in  $SO(10)$  does the gauge symmetry distinguish quark and lepton multiplets from Higgs multiplets.

Finally, larger symmetry groups have been considered. For example,  $E(6)$  has a fundamental representation  $\mathbf{27}$  which under  $SO(10)$  transforms as a  $[\mathbf{16} + \mathbf{10} + \mathbf{1}]$ . The breaking pattern  $E(6) \rightarrow SU(3)_C \times SU(3)_L \times SU(3)_R$  is also possible. With the additional permutation symmetry  $Z(3)$  interchanging the three  $SU(3)$ s we obtain so-called ‘‘trinification’’ [5] with a universal gauge coupling. The latter breaking pattern has been used in phenomenological analyses of the heterotic string [6]. However, in larger symmetry groups, such as  $E(6)$ ,  $SU(6)$ , etc., there are now many more states which have not been observed and must be removed from the effective low energy theory. In particular, three families of  $\mathbf{27}$ s in  $E(6)$  contain three Higgs type multiplets transforming as  $\mathbf{10}$ s of  $SO(10)$ . This makes these larger symmetry groups

unattractive starting points for model building.

### 1.3 String Theory and Orbifold GUTs

Orbifold compactification of the heterotic string [7, 8, 9], and recent field theoretic constructions known as orbifold GUTs [10], contain grand unified symmetries realized in 5 and 6 dimensions. However, upon compactifying all but four of these extra dimensions, only the MSSM is recovered as a symmetry of the effective four dimensional field theory.<sup>1</sup> These theories can retain many of the nice features of four dimensional SUSY GUTs, such as charge quantization, gauge coupling unification and sometimes even Yukawa unification; while at the same time resolving some of the difficulties of 4d GUTs, in particular problems with unwieldy Higgs sectors necessary for spontaneously breaking the GUT symmetry, and problems with doublet-triplet Higgs splitting or rapid proton decay. We will comment further on the corrections to the four dimensional GUT picture due to orbifold GUTs in the following sections.

### 1.4 Gauge coupling unification

The biggest paradox of grand unification is to understand how it is possible to have a universal gauge coupling  $g_G$  in a grand unified theory [GUT] and yet have three unequal gauge couplings at the weak scale with  $g_3 > g > g'$ . The solution is given in terms of the concept of an effective field theory [EFT] [16]. The GUT symmetry is spontaneously broken at the scale  $M_G$  and all particles not in the SM obtain mass of order  $M_G$ . When calculating Green's functions with external energies  $E \gg M_G$ , we can neglect the mass of all particles in the loop and hence all particles contribute to the renormalization group running of the universal gauge coupling. However, for  $E \ll M_G$  one can consider an effective field theory including only the states with mass  $< E \ll M_G$ . The gauge symmetry of the EFT is  $SU(3)_C \times SU(2)_L \times U(1)_Y$  and the three gauge couplings renormalize independently. The states of the EFT include only those of the SM; 12 gauge bosons, 3 families of quarks and leptons and one or more Higgs doublets. At  $M_G$  the two effective theories [the GUT itself is most likely the EFT of a more fundamental theory defined at a higher scale] must give identical results; hence we have the boundary conditions  $g_3 = g_2 = g_1 \equiv g_G$  where at any scale

---

<sup>1</sup>Also, in recent years there has been a great deal of progress in constructing three and four family models in Type IIA string theory with intersecting D6 branes [11]. Although these models can incorporate  $SU(5)$  or a Pati-Salam symmetry group in four dimensions, they typically have problems with gauge coupling unification. In the former case this is due to charged exotics which affect the RG running, while in the latter case the  $SU(4) \times SU(2)_L \times SU(2)_R$  symmetry never unifies. Note, heterotic string theory models also exist whose low energy effective 4d field theory is a SUSY GUT [12]. These models have all the virtues and problems of 4d GUTs. Finally, many heterotic string models have been constructed with the standard model gauge symmetry in 4d and no intermediate GUT symmetry in less than 10d. Recently some minimal 3 family supersymmetric models have been constructed [13, 14]. These theories may retain some of the symmetry relations of GUTs, however the unification scale would typically be the string scale, of order  $5 \times 10^{17}$  GeV, which is inconsistent with low energy data. A way out of this problem was discovered in the context of the strongly coupled heterotic string, defined in an effective 11 dimensions [15]. In this case the 4d Planck scale (which controls the value of the string scale) now unifies with the GUT scale.

$\mu < M_G$  we have  $g_2 \equiv g$  and  $g_1 = \sqrt{5/3} g'$ . Then using two low energy couplings, such as  $\alpha_s(M_Z)$ ,  $\alpha_{EM}(M_Z)$ , the two independent parameters  $\alpha_G$ ,  $M_G$  can be fixed. The third gauge coupling,  $\sin^2 \theta_W$  in this case, is then predicted. This was the procedure up until about 1991 [17, 18]. Subsequently, the uncertainties in  $\sin^2 \theta_W$  were reduced ten fold. Since then,  $\alpha_{EM}(M_Z)$ ,  $\sin^2 \theta_W$  have been used as input to predict  $\alpha_G$ ,  $M_G$  and  $\alpha_s(M_Z)$  [19].

We emphasize that the above boundary condition is only valid when using one loop renormalization group [RG] running. With precision electroweak data, however, it is necessary to use two loop RG running. Hence one must include one loop threshold corrections to gauge coupling boundary conditions at both the weak and GUT scales. In this case it is always possible to define the GUT scale as the point where  $\alpha_1(M_G) = \alpha_2(M_G) \equiv \tilde{\alpha}_G$  and  $\alpha_3(M_G) = \tilde{\alpha}_G (1 + \epsilon_3)$ . The threshold correction  $\epsilon_3$  is a logarithmic function of all states with mass of order  $M_G$  and  $\tilde{\alpha}_G = \alpha_G + \Delta$  where  $\alpha_G$  is the GUT coupling constant above  $M_G$  and  $\Delta$  is a one loop threshold correction. To the extent that gauge coupling unification is perturbative, the GUT threshold corrections are small and calculable. This presumes that the GUT scale is sufficiently below the Planck scale or any other strong coupling extension of the GUT, such as a strongly coupled string theory.

Supersymmetric grand unified theories [SUSY GUTs] are an extension of non-SUSY GUTs [20]. The key difference between SUSY GUTs and non-SUSY GUTs is the low energy effective theory. The low energy effective field theory in a SUSY GUT is assumed to satisfy N=1 supersymmetry down to scales of order the weak scale in addition to the SM gauge symmetry. Hence the spectrum includes all the SM states plus their supersymmetric partners. It also includes one pair (or more) of Higgs doublets; one to give mass to up-type quarks and the other to down-type quarks and charged leptons. Two doublets with opposite hypercharge  $Y$  are also needed to cancel fermionic triangle anomalies. Finally, it is important to recognize that a low energy SUSY breaking scale (the scale at which the SUSY partners of SM particles obtain mass) is necessary to solve the gauge hierarchy problem.

Simple non-SUSY  $SU(5)$  is ruled out; initially by the increased accuracy in the measurement of  $\sin^2 \theta_W$  and by early bounds on the proton lifetime (see below) [18]. However, by now LEP data [19] has conclusively shown that SUSY GUTs is the *new standard model*; by which we mean the theory used to guide the search for new physics beyond the present SM (see Fig. 1). SUSY extensions of the SM have the property that their effects decouple as the effective SUSY breaking scale is increased. Any theory beyond the SM must have this property simply because the SM works so well. However, the SUSY breaking scale cannot be increased with impunity, since this would reintroduce a gauge hierarchy problem. Unfortunately there is no clear-cut answer to the question, when is the SUSY breaking scale too high. A conservative bound would suggest that the third generation quarks and leptons must be lighter than about 1 TeV, in order that the one loop corrections to the Higgs mass from Yukawa interactions remains of order the Higgs mass bound itself.

At present gauge coupling unification within SUSY GUTs works extremely well. Exact unification at  $M_G$ , with two loop renormalization group running from  $M_G$  to  $M_Z$ , and one loop threshold corrections at the weak scale, fits to within 3  $\sigma$  of the present precise low energy data. A small threshold correction at  $M_G$  ( $\epsilon_3 \sim -3\%$ )




Figure 1: Gauge coupling unification in non-SUSY GUTs on the left vs. SUSY GUTs on the right using the LEP data as of 1991. Note, the difference in the running for SUSY is the inclusion of supersymmetric partners of standard model particles at scales of order a TeV (Fig. taken from Ref. [21]). Given the present accurate measurements of the three low energy couplings, in particular  $\alpha_s(M_Z)$ , GUT scale threshold corrections are now needed to precisely fit the low energy data. The dark blob in the plot on the right represents these model dependent corrections.

to - 4%) is sufficient to fit the low energy data precisely [22, 23, 24].<sup>2</sup> This may be compared to non-SUSY GUTs where the fit misses by  $\sim 12 \sigma$  and a precise fit requires new weak scale states in incomplete GUT multiplets or multiple GUT breaking scales.<sup>3</sup>

Following the analysis of Ref. [24] let us try to understand the need for the GUT threshold correction and its order of magnitude. The renormalization group equations relate the low energy gauge coupling constants  $\alpha_i(M_Z)$ ,  $i = 1, 2, 3$  to the value of the

<sup>2</sup>This result implicitly assumes universal GUT boundary conditions for soft SUSY breaking parameters at  $M_G$ . In the simplest case we have a universal gaugino mass  $M_{1/2}$ , a universal mass for squarks and sleptons  $m_{16}$  and a universal Higgs mass  $m_{10}$ , as motivated by  $SO(10)$ . In some cases, threshold corrections to gauge coupling unification can be exchanged for threshold corrections to soft SUSY parameters. See for example, Ref. [25] and references therein.

<sup>3</sup>Non-SUSY GUTs with a more complicated breaking pattern can still fit the data. For example, non-SUSY  $SO(10) \rightarrow SU(4)_C \times SU(2)_L \times SU(2)_R \rightarrow SM$  with the second breaking scale of order an intermediate scale, determined by light neutrino masses using the see-saw mechanism, can fit the low energy data for gauge couplings [26] and at the same time survive nucleon decay bounds [27], discussed in Section 1.5.

unification scale  $\Lambda_U$  and the GUT coupling  $\alpha_U$  by the expression

$$\frac{1}{\alpha_i(M_Z)} = \frac{1}{\alpha_U} + \frac{b_i}{2\pi} \log\left(\frac{\Lambda_U}{M_Z}\right) + \delta_i \quad (1)$$

where  $\Lambda_U$  is the GUT scale evaluated at one loop and the threshold corrections,  $\delta_i$ , are given by  $\delta_i = \delta_i^{(2)} + \delta_i^{(l)} + \delta_i^{(g)}$  with  $\delta_i^{(2)}$  representing two loop running effects,  $\delta_i^{(l)}$  the light threshold corrections at the SUSY breaking scale and  $\delta_i^{(g)} = \delta_i^{(h)} + \delta_i^{(b)}$  representing GUT scale threshold corrections. Note, in this analysis, the two loop RG running is treated on the same footing as weak and GUT scale threshold corrections. One then obtains the prediction

$$(\alpha_3(M_Z) - \alpha_3^{LO}(M_Z))/\alpha_3^{LO}(M_Z) = -\alpha_3^{LO}(M_Z) \delta_s \quad (2)$$

where  $\alpha_3^{LO}(M_Z)$  is the leading order one loop RG result and  $\delta_s = \frac{1}{7}(5\delta_1 - 12\delta_2 + 7\delta_3)$  is the net threshold correction. [A similar formula applies at the GUT scale with the GUT threshold correction,  $\epsilon_3$ , given by  $\epsilon_3 = -\tilde{\alpha}_G \delta_s^{(g)}$ .] Given the experimental inputs [28]:

$$\begin{aligned} \alpha_{em}^{-1}(M_Z) &= 127.906 \pm 0.019 \\ \sin^2\theta_W(M_Z) &= 0.2312 \pm 0.0002 \\ \alpha_3(M_Z) &= 0.1187 \pm 0.0020 \end{aligned} \quad (3)$$

and taking into account the light threshold corrections, assuming an ensemble of 10 SUSY spectra [24] (corresponding to the Snow Mass benchmark points), we have

$$\alpha_3^{LO}(M_Z) \approx 0.118 \quad (4)$$

and

$$\begin{aligned} \delta_s^{(2)} &\approx -0.82 \\ \delta_s^{(l)} &\approx -0.50 + \frac{19}{28\pi} \log \frac{M_{SUSY}}{M_Z}. \end{aligned} \quad (5)$$

For  $M_{SUSY} = 1$  TeV, we have  $\delta_s^{(2)} + \delta_s^{(l)} \approx -0.80$ . Since the one loop result  $\alpha_3^{LO}(M_Z)$  is very close to the experimental value, we need  $\delta_s \approx 0$  or equivalently,  $\delta_s^{(g)} \approx 0.80$ . This corresponds, at the GUT scale, to  $\epsilon_3 \approx -3\%$ .<sup>4</sup>

---

<sup>4</sup>In order to fit the low energy data for gauge coupling constants we require a relative shift in  $\alpha_3(M_G)$  of order 3% due to GUT scale threshold corrections. If these GUT scale corrections were not present, however, weak scale threshold corrections of order 9% (due to the larger value of  $\alpha_3$  at  $M_Z$ ) would be needed to resolve the discrepancy with the data for exact gauge coupling unification at  $M_G$ . Leaving out the fact that any consistent GUT necessarily contributes threshold corrections at the GUT scale, it is much more difficult to find the necessary larger corrections at the weak scale. For example, we need  $M_{SUSY} \approx 40$  TeV for the necessary GUT scale threshold correction to vanish.

In four dimensional SUSY GUTs, the threshold correction  $\epsilon_3$  receives a positive contribution from Higgs doublets and triplets.<sup>5</sup> Thus a larger, negative contribution must come from the GUT breaking sector of the theory. This is certainly possible in specific SO(10) [29] or SU(5) [30] models, but it is clearly a significant constraint on the 4d GUT sector of the theory. In five or six dimensional orbifold GUTs, on the other hand, the “GUT scale” threshold correction comes from the Kaluza-Klein modes between the compactification scale,  $M_c$ , and the effective cutoff scale  $M_*$ .<sup>6</sup> Thus, in orbifold GUTs, gauge coupling unification at two loops is only consistent with the low energy data with a fixed value for  $M_c$  and  $M_*$ .<sup>7</sup> Typically, one finds  $M_c < M_G = 3 \times 10^{16}$  GeV, where  $M_G$  is the 4d GUT scale. Since the grand unified gauge bosons, responsible for nucleon decay, get mass at the compactification scale, the result  $M_c < M_G$  for orbifold GUTs has significant consequences for nucleon decay.

A few final comments are in order. We do not consider the scenario of split supersymmetry [33] in this review. In this scenario squarks and sleptons have mass at a scale  $\tilde{m} \gg M_Z$ , while gauginos and Higgsinos have mass of order the weak scale. Gauge coupling unification occurs at a scale of order  $10^{16}$  GeV, *provided that the scale  $\tilde{m}$  lies in the range  $10^3 - 10^{11}$  GeV* [34]. A serious complaint concerning the split SUSY scenario is that it does not provide a solution to the gauge hierarchy problem. Moreover, it is only consistent with grand unification if it also postulates an “intermediate” scale,  $\tilde{m}$ , for scalar masses. In addition, it is in conflict with  $b - \tau$  Yukawa unification, unless  $\tan \beta$  is fine-tuned to be close to 1 [34].<sup>8</sup>

We have also neglected to discuss non-supersymmetric GUTs in four dimensions which still survive once one allows for several scales of GUT symmetry breaking [26]. Finally, it has been shown that non-supersymmetric GUTs in warped 5 dimensional orbifolds can be consistent with gauge coupling unification, assuming that the right-handed top quark and the Higgs doublets are composite-like objects with a compositeness scale of order a TeV [36].

## 1.5 Nucleon Decay

Baryon number is necessarily violated in any GUT [37]. In  $SU(5)$  nucleons decay via the exchange of gauge bosons with GUT scale masses, resulting in dimension

---

<sup>5</sup>Note, the Higgs contribution is given by  $\epsilon_3 = \frac{3\tilde{\alpha}_G}{5\pi} \log \left| \frac{\tilde{M}_t \gamma}{M_G} \right|$  where  $\tilde{M}_t$  is the effective color triplet Higgs mass (setting the scale for dimension 5 baryon and lepton number violating operators) and  $\gamma = \lambda_b/\lambda_t$  at  $M_G$ . Since  $\tilde{M}_t$  is necessarily greater than  $M_G$ , the Higgs contribution to  $\epsilon_3$  is positive.

<sup>6</sup>In string theory, the cutoff scale is the string scale.

<sup>7</sup>It is interesting to note that a ratio  $M_*/M_c \sim 100$ , needed for gauge coupling unification to work in orbifold GUTs is typically the maximum value for this ratio consistent with perturbativity [31]. In addition, in orbifold GUTs brane-localized gauge kinetic terms may destroy the successes of gauge coupling unification. However, for values of  $M_*/M_c = M_*\pi R \gg 1$  the unified bulk gauge kinetic terms can dominate over the brane-localized terms [32].

<sup>8</sup> $b - \tau$  Yukawa unification only works for  $\tilde{m} < 10^4$  for  $\tan \beta \geq 1.5$ . This is because the effective theory between the gaugino mass scale and  $\tilde{m}$  includes only one Higgs doublet, as in the standard model. In this case, the large top quark Yukawa coupling tends to increase the ratio  $\lambda_b/\lambda_\tau$  as one runs down in energy below  $\tilde{m}$ . This is opposite to what happens in MSSM where the large top quark Yukawa coupling decreases the ratio  $\lambda_b/\lambda_\tau$  [35].



6 baryon number violating operators suppressed by  $(1/M_G^2)$ . The nucleon lifetime is calculable and given by  $\tau_N \propto M_G^4/(\alpha_G^2 m_p^5)$ . The dominant decay mode of the proton (and the baryon violating decay mode of the neutron), via gauge exchange, is  $p \rightarrow e^+ \pi^0$  ( $n \rightarrow e^+ \pi^-$ ). In any simple gauge symmetry, with one universal GUT coupling and scale  $(\alpha_G, M_G)$ , the nucleon lifetime from gauge exchange is calculable. Hence, the GUT scale may be directly observed via the extremely rare decay of the nucleon. Experimental searches for nucleon decay began with the Kolar Gold Mine, Homestake, Soudan, NUSEX, Frejus, HPW, and IMB detectors [17]. The present experimental bounds come from Super-Kamiokande and Soudan II. We discuss these results shortly. Non-SUSY GUTs are also ruled out by the non-observation of nucleon decay [18]. In SUSY GUTs, the GUT scale is of order  $3 \times 10^{16}$  GeV, as compared to the GUT scale in non-SUSY GUTs which is of order  $10^{15}$  GeV. Hence the dimension 6 baryon violating operators are significantly suppressed in SUSY GUTs [20] with  $\tau_p \sim 10^{34-38}$  yrs.

However, in SUSY GUTs there are additional sources for baryon number violation – dimension 4 and 5 operators [38]. Although our notation does not change, when discussing SUSY GUTs all fields are implicitly bosonic superfields and the operators considered are the so-called F terms which contain two fermionic components and the rest scalars or products of scalars. Within the context of  $SU(5)$  the dimension 4 and 5 operators have the form  $(\mathbf{10} \bar{\mathbf{5}} \bar{\mathbf{5}}) \supset (u^c d^c d^c) + (Q L d^c) + (e^c L L)$  and  $(\mathbf{10} \mathbf{10} \mathbf{10} \bar{\mathbf{5}}) \supset (Q Q Q L) + (u^c u^c d^c e^c) + B$  and  $L$  conserving terms, respectively. The dimension 4 operators are renormalizable with dimensionless couplings; similar to Yukawa couplings. On the other hand, the dimension 5 operators have a dimensionful coupling of order  $(1/M_G)$ .

The dimension 4 operators violate baryon number or lepton number, respectively, but not both. The nucleon lifetime is extremely short if both types of dimension 4 operators are present in the low energy theory. However both types can be eliminated by requiring R parity. In  $SU(5)$  the Higgs doublets reside in a  $\mathbf{5}_H$ ,  $\bar{\mathbf{5}}_H$  and R parity distinguishes the  $\bar{\mathbf{5}}$  (quarks and leptons) from  $\bar{\mathbf{5}}_H$  (Higgs). R parity [39] (or its cousin, family reflection symmetry (see Dimopoulos and Georgi [20] and DRW [40]) takes  $F \rightarrow -F$ ,  $H \rightarrow H$  with  $F = \{\mathbf{10}, \bar{\mathbf{5}}\}$ ,  $H = \{\bar{\mathbf{5}}_H, \mathbf{5}_H\}$ . This forbids the dimension 4 operator  $(\mathbf{10} \bar{\mathbf{5}} \bar{\mathbf{5}})$ , but allows the Yukawa couplings of the form  $(\mathbf{10} \bar{\mathbf{5}} \bar{\mathbf{5}}_H)$  and  $(\mathbf{10} \mathbf{10} \mathbf{5}_H)$ . It also forbids the dimension 3, lepton number violating, operator  $(\bar{\mathbf{5}} \mathbf{5}_H) \supset (L H_u)$  with a coefficient with dimensions of mass which, like the  $\mu$  parameter, could be of order the weak scale and the dimension 5, baryon number violating, operator  $(\mathbf{10} \mathbf{10} \mathbf{10} \bar{\mathbf{5}}_H) \supset (Q Q Q H_d) + \dots$ .

Note, in the MSSM it is possible to retain R parity violating operators at low energy as long as they violate either baryon number or lepton number only but not both. Such schemes are natural if one assumes a low energy symmetry, such as lepton number, baryon number or a baryon parity [41]. However these symmetries cannot be embedded in a GUT. Thus, in a SUSY GUT, only R parity can prevent unwanted dimension four operators. Hence, by naturalness arguments, R parity must be a symmetry in the effective low energy theory of any SUSY GUT. This does not mean to say that R parity is guaranteed to be satisfied in any GUT.

Note also, R parity distinguishes Higgs multiplets from ordinary families. In  $SU(5)$ , Higgs and quark/lepton multiplets have identical quantum numbers; while in  $E(6)$ , Higgs and families are unified within the fundamental  $\mathbf{27}$  representation. Only

in  $SO(10)$  are Higgs and ordinary families distinguished by their gauge quantum numbers. Moreover the  $Z(4)$  center of  $SO(10)$  distinguishes **10**s from **16**s and can be associated with R parity [42].

Dimension 5 baryon number violating operators may be forbidden at tree level by symmetries in  $SU(5)$ , etc. These symmetries are typically broken however by the VEVs responsible for the color triplet Higgs masses. Consequently these dimension 5 operators are generically generated via color triplet Higgsino exchange. Hence, the color triplet partners of Higgs doublets must necessarily obtain mass of order the GUT scale. The dominant decay modes from dimension 5 operators are  $p \rightarrow K^+ \bar{\nu}$  ( $n \rightarrow K^0 \bar{\nu}$ ). This is due to a simple symmetry argument; the operators  $(Q_i Q_j Q_k L_l)$ ,  $(u_i^c u_j^c d_k^c e_l^c)$  (where  $i, j, k, l = 1, 2, 3$  are family indices and color and weak indices are implicit) must be invariant under  $SU(3)_C$  and  $SU(2)_L$ . As a result their color and weak doublet indices must be anti-symmetrized. However since these operators are given by bosonic superfields, they must be totally symmetric under interchange of all indices. Thus the first operator vanishes for  $i = j = k$  and the second vanishes for  $i = j$ . Hence a second or third generation member must exist in the final state [40].

Recent Super-Kamiokande bounds on the proton lifetime severely constrain these dimension 6 and 5 operators with  $\tau_{(p \rightarrow e^+ \pi^0)} > 5.0 \times 10^{33}$  yrs (79.3 ktyr exposure),  $\tau_{(n \rightarrow e^+ \pi^-)} > 5 \times 10^{33}$  yrs (61 ktyr), and  $\tau_{(p \rightarrow K^+ \bar{\nu})} > 2.3 \times 10^{33}$  yrs (92 ktyr),  $\tau_{(n \rightarrow K^0 \bar{\nu})} > 1.3 \times 10^{32}$  yrs (92 ktyr) at (90% CL) based on the listed exposures [43]. These constraints are now sufficient to rule out minimal SUSY  $SU(5)$  [44].<sup>9</sup> Non-minimal Higgs sectors in  $SU(5)$  or  $SO(10)$  theories still survive [23, 30]. The upper bound on the proton lifetime from these theories are approximately a factor of 5 above the experimental bounds. They are also being pushed to their theoretical limits. Hence if SUSY GUTs are correct, then nucleon decay must be seen soon.

Is there a way out of this conclusion? Orbifold GUTs and string theories, see Sect. 1.3, contain grand unified symmetries realized in higher dimensions. In the process of compactification and GUT symmetry breaking, color triplet Higgs states are removed (projected out of the massless sector of the theory). In addition, the same projections typically rearrange the quark and lepton states so that the massless states which survive emanate from different GUT multiplets. In these models, proton decay due to dimension 5 operators can be severely suppressed or eliminated completely. However, proton decay due to dimension 6 operators may be enhanced, since the gauge bosons mediating proton decay obtain mass at the compactification scale,  $M_c$ , which is less than the 4d GUT scale (see the discussion at the end of Section 1.4), or suppressed, if the states of one family come from different irreducible representations. Which effect dominates is a model dependent issue. In some complete 5d orbifold GUT models [47, 24] the lifetime for the decay  $\tau(p \rightarrow e^+ \pi^0)$  can be near the excluded bound of  $5 \times 10^{33}$  years with, however, large model dependent and/or theoretical uncertainties. In other

---

<sup>9</sup>This conclusion relies on the mild assumption that the three-by-three matrices diagonalizing squark and slepton mass matrices are not so different from their fermionic partners. It has been shown that if this caveat is violated, then dimension five proton decay in minimal SUSY  $SU(5)$  may not be ruled out [45]. This is however a very fine-tuned resolution of the problem. Another possible way out is to allow for a more complicated  $SU(5)$  breaking Higgs sector in the otherwise minimal model [46].

cases, the modes  $p \rightarrow K^+ \bar{\nu}$  and  $p \rightarrow K^0 \mu^+$  may be dominant [24]. To summarize, in either 4d or orbifold string/field theories, nucleon decay remains a premier signature for SUSY GUTs. Moreover, the observation of nucleon decay may distinguish extra-dimensional orbifold GUTs from four dimensional ones.

Before concluding the topic of baryon number violation, consider the status of  $\Delta B = 2$  neutron- anti-neutron oscillations. Generically the leading operator for this process is the dimension 9 six quark operator  $G_{(\Delta B=2)} (u^c d^c d^c u^c d^c d^c)$  with dimensionful coefficient  $G_{(\Delta B=2)} \sim 1/M^5$ . The present experimental bound  $\tau_{n-\bar{n}} \geq 0.86 \times 10^8$  sec. at 90% CL [48] probes only up to the scale  $M \leq 10^6$  GeV. For  $M \sim M_G$ ,  $n - \bar{n}$  oscillations appear to be unobservable for any GUT (for a recent discussion see [49]).

## 1.6 Yukawa coupling unification

### 1.6.1 3rd generation, $b - \tau$ or $t - b - \tau$ unification

If quarks and leptons are two sides of the same coin, related by a new grand unified gauge symmetry, then that same symmetry relates the Yukawa couplings (and hence the masses) of quarks and leptons. In  $SU(5)$ , there are two independent renormalizable Yukawa interactions given by  $\lambda_t (\mathbf{10} \mathbf{10} \mathbf{5}_H) + \lambda (\mathbf{10} \mathbf{5} \mathbf{5}_H)$ . These contain the SM interactions  $\lambda_t (\mathbf{Q} \mathbf{u}^c \mathbf{H}_u) + \lambda (\mathbf{Q} \mathbf{d}^c \mathbf{H}_d + \mathbf{e}^c \mathbf{L} \mathbf{H}_d)$ . Hence, at the GUT scale we have the tree level relation,  $\lambda_b = \lambda_\tau \equiv \lambda$  [35]. In  $SO(10)$  there is only one independent renormalizable Yukawa interaction given by  $\lambda (\mathbf{16} \mathbf{16} \mathbf{10}_H)$  which gives the tree level relation,  $\lambda_t = \lambda_b = \lambda_\tau \equiv \lambda$  [50, 51]. Note, in the discussion above we assume the minimal Higgs content with Higgs in  $\mathbf{5}$ ,  $\bar{\mathbf{5}}$  for  $SU(5)$  and  $\mathbf{10}$  for  $SO(10)$ . With Higgs in higher dimensional representations there are more possible Yukawa couplings [59, 60, 61].

In order to make contact with the data, one now renormalizes the top, bottom and  $\tau$  Yukawa couplings, using two loop RG equations, from  $M_G$  to  $M_Z$ . One then obtains the running quark masses  $m_t(M_Z) = \lambda_t(M_Z) v_u$ ,  $m_b(M_Z) = \lambda_b(M_Z) v_d$  and  $m_\tau(M_Z) = \lambda_\tau(M_Z) v_d$  where  $\langle H_u^0 \rangle \equiv v_u = \sin \beta v / \sqrt{2}$ ,  $\langle H_d^0 \rangle \equiv v_d = \cos \beta v / \sqrt{2}$ ,  $v_u/v_d \equiv \tan \beta$  and  $v \sim 246$  GeV is fixed by the Fermi constant,  $G_\mu$ .

Including one loop threshold corrections at  $M_Z$  and additional RG running, one finds the top, bottom and  $\tau$  pole masses. In SUSY,  $b - \tau$  unification has two possible solutions with  $\tan \beta \sim 1$  or  $40 - 50$ . The small  $\tan \beta$  solution is now disfavored by the LEP limit,  $\tan \beta > 2.4$  [52].<sup>10</sup> The large  $\tan \beta$  limit overlaps the  $SO(10)$  symmetry relation.

When  $\tan \beta$  is large there are significant weak scale threshold corrections to down quark and charged lepton masses from either gluino and/or chargino loops [54]. Yukawa unification (consistent with low energy data) is only possible in a restricted region of SUSY parameter space with important consequences for SUSY searches [55].

<sup>10</sup>However, this bound disappears if one takes  $M_{SUSY} = 2$  TeV and  $m_t = 180$  GeV [53].

### 1.6.2 Three families

Simple Yukawa unification is not possible for the first two generations of quarks and leptons. Consider the  $SU(5)$  GUT scale relation  $\lambda_b = \lambda_\tau$ . If extended to the first two generations one would have  $\lambda_s = \lambda_\mu$ ,  $\lambda_d = \lambda_e$  which gives  $\lambda_s/\lambda_d = \lambda_\mu/\lambda_e$ . The last relation is a renormalization group invariant and is thus satisfied at any scale. In particular, at the weak scale one obtains  $m_s/m_d = m_\mu/m_e$  which is in serious disagreement with the data with  $m_s/m_d \sim 20$  and  $m_\mu/m_e \sim 200$ . An elegant solution to this problem was given by Georgi and Jarlskog [56]. Of course, a three family model must also give the observed CKM mixing in the quark sector. Note, although there are typically many more parameters in the GUT theory above  $M_G$ , it is possible to obtain effective low energy theories with many fewer parameters making strong predictions for quark and lepton masses.

It is important to note that grand unification alone is not sufficient to obtain predictive theories of fermion masses and mixing angles. Other ingredients are needed. In one approach additional global family symmetries are introduced (non-abelian family symmetries can significantly reduce the number of arbitrary parameters in the Yukawa matrices). These family symmetries constrain the set of effective higher dimensional fermion mass operators. In addition, sequential breaking of the family symmetry is correlated with the hierarchy of fermion masses. Three-family models exist which fit all the data, including neutrino masses and mixing [57]. In a completely separate approach for  $SO(10)$  models, the Standard Model Higgs bosons are contained in the higher dimensional Higgs representations including the **10**,  **$\overline{126}$**  and/or **120**. Such theories have been shown to make predictions for neutrino masses and mixing angles [59, 60, 61].

## 1.7 Neutrino Masses

Atmospheric and solar neutrino oscillations require neutrino masses. Adding three “sterile” neutrinos  $\nu^c$  with the Yukawa coupling  $\lambda_\nu$  ( $\nu^c \mathbf{L} \mathbf{H}_u$ ), one easily obtains three massive Dirac neutrinos with mass  $m_\nu = \lambda_\nu v_u$ .<sup>11</sup> However in order to obtain a tau neutrino with mass of order 0.1 eV, one needs  $\lambda_{\nu_\tau}/\lambda_\tau \leq 10^{-10}$ . The see-saw mechanism, on the other hand, can naturally explain such small neutrino masses [62, 63]. Since  $\nu^c$  has no SM quantum numbers, there is no symmetry (other than global lepton number) which prevents the mass term  $\frac{1}{2} \nu^c M \nu^c$ . Moreover one might expect  $M \sim M_G$ . Heavy “sterile” neutrinos can be integrated out of the theory, defining an effective low energy theory with only light active Majorana neutrinos with the effective dimension 5 operator  $\frac{1}{2} (\mathbf{L} \mathbf{H}_u) \lambda_\nu^T M^{-1} \lambda_\nu (\mathbf{L} \mathbf{H}_u)$ . This then leads to a  $3 \times 3$  Majorana neutrino mass matrix  $\mathbf{m} = m_\nu^T M^{-1} m_\nu$ .

Atmospheric neutrino oscillations require neutrino masses with  $\Delta m_\nu^2 \sim 3 \times 10^{-3}$  eV<sup>2</sup> with maximal mixing, in the simplest two neutrino scenario. With hierarchical neutrino masses  $m_{\nu_\tau} = \sqrt{\Delta m_\nu^2} \sim 0.055$  eV. Moreover via the “see-saw” mechanism  $m_{\nu_\tau} = m_t(m_t)^2/(3M)$ . Hence one finds  $M \sim 2 \times 10^{14}$  GeV; remarkably close to the GUT scale. Note we have related the neutrino Yukawa coupling to the top quark

---

<sup>11</sup>Note, these “sterile” neutrinos are quite naturally identified with the right-handed neutrinos necessarily contained in complete families of  $SO(10)$  or Pati-Salam.

Yukawa coupling  $\lambda_{\nu\tau} = \lambda_t$  at  $M_G$  as given in  $SO(10)$  or  $SU(4) \times SU(2)_L \times SU(2)_R$ . However at low energies they are no longer equal and we have estimated this RG effect by  $\lambda_{\nu\tau}(M_Z) \approx \lambda_t(M_Z)/\sqrt{3}$ .

## 1.8 Selected Topics

### 1.8.1 Magnetic Monopoles

In the broken phase of a GUT there are typically localized classical solutions carrying magnetic charge under an unbroken  $U(1)$  symmetry [64]. These magnetic monopoles with mass of order  $M_G/\alpha_G$  are produced during the GUT phase transition in the early universe. The flux of magnetic monopoles is experimentally found to be less than  $\sim 10^{-16} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$  [65]. Many more are however predicted, hence the GUT monopole problem. In fact, one of the original motivations for an inflationary universe is to solve the monopole problem by invoking an epoch of rapid inflation after the GUT phase transition [66]. This would have the effect of diluting the monopole density as long as the reheat temperature is sufficiently below  $M_G$ . Other possible solutions to the monopole problem include: sweeping them away by domain walls [67],  $U(1)$  electromagnetic symmetry breaking at high temperature [68] or GUT symmetry non-restoration [69]. Parenthetically, it was also shown that GUT monopoles can catalyze nucleon decay [70]. A significantly lower bound on the monopole flux can then be obtained by considering X-ray emission from radio pulsars due to monopole capture and the subsequent nucleon decay catalysis [71].

### 1.8.2 Baryogenesis via Leptogenesis

Baryon number violating operators in  $SU(5)$  or  $SO(10)$  preserve the global symmetry  $B - L$ . Hence the value of the cosmological  $B - L$  density is an initial condition of the theory and is typically assumed to be zero. On the other hand, anomalies of the electroweak symmetry violate  $B + L$  while also preserving  $B - L$ . Hence thermal fluctuations in the early universe, via so-called sphaleron processes, can drive  $B + L$  to zero, washing out any net baryon number generated in the early universe at GUT temperatures.

One way out of this dilemma is to generate a net  $B - L$  dynamically in the early universe. We have just seen that neutrino oscillations suggest a new scale of physics of order  $10^{14}$  GeV. This scale is associated with heavy Majorana neutrinos with mass  $M$ . If in the early universe, the decay of the heavy neutrinos is out of equilibrium and violates both lepton number and CP, then a net lepton number may be generated. This lepton number will then be partially converted into baryon number via electroweak processes [72].

### 1.8.3 GUT symmetry breaking

The grand unification symmetry is necessarily broken spontaneously. Scalar potentials (or superpotentials) exist whose vacua spontaneously break  $SU(5)$  and  $SO(10)$ . These potentials are ad hoc (just like the Higgs potential in the SM) and therefore it is hoped that they may be replaced with better motivated sectors. Gauge coupling

unification now tests GUT breaking sectors, since it is one of the two dominant corrections to the GUT threshold correction  $\epsilon_3$ . The other dominant correction comes from the Higgs sector and doublet-triplet splitting. This latter contribution is always positive  $\epsilon_3 \propto \ln(M_T/M_G)$  (where  $M_T$  is an effective color triplet Higgs mass), while the low energy data requires  $\epsilon_3 < 0$ . Hence the GUT breaking sector must provide a significant (of order -8%) contribution to  $\epsilon_3$  to be consistent with the Super-K bound on the proton lifetime [29, 23, 30, 57].

In string theory (and GUTs in extra-dimensions), GUT breaking may occur due to boundary conditions in the compactified dimensions [7, 10]. This is still ad hoc. The major benefits are that it does not require complicated GUT breaking sectors.

#### 1.8.4 Doublet-triplet splitting

The minimal supersymmetric standard model has a  $\mu$  problem; why is the coefficient of the bilinear Higgs term in the superpotential  $\mu (\mathbf{H}_u \mathbf{H}_d)$  of order the weak scale when, since it violates no low energy symmetry, it could be as large as  $M_G$ . In a SUSY GUT, the  $\mu$  problem is replaced by the problem of *doublet-triplet* splitting — giving mass of order  $M_G$  to the color triplet Higgs and mass  $\mu$  to the Higgs doublets. Several mechanisms for natural doublet-triplet splitting have been suggested, such as the sliding singlet, missing partner or missing VEV [73], and pseudo-Nambu-Goldstone boson mechanisms. Particular examples of the missing partner mechanism for  $SU(5)$  [30], the missing VEV mechanism for  $SO(10)$  [57, 23] and the pseudo-Nambu-Goldstone boson mechanism for  $SU(6)$  [74] have been shown to be consistent with gauge coupling unification and proton decay. There are also several mechanisms for explaining why  $\mu$  is of order the SUSY breaking scale [75]. Finally, for a recent review of the  $\mu$  problem and some suggested solutions in SUSY GUTs and string theory, see Ref. [76, 9] and references therein.

Once again, in string theory (and orbifold GUTs), the act of breaking the GUT symmetry via orbifolding projects certain states out of the theory. It has been shown that it is possible to remove the color triplet Higgs while retaining the Higgs doublets in this process. Hence the doublet-triplet splitting problem is finessed. As discussed earlier (see Section 1.5), this has the effect of eliminating the contribution of dimension 5 operators to nucleon decay.

## 2 Conclusion

Grand unification of the strong and electroweak interactions requires that the three low energy gauge couplings unify (up to small threshold corrections) at a unique scale,  $M_G$ . Supersymmetric grand unified theories provide, by far, the most predictive and economical framework allowing for perturbative unification.

The three pillars of SUSY GUTs are:

- gauge coupling unification at  $M_G \sim 3 \times 10^{16}$  GeV;
- low energy supersymmetry [with a large SUSY desert], and
- nucleon decay.

The first prediction has already been verified (see Fig. 1). Perhaps the next two will be seen soon. Whether or not Yukawa couplings unify is more model dependent. Nevertheless, the “digital” 16 dimensional representation of quarks and leptons in  $SO(10)$  is very compelling and may yet lead to an understanding of fermion masses and mixing angles.

In any event, the experimental verification of the first three pillars of SUSY GUTs would forever change our view of Nature. Moreover, the concomitant evidence for a vast SUSY desert would expose a huge lever arm for discovery. For then it would become clear that experiments probing the TeV scale could reveal physics at the GUT scale and perhaps beyond. Of course, some questions will still remain: Why do we have three families of quarks and leptons? How is the grand unified symmetry and possible family symmetries chosen by Nature? At what scale might stringy physics become relevant? Etc.

## References

- [1] J. Pati and A. Salam, *Phys. Rev.* **D8** 1240 (1973). For more discussion on the standard charge assignments in this formalism, see A. Davidson, *Phys. Rev.***D20**, 776 (1979) and R.N. Mohapatra and R.E. Marshak, *Phys. Lett.***B91**, 222 (1980).
- [2] H. Georgi and S.L. Glashow, *Phys. Rev. Lett.* **32** 438 (1974).
- [3] H. Georgi, *Particles and Fields*, Proceedings of the APS Div. of Particles and Fields, ed C. Carlson, p. 575 (1975); H. Fritzsch and P. Minkowski, *Ann. Phys.* **93**, 193 (1975).
- [4] S.M. Barr, *Phys. Lett.* **B112**, 219 (1982).
- [5] A. de Rujula, H. Georgi and S.L. Glashow, p. 88, 5th Workshop on Grand Unification, edited by K. Kang, H. Fried and P. Frampton, World Scientific, Singapore (1984). See also earlier paper by Y. Achiman and B. Stech, p. 303, *New Phenomena in Lepton-Hadron Physics*, edited by D.E.C. Fries and J. Wess, Plenum, NY (1979).
- [6] B.R. Greene, K.H. Kirklin, P.J. Miron, G.G. Ross, *Nucl. Phys.* **B278**, 667 (1986), *Nucl. Phys.* **B292**, 606 (1987); B.R. Greene, C.A. Lutken, G.G. Ross, *Nucl. Phys.* **B325**, 101 (1989); J. E. Kim, *Phys. Lett. B* **591**, 119 (2004) [arXiv:hep-ph/0403196].
- [7] P. Candelas, G.T. Horowitz, A. Strominger, E. Witten, *Nucl. Phys.* **B258**, 46 (1985); L.J. Dixon, J.A. Harvey, C. Vafa, E. Witten, *Nucl. Phys.* **B261**, 678 (1985), *Nucl. Phys.* **B274**, 285 (1986); L. E. Ibanez, H. P. Nilles, F. Quevedo, *Phys. Lett.***B187**, 25 (1987); L. E. Ibanez, J. E. Kim, H. P. Nilles, F. Quevedo, *Phys. Lett.***B191**, 282 (1987).
- [8] T. Kobayashi, S. Raby, R.-J. Zhang, *Phys. Lett.***B593**, 262 (2004); S. Forste, H. P. Nilles, P. Vaudrevange, A. Wingerter, *Phys. Rev.***D70**, 106008 (2004);

- T. Kobayashi, S. Raby and R. J. Zhang, *Nucl. Phys.***B704**, 3 (2005); W. Buchmuller, K. Hamaguchi, O. Lebedev and M. Ratz, *Nucl. Phys.***B712**, 139 (2005).
- [9] E. Witten, arXiv:hep-ph/0201018; M. Dine, Y. Nir and Y. Shadmi, *Phys. Rev.* **D66**, 115001 (2002) [arXiv:hep-ph/0206268].
- [10] Y. Kawamura, *Progr. Theor. Phys.***103**, 613 (2000); *ibid.* **105**, 999 (2001); G. Altarelli, F. Feruglio, *Phys. Lett.***B511**, 257 (2001); L. J. Hall, Y. Nomura, *Phys. Rev.* **D64**, 055003 (2001); A. Hebecker, J. March-Russell, *Nucl. Phys.* **B613**, 3 (2001); T. Asaka, W. Buchmuller, L. Covi, *Phys. Lett.* **B523**, 199 (2001); L. J. Hall, Y. Nomura, T. Okui, D. R. Smith, *Phys. Rev.* **D65**, 035008 (2002); R. Dermisek and A. Mafi, *Phys. Rev.* **D65**, 055002 (2002); H. D. Kim and S. Raby, *JHEP* **0301**, 056 (2003).
- [11] For a recent review see, R. Blumenhagen, M. Cvetič, P. Langacker and G. Shiu, “Toward realistic intersecting D-brane models,” [arXiv:hep-th/0502005].
- [12] G. Aldazabal, A. Font, L. E. Ibanez and A. M. Uranga, *Nucl. Phys.***B452**, 3 (1995); Z. Kakushadze, S.H.H. Tye, *Phys. Rev.* **D54**, 7520 (1996); Z. Kakushadze, G. Shiu, S.H.H. Tye, Y. Vtorov-Karevsky, *Int. J. Mod. Phys.* **A13**, 2551 (1998).
- [13] G. B. Cleaver, A. E. Faraggi and D. V. Nanopoulos, *Int. J. Mod. Phys.* **A16**, 425 (2001) [arXiv:hep-ph/9904301]; G. B. Cleaver, A. E. Faraggi, D. V. Nanopoulos and J. W. Walker, *Nucl. Phys.* **B593**, 471 (2001) [arXiv:hep-ph/9910230].
- [14] V. Braun, Y. H. He, B. A. Ovrut and T. Pantev, *Phys. Lett.* **B618**, 252 (2005) [arXiv:hep-th/0501070]; V. Braun, Y. H. He, B. A. Ovrut and T. Pantev, *JHEP* **0506**, 039 (2005) [arXiv:hep-th/0502155].
- [15] E. Witten, *Nucl. Phys.* **B471**, 135 (1996) [arXiv:hep-th/9602070].
- [16] H. Georgi, H. Quinn and S. Weinberg, *Phys. Rev. Lett.* **33** 451 (1974); see also the definition of effective field theories by S. Weinberg, *Phys. Lett.* **91B** 51 (1980).
- [17] See talks on proposed and running nucleon decay experiments, and theoretical talks by P. Langacker, p. 131 and W.J. Marciano and A. Sirlin, p. 151, in The Second Workshop on Grand Unification, J.P. Leveille, L.R. Sulak and D.G. Unger, editors, Birkhäuser, Boston (1981).
- [18] W.J. Marciano, p. 190, Eighth Workshop on Grand Unification, K. Wali, editor, World Scientific Publishing Co., Singapore (1987).
- [19] U. Amaldi, W. de Boer and H. Fürstenau, *Phys. Lett.* **B260**, 447 (1991); J. Ellis, S. Kelly and D.V. Nanopoulos, *Phys. Lett.* **B260**, 131 (1991); P. Langacker and M. Luo, *Phys. Rev.* **D44**, 817 (1991); P. Langacker and N. Polonsky, *Phys. Rev.* **D47**, 4028 (1993); M. Carena, S. Pokorski and C.E.M. Wagner, *Nucl. Phys.* **B406**, 59 (1993); see also the review by S. Dimopoulos, S. Raby and F. Wilczek, *Physics Today*, 25–33, October (1991).



- [20] S. Dimopoulos, S. Raby and F. Wilczek, *Phys. Rev.* **D24**, 1681 (1981); S. Dimopoulos and H. Georgi, *Nucl. Phys.* **B193**, 150 (1981); L. Ibanez and G.G. Ross, *Phys. Lett.* **105B**, 439 (1981); N. Sakai, *Z. Phys.* **C11**, 153 (1981); M. B. Einhorn and D. R. T. Jones, *Nucl. Phys.* **B196**, 475 (1982); W. J. Marciano and G. Senjanovic, *Phys. Rev.* **D 25**, 3092 (1982).
- [21] D.I. Kazakov, Lectures given at the European School on High Energy Physics, Aug.-Sept. 2000, Caramulo, Portugal [hep-ph/0012288v2].
- [22] V. Lucas and S. Raby, *Phys. Rev.* **D54**, 2261 (1996) [arXiv:hep-ph/9601303]; T. Blazek, M. Carena, S. Raby and C. E. M. Wagner, *Phys. Rev.* **D56**, 6919 (1997) [arXiv:hep-ph/9611217]; G. Altarelli, F. Feruglio and I. Masina, *JHEP* **0011**, 040 (2000) [arXiv:hep-ph/0007254].
- [23] R. Dermíšek, A. Mafi and S. Raby, *Phys. Rev.* **D63**, 035001 (2001); K.S. Babu, J.C. Pati and F. Wilczek, *Nucl. Phys.* **B566**, 33 (2000).
- [24] M. L. Alciati, F. Feruglio, Y. Lin and A. Varagnolo, *JHEP* **0503**, 054 (2005) [arXiv:hep-ph/0501086].
- [25] G. Anderson, C. H. Chen, J. F. Gunion, J. Lykken, T. Moroi and Y. Yamada, eConf **C960625**, SUP107 (1996) [arXiv:hep-ph/9609457].
- [26] R.N. Mohapatra and M.K. Parida, *Phys. Rev.* **D47**, 264 (1993).
- [27] D.G. Lee, R.N. Mohapatra, M.K. Parida and M. Rani, *Phys. Rev.* **D51**, 229 (1995).
- [28] S. Eidelman *et al.* [Particle Data Group], *Phys. Lett.* **B592**, 1 (2004).
- [29] K.S. Babu and S.M. Barr, *Phys. Rev.* **D48**, 5354 (1993); V. Lucas and S. Raby, *Phys. Rev.* **D54**, 2261 (1996); S.M. Barr and S. Raby, *Phys. Rev. Lett.* **79**, 4748 (1997) and references therein.
- [30] G. Altarelli, F. Feruglio, I. Masina, *JHEP* **0011**, 040 (2000). See also earlier papers by A. Masiero, D.V. Nanopoulos, K. Tamvakis and T. Yanagida, *Phys. Lett.* **B115**, 380 (1982); B. Grinstein, *Nucl. Phys.* **B206**, 387 (1982).
- [31] K.R. Dienes, E. Dudas, T. Gherghetta, *Phys. Rev. Lett.* **91**, 061601 (2003).
- [32] L. J. Hall, Y. Nomura, *Phys. Rev.* **D64**, 055003 (2001).
- [33] N. Arkani-Hamed and S. Dimopoulos, *JHEP* **0506**, 073 (2005) [arXiv:hep-th/0405159].
- [34] G. F. Giudice and A. Romanino, *Nucl. Phys.* **B699**, 65 (2004) [Erratum-ibid. **B 706**, 65 (2005)] [arXiv:hep-ph/0406088].

- [35] M. Chanowitz, J. Ellis and M.K. Gaillard, *Nucl. Phys.* **B135**, 66 (1978). For the corresponding SUSY analysis, see M. Einhorn and D.R.T. Jones, *Nucl. Phys.* **B196**, 475 (1982); K. Inoue, A. Kakuto, H. Komatsu and S. Takeshita, *Prog. Theor. Phys.* **67**, 1889 (1982); L. E. Ibanez and C. Lopez, *Phys. Lett.* **B126**, 54 (1983); *Nucl. Phys.* **B233**, 511 (1984).
- [36] K. Agashe, R. Contino and R. Sundrum, [arXiv:hep-ph/0502222].
- [37] M. Gell-Mann, P. Ramond and R. Slansky, in *Supergravity*, ed. P. van Nieuwenhuizen and D.Z. Freedman, North-Holland, Amsterdam, 1979, p. 315.
- [38] S. Weinberg, *Phys. Rev.* **D26**, 287 (1982); N. Sakai and T Yanagida, *Nucl. Phys.* **B197**, 533 (1982).
- [39] G. Farrar and P. Fayet, *Phys. Lett.* **B76**, 575 (1978).
- [40] S. Dimopoulos, S. Raby and F. Wilczek, *Phys. Lett.* **112B**, 133 (1982); J. Ellis, D.V. Nanopoulos and S. Rudaz, *Nucl. Phys.* **B202**, 43 (1982).
- [41] L.E. Ibanez and G.G. Ross, *Nucl. Phys.* **B368**, 3 (1992).
- [42] For a recent discussion, see C.S. Aulakh, B. Bajc, A. Melfo, A. Rasin and G. Senjanovic, *Nucl. Phys.* **B597**, 89 (2001).
- [43] See talks by Matthew Earl, NNN workshop, Irvine, February (2000); Y. Totsumura, SUSY2K, CERN, June (2000); Y. Suzuki, International Workshop on Neutrino Oscillations and their Origins, Tokyo, Japan, December (2000) and Baksan School, Baksan Valley, Russia, April (2001), hep-ex/0110005; K. Kobayashi [Super-Kamiokande Collaboration], "Search for nucleon decay from Super-Kamiokande," *Prepared for 27th International Cosmic Ray Conference (ICRC 2001), Hamburg, Germany, 7-15 Aug 2001*. For published results see Super-Kamiokande Collaboration: Y. Hayato, M. Earl, et. al, *Phys. Rev. Lett.* **83**, 1529 (1999); K. Kobayashi *et al.* [Super-Kamiokande Collaboration], [arXiv:hep-ex/0502026].
- [44] T. Goto and T. Nihei, *Phys. Rev.* **D59**, 115009 (1999) [arXiv:hep-ph/9808255]; H. Murayama and A. Pierce, *Phys. Rev.* **D65**, 055009 (2002) [arXiv:hep-ph/0108104].
- [45] B. Bajc, P. Fileviez Perez and G. Senjanovic, *Phys. Rev.* **D66**, 075005 (2002) [arXiv:hep-ph/0204311].
- [46] J. L. Chkareuli and I. G. Gogoladze, *Phys. Rev.* **D58**, 055011 (1998) [arXiv:hep-ph/9803335].
- [47] L. J. Hall and Y. Nomura, *Phys. Rev.* **D66**, 075004 (2002); H. D. Kim, S. Raby and L. Schradin, *JHEP* **0505**, 036 (2005).
- [48] M. Baldoceolin et al., *Zeit. fur Phys.* **C63**, 409 (1994).

- [49] K. S. Babu and R. N. Mohapatra, *Phys. Lett.* **B518**, 269 (2001) [arXiv:hep-ph/0108089].
- [50] H. Georgi and D.V. Nanopoulos, *Nucl. Phys.* **B159**, 16 (1979); J. Harvey, P. Ramond and D.B. Reiss, *Phys. Lett.* **92B**, 309 (1980); *Nucl. Phys.* **B199**, 223 (1982).
- [51] T. Banks, *Nucl. Phys.* **B303**, 172 (1988); M. Olechowski and S. Pokorski, *Phys. Lett.* **B214**, 393 (1988); S. Pokorski, *Nucl. Phys.* **B13** (Proc. Supp.), 606 (1990); B. Ananthanarayan, G. Lazarides and Q. Shafi, *Phys. Rev.* **D44**, 1613 (1991); Q. Shafi and B. Ananthanarayan, ICTP Summer School lectures (1991); S. Dimopoulos, L.J. Hall and S. Raby, *Phys. Rev. Lett.* **68**, 1984 (1992), *Phys. Rev.* **D45**, 4192 (1992); G. Anderson et al., *Phys. Rev.* **D47**, 3702 (1993); B. Ananthanarayan, G. Lazarides and Q. Shafi, *Phys. Lett.* **B300**, 245 (1993); G. Anderson et al., *Phys. Rev.* **D49**, 3660 (1994); B. Ananthanarayan, Q. Shafi and X.M. Wang, *Phys. Rev.* **D50**, 5980 (1994).
- [52] LEP Higgs Working Group and ALEPH collaboration and DELPHI collaboration and L3 collaboration and OPAL Collaboration, Preliminary results, [hep-ex/0107030] (2001).
- [53] M. Carena and H. E. Haber, *Prog. Part. Nucl. Phys.* **50**, 63 (2003) [arXiv:hep-ph/0208209].
- [54] L.J. Hall, R. Rattazzi and U. Sarid, *Phys. Rev.* **D50**, 7048 (1994); M. Carena, M. Olechowski, S. Pokorski and C.E.M. Wagner, *Nucl. Phys.* **B419**, 213 (1994); R. Rattazzi and U. Sarid, *Nucl. Phys.* **B501**, 297 (1997).
- [55] T. Blazek, R. Dermisek and S. Raby, *Phys. Rev. Lett.* **88**, 111804 (2002) [arXiv:hep-ph/0107097]; *Phys. Rev.* **D65**, 115004 (2002) [arXiv:hep-ph/0201081]; K. Tobe and J. D. Wells, *Nucl. Phys.* **B663**, 123 (2003) [arXiv:hep-ph/0301015]; D. Auto, H. Baer, C. Balazs, A. Belyaev, J. Ferrandis and X. Tata, *JHEP* **0306**, 023 (2003) [arXiv:hep-ph/0302155].
- [56] H. Georgi and C. Jarlskog, *Phys. Lett.* **86B** 297 (1979).
- [57] K.S. Babu and R.N. Mohapatra, *Phys. Rev. Lett.* **74**, 2418 (1995); V. Lucas and S. Raby, *Phys. Rev.* **D54**, 2261 (1996); T. Blažek, M. Carena, S. Raby and C. Wagner, *Phys. Rev.* **D56**, 6919 (1997); R. Barbieri, L.J. Hall, S. Raby and A. Romanino, *Nucl. Phys.* **B493**, 3 (1997); T. Blazek, S. Raby and K. Tobe, *Phys. Rev.* **D60**, 113001 (1999), *Phys. Rev.* **D62** 055001 (2000); Q. Shafi and Z. Tavartkiladze, *Phys. Lett.* **B487**, 145 (2000); C.H. Albright and S.M. Barr, *Phys. Rev. Lett.* **85**, 244 (2000); K.S. Babu, J.C. Pati and F. Wilczek, *Nucl. Phys.* **B566**, 33 (2000); G. Altarelli, F. Feruglio, I. Masina, Ref. [30]; Z. Berezhiani and A. Rossi, *Nucl. Phys.* **B594**, 113 (2001); C. H. Albright and S. M. Barr, *Phys. Rev.* **D64**, 073010 (2001) [arXiv:hep-ph/0104294];
- [58] R. Dermisek and S. Raby, *Phys. Lett.* **B622**, 327 (2005) [arXiv:hep-ph/0507045].

- [59] G. Lazarides, Q. Shafi and C. Wetterich, *Nucl. Phys.* **B181**, 287 (1981); T. E. Clark, T. K. Kuo and N. Nakagawa, *Phys. Lett.* **B115**, 26 (1982); K. S. Babu and R. N. Mohapatra, *Phys. Rev. Lett.* **70**, 2845 (1993) [arXiv:hep-ph/9209215].
- [60] B. Bajc, G. Senjanovic and F. Vissani, *Phys. Rev. Lett.* **90**, 051802 (2003) [arXiv:hep-ph/0210207].
- [61] H. S. Goh, R. N. Mohapatra and S. P. Ng, *Phys. Lett.* **B570**, 215 (2003) [arXiv:hep-ph/0303055]; H. S. Goh, R. N. Mohapatra and S. P. Ng, *Phys. Rev.* **D68**, 115008 (2003) [arXiv:hep-ph/0308197]; B. Dutta, Y. Mimura and R. N. Mohapatra, *Phys. Rev.* **D69**, 115014 (2004) [arXiv:hep-ph/0402113]; S. Bertolini and M. Malinsky, [arXiv:hep-ph/0504241]; K. S. Babu and C. Macesanu, [arXiv:hep-ph/0505200].
- [62] P. Minkowski, *Phys. Lett.* **B67**, 421 (1977).
- [63] T. Yanagida, in Proceedings of the Workshop on the unified theory and the baryon number of the universe, ed. O. Sawada and A. Sugamoto, KEK report No. 79-18, Tsukuba, Japan, 1979; S. Glashow, Quarks and leptons, published in Proceedings of the Cargèse Lectures, M. Levy (ed.), Plenum Press, New York, (1980); M. Gell-Mann, P. Ramond and R. Slansky, in Supergravity, ed. P. van Nieuwenhuizen and D.Z. Freedman, North-Holland, Amsterdam, (1979), p. 315; R.N. Mohapatra and G. Senjanovic, *Phys. Rev. Lett.* **44**, 912 (1980).
- [64] G. 't Hooft, *Nucl. Phys.* **B79**, 276 (1974); A.M. Polyakov, *Pis'ma Zh. Eksp. Teor. Fiz.* **20**, 430 (1974) [*JETP Lett.* **20**, 194 (1974)]. For a pedagogical introduction, see S. Coleman, in *Aspects of Symmetry*, Selected Erice Lectures, Cambridge University Press, Cambridge, (1985) and P. Goddard and D. Olive, *Rep. Prog. Phys.* **41**, 1357 (1978).
- [65] I. De Mitri, MACRO Collaboration, *Nucl. Phys.* (Proc. Suppl.), **B95**, 82 (2001).
- [66] For a review, see A.D. Linde, *Particle Physics and Inflationary Cosmology*, Harwood Academic, Switzerland (1990).
- [67] G. R. Dvali, H. Liu and T. Vachaspati, *Phys. Rev. Lett.* **80**, 2281 (1998) [arXiv:hep-ph/9710301].
- [68] P. Langacker and S. Y. Pi, *Phys. Rev. Lett.* **45**, 1 (1980).
- [69] G. R. Dvali, A. Melfo and G. Senjanovic, *Phys. Rev. Lett.* **75**, 4559 (1995) [arXiv:hep-ph/9507230].
- [70] V. Rubakov, *Nucl. Phys.* **B203**, 311 (1982), Institute of Nuclear Research Report No. P-0211, Moscow (1981), unpublished; C. Callan, *Phys. Rev.* **D26**, 2058 (1982); F. Wilczek, *Phys. Rev. Lett.* **48**, 1146 (1982). See also, S. Dawson and A.N. Schellekens, *Phys. Rev.* **D27**, 2119 (1983).
- [71] K. Freese, M. S. Turner and D. N. Schramm, *Phys. Rev. Lett.* **51**, 1625 (1983).

- [72] M. Fukugita and T. Yanagida, *Phys. Lett.* **B174**, 45 (1986); see also the recent review by W. Buchmuller, R. D. Peccei and T. Yanagida, [arXiv:hep-ph/0502169] and references therein.
- [73] S. Dimopoulos and F. Wilczek, Proceedings Erice Summer School, Ed. A. Zichichi (1981); K.S. Babu and S.M. Barr, *Phys. Rev.* **D50**, 3529 (1994).
- [74] R. Barbieri, G.R. Dvali, A. Strumia, *Nucl. Phys.* **B391**, 487 (1993); Z. Berezhiani, C. Csaki, and L. Randall, *Nucl. Phys.* **B444**, 61 (1995); Q. Shafi, Z. Tavartkiladze, *Phys. Lett.* **B522**, 102 (2001).
- [75] G.F. Giudice and A. Masiero, *Phys. Lett.* **B206**, 480 (1988); J.E. Kim and H.P. Nilles, *Mod. Phys. Lett.* **A9**, 3575 (1994).
- [76] L. Randall and C. Csaki, [arXiv:hep-ph/9508208].