

Using Poisson mixed-effects model to quantify transcript-level gene expression in RNA-Seq

Ming Hu¹, Yu Zhu^{2,*}, Jeremy M G Taylor³, Jun S Liu¹ and Zhaohui Qin^{4,*}

¹Department of Statistics, Harvard University, Cambridge, MA 02138.

²Department of Statistics, Purdue University, West Lafayette, IN 47907.

³Department of Biostatistics, University of Michigan, Ann Arbor, MI 48109

⁴Department of Biostatistics and Bioinformatics, Emory University, Atlanta, GA 30322

Associate Editor: Prof. Ivo Hofacker

ABSTRACT

Motivation: RNA sequencing (RNA-Seq) is a powerful new technology for mapping and quantifying transcriptomes using ultra high-throughput next generation sequencing technologies. Using deep sequencing, gene expression levels of all transcripts including novel ones can be quantified digitally. Although extremely promising, the massive amounts of data generated by RNA-Seq, substantial biases, and uncertainty in short read alignment pose challenges for data analysis. In particular, large base-specific variation and between-base dependence make simple approaches, such as those that use averaging to normalize RNA-Seq data and quantify gene expressions, ineffective.

Results: In this study, we propose a Poisson mixed-effects (or in short, POME) model to characterize base-level read coverage within each transcript. The underlying expression level is included as a key parameter in this model. Because the proposed model is capable of incorporating base-specific variation as well as between-base dependence that affect read coverage profile throughout the transcript, it can lead to improved quantification of the true underlying expression level.

Availability and Implementation: POME can be freely downloaded at <http://www.stat.purdue.edu/~yuzhu/pome.html>.

Contact: yuzhu@purdue.edu or zhaohui.qin@emory.edu.

Supplementary information: Supplementary data are available at *Bioinformatics* online.

*To whom correspondence should be addressed.

1 INTRODUCTION

The transcriptome is the complete set of transcripts in a cell under any given developmental stage or physiological condition. Comprehensively cataloging all the components in the transcriptome is a grand challenge in the post-genome era. In the past decade, microarray technology has played a prominent role in advancing our understanding of transcriptome complexity by allowing scientists to simultaneously monitor the expression of almost all the genes in the genome (Lockhart, et al., 1996; Schena, et al., 1995). Despite its overwhelming success, microarray technology has its limitations. For example, designing probes on the chip requires

knowledge of the genome sequence and annotation; hence novel transcripts will be missed. Additionally, cross-hybridization, background signal and saturation result in a reduction of microarray's dynamic range and accuracy.

A recently developed sequencing-based technology for measuring gene expression, termed RNA-Seq, has the potential to overcome these limitations (Mortazavi, et al., 2008; Wang, et al., 2009). The ultra-high-throughput next generation sequencing technologies capable of producing millions of sequence reads dramatically increase the throughput in DNA sequencing compared to conventional Sanger technology and at a much lower cost. An array of studies has been published that successfully apply these new sequencing technologies to measure mRNA expression levels in cells from various species (Cloonan, et al., 2008; Lister, et al., 2008; Maher, et al., 2009; Marioni, et al., 2008; Morin, et al., 2008; Mortazavi, et al., 2008; Nagalakshmi, et al., 2008; Prensner, et al., 2011; Trapnell, et al., 2010; Wilhelm, et al., 2008). Since Illumina's platforms have been widely adopted in RNA-Seq experiments, we focused on short read RNA-Seq data generated from the Illumina platform in this study.

In RNA-Seq experiments using the aforementioned sequencing technologies, RNA molecules are first converted to a library of cDNA fragments with adaptors attached to both ends. Each molecule, often after amplification, is then sequenced using one of the next generation sequencing technologies. Following sequencing, the resulting reads are aligned to known transcripts or *de novo* assembled together to produce a genome-scale transcriptional profile.

A fundamental question in RNA-Seq data analysis is to derive expression level from raw sequencing output data. This is the basis of almost all further investigation such as detecting differentially expressed genes, alternative splicing events, etc. Sophisticated and tailor-made data analysis methods are needed to fully realize the power of the new sequencing technologies.

A natural idea is to use the total number of reads mapped inside each transcript, or region of interest (ROI), to represent the expression level. Mortazavi et al. proposed to use the number of reads per kilobase of a transcript per million mapped reads (RPKM) as the transcript's expression level (Mortazavi, et al., 2008). The RPKM

method is easy to implement and takes into consideration the transcript length and the total number of uniquely mapped reads. However, the RPKM method is oversimplified, as it ignores the variability of read coverage within a ROI demonstrated by real RNA-Seq data. More sophisticated methods are required to account for the complexity and uncertainties associated with read mapping and read depth within a ROI. In that regard, model-based methods can potentially improve upon the RPKM method by explicitly modeling the varying sequencing read coverage within a ROI. A number of models have been proposed in the literature, such as Poisson model (Marioni, et al., 2008), generalized Poisson model (Srivastava and Chen, 2010) and negative binomial model (Bullard, et al., 2010). However, these models are not sophisticated enough to capture all the variability demonstrated in the observed RNA-Seq data. For example, they do not consider variation in sequencing read coverage within a ROI resulted from local genomic feature and fluctuation in the base-level PCR amplification rate. To accommodate this type of variation, Li et al. proposed to use Poisson distribution with varying intensity rate to model read counts covering different positions in a ROI and developed a method called mseq to quantify its expression level (Li, et al., 2010). The mseq method utilizes neighborhood sequence information and can better explain base-level read counts variation in RNA-Seq data.

Despite improvement over other existing methods, the mseq method assumes that the observed base-level read counts are stochastically independent with each other in a ROI. Analyzing real data indicated that this assumption is not valid in a substantial proportion of the ROIs. We found that in highly expressed transcripts, between-base or spatial dependence is not negligible (see Results). Ignoring this dependence may lead to less accurate estimation of the true expression levels.

The presence of location-specific variation along with between-location correlation is a outstanding characteristic of many spatial data generated in geostatistics, spatial epidemiology, and image processing and has been studied in the literature of spatial statistics (Best, et al., 2005; Diggle, et al., 1998; Wakefield, 2007). Typically, *Poisson mixed-effects models* are used to analyze such spatial count data. In this study, in order to model base-specific read coverage while accounting for their dependency simultaneously, we apply the spatial Poisson mixed-effects model to characterize transcript level RNA-Seq data.

2 METHODS

2.1 Poisson mixed-effects model

Let Y_{jk} represent the number of reads whose mapping starts at the j th base of a specified transcript in the k th sample. Here $j = 1, \dots, n$, n is the length of the transcript, and $k = 1, \dots, m$, m is the number of samples. The definition of Y_{jk} is the same as in mseq (Li, et al., 2010). Further let θ_k represent the expression index of the transcript in the k th sample, which is of primary interest. Our goal is to build a statistical model to capture the base-specific variation and between-base correlation in Y_{jk} . This idea is motivated by the model-based expression index

(MBEI) model proposed by Li and Wong to model probe-level microarray gene expression data (Li and Wong, 2001).

In order to avoid over-fitting using a complex model, we propose the following Poisson mixed-effects (or in short, POME) model for Y_{jk} :

$$Y_{jk} | \theta_k, U_{jk}, V_{jk} \sim \text{Poisson}(n\theta_k \exp\{U_{jk} + V_{jk}\}) \quad (1)$$

In addition to the fixed effect θ_k , which is the expression index that is of primary interest, there are two random effects terms in this model: U_{jk} 's and V_{jk} 's. As in spatial statistics, V_{jk} 's are assumed to be independent and identically distributed as $N(0, \sigma_v^2)$, and used to account for unstructured variability, which may be attributed to some latent factors for over-dispersion; U_{jk} 's are used to represent the correlation between the read counts of different base pairs. Like in spatial statistics, there are various ways to specify this correlation structure. In POME, we chose the intrinsic conditional autoregressive (ICAR) structure (Besag, 1974). The ICAR structure specifies between-base correlation using a Gaussian Markov random field. Originally, the ICAR structure was proposed for image processing and was later used for disease mapping in spatial epidemiology (Clayton and Kaldor, 1987). For a fixed base j , first we define its neighborhood, denoted by ∂j as the collection of the two bases $j-1$ and $j+1$ that are adjacent to base j . Other definitions of neighborhood are possible, for example, bases that are not immediately adjacent to base j can be included in ∂j (Cressie and Chan, 1989). Second, we define a weight matrix $W = (w_{ij})$ as follows. For $1 \leq i, j \leq n$, $w_{ii} = 0$; $w_{ij} = 1$ if $i \in \partial j$; and $w_{ij} = 0$ otherwise. Let $U_{(-j)k}$ denote the collection of U_{ik} 's with $i \neq j$. The conditional distribution of U_{jk} given $U_{(-j)k}$ is assumed to be:

$$U_{jk} | U_{(-j)k} \sim N\left(\frac{\sum_{i \in \partial j} w_{ij} U_{ik}}{\sum_{i \in \partial j} w_{ij}}, \frac{\sigma_u^2}{\sum_{i \in \partial j} w_{ij}}\right) \quad (2)$$

The ICAR structure induces correlation between different bases.

Another option for the correlation structure is the joint structure (Wakefield, 2007). The joint structure assumes that in the k th sample, U_{jk} 's follow the multivariate Gaussian distribution $N(0, \sigma_u^2 \Sigma)$, where $\Sigma = (\sigma_{ij})$ is the correlation matrix of U_{jk} 's and $\sigma_{ij} = \rho^{d_{ij}}$, $0 < \rho < 1$, d_{ij} is the distance or the number of base pairs between the i th base and the j th base, i.e., $d_{ij} = |i - j|$. When the ICAR structure in POME is replaced by

the joint structure, the resulting model is referred to as the joint model.

The discussion about the pros and cons of the joint versus ICAR structure can be found in the disease mapping literature (Best, et al., 2005; Wakefield, 2007). We choose the ICAR structure in POME because read coverage for some transcripts may be sparse and the excessive zero counts pose a challenge for the joint model. POME is flexible in that covariates or deterministic patterns that affect θ_k can be incorporated into the POME model in a straightforward fashion.

In the literature on disease mapping, Markov chain Monte Carlo (MCMC) techniques (Gilks, et al., 1998; Liu, 2001) are the predominant methods used for fitting the POME model and performing subsequent statistical inference, following (Besag, et al., 1991) and (Diggle, et al., 1998). In the literature on image processing, however, maximum likelihood methods are also used for model fitting and inference; see (Zhu, et al., 2009) for example. The preference for Bayesian computational methods is due to the fact that the random effects U_{jk} 's and V_{jk} 's are not directly observable; and it takes high-dimensional integration to integrate them out, which can be computationally challenging. In this study, following the tradition in disease mapping, we use Bayesian computational methods when applying the POME model for transcript level RNA-Seq data analysis.

2.2 Model implementation

The MCMC methods are employed to carry out the fitting of the POME model and subsequent statistical inference. We start from assigning appropriate priors for the model parameters. The marginal distributions of the random effects U_{jk} and V_{jk} are $N(0, \sigma_u^2)$ and $N(0, \sigma_v^2)$, respectively. Following the approach by Wakefield (Wakefield, 2007), we define the total precision as $\tau = (\sigma_u^2 + \sigma_v^2)^{-1}$, and specify a Gamma prior for it, which is $\tau \sim \Gamma(a, b)$. Let $p = \sigma_u^2 / (\sigma_u^2 + \sigma_v^2)$ represent the proportion of base-specific variation in the total variation. We assign a Beta prior $Beta(c, d)$ to p .

If the joint structure is assumed for U_{jk} 's, there is another parameter ρ determining the extent of between-bases correlation. We assign another Beta prior $Beta(e, f)$ to ρ . A non-informative prior $I(0, +\infty)$ is assigned for θ_k .

As the default, we specify $c = d = e = f = 1$ so that the priors for p and ρ are uniform distributions on $[0, 1]$. The total precision $\tau = (\sigma_u^2 + \sigma_v^2)^{-1}$ plays a crucial role in the model and the final result is sensitive to the prior assigned to τ . We adopt an empirical Bayesian method to further specify a and b in the prior $\Gamma(a, b)$ for τ by setting $a = 1$ and $b = \max_{1 \leq k \leq m} \hat{\tau}_k$ (Supplementary Material Section 1).

It is straightforward to derive the joint posterior distribution involving all the fixed and random effects or parameters (Supplementary Material Section 2), and we use Gibbs sampler to iteratively sample parameters from the conditional posterior distribu-

tions. Since we use the conjugate priors, the conditional posterior distributions for θ_k and τ are Gamma distributions, which are easy to sample from. The conditional posterior distributions for random effects U_{jk} 's and V_{jk} 's are rather complicated and not in closed forms. Since both are log-concave functions, we use the adaptive rejection sampling (ARS) method (Gilks and Wild, 1992) to draw samples from them. For the other parameters, p and ρ , their conditional posterior distributions are nearly log-concave. We use the adaptive rejection Metropolis-Hasting sampling (ARMS) technique (Gilks, et al., 1995) to draw samples from these complicated distributions. One important issue in using ARS and ARMS is to assign appropriate ranges for the parameters. In ARS, we use the interval $[-5, 5]$ for random effects U_{jk} 's and V_{jk} 's. For the parameters p and ρ , we use their natural range $[0, 1]$ in ARMS. In addition, to make the fixed parameter θ_k identifiable, we impose the following two constraints on the random effects in each iteration of the Gibbs sampler:

$$\sum_{j=1}^n U_{jk} = 0, \sum_{j=1}^n V_{jk} = 0, 1 \leq k \leq m \quad (3)$$

When fitting the POME model, we ran 10,000 MCMC iterations for each transcript. The first 9,000 samplers were dropped as the burn-in stage, and then every 10th sample in the last 1,000 samplers were used to calculate the posterior means.

3 RESULTS

3.1 Data description

To evaluate the performance of POME, we choose a published real RNA-Seq data, obtained from twelve prostate cancer cell lines and tissues (Sam, et al., 2011). More details about the samples can be found in Table S1 in the Supplementary Material. A brief description of the reads mapping procedure used by Sam, et al., 2011 is provided in the Supplementary Material Section 3. We choose this dataset because these samples were profiled using two types of sequencing instruments with different technologies: Illumina Genome Analyzer using sequencing by synthesis technology and Helicos HeliScope using single-molecule sequencing technology. The Helicos technology represents a new wave of next generation sequencing technologies in which samples were profiled directly without the polymerase chain reaction (PCR) step, thus eliminating the overrepresentation or underrepresentation biases introduced by the copying process of PCR, a necessary step in Illumina sequencing technology. Given this, we use the expression measure obtained from Helicos as the gold standard when evaluating expression measures reported by different algorithms analyzing RNA-Seq data generated by the Illumina platform.

Transcript-level expression measures from Helicos and mapped Illumina reads information are kindly provided by the Chinnaiyan Lab. The details of the involved data processing procedures can be found in Sam et al. 2011. We plotted log 2 mean versus log 2 variance of the Illumina read coverage of these transcripts in all twelve cell lines (Figure S1). We observed that the variance of read count

is much larger than the mean, indicating the presence of substantial over-dispersion, especially for highly expressed transcripts. This is consistent with the observation of Li et al. (Li, et al., 2010). Furthermore, when we selected the most highly expressed transcripts (top 500 according to the gold standard Helicos measures, of which 439 also satisfied the minimum read coverage criterion in Illumina (>0.3 RPKM noise level used in the Sam et al. study) and minimum effective length criterion (with more than 100 non-zero Y_{jk} 's)). When examining these transcripts in the LnCaP_0 sample (Supplementary Material Table S1), we found that the median of the lag one autocorrelations between base-level read counts of these transcripts is 0.11, the 3rd quartile is 0.18, and the maximum correlation is 0.57. We checked on other samples and observed similar patterns. These findings indicate that dependence between read counts of adjacent base pairs is real and needs to be considered.

3.2 Simulation study

3.2.1 Simulated data from the joint POME model

We conducted a simulation study to compare the performance of the POME method against the commonly used methods in quantifying transcript-level gene expression: RPKM (Mortazavi, et al., 2008), mseq (Li, et al., 2010) and GPseq (Srivastava and Chen, 2010).

We simulated the read coverage profiles from the spatial Poisson mixed effects model with the joint correlation structure, i.e., the joint model which is different from POME. To make our simulation study more realistic, we first used the joint model to fit observed read count data in the 439 transcripts of high expression levels. We then sampled putative sequencing coverage profiles for these transcripts from the joint model. The proportion of base-specific variation in the total variation p was drawn randomly from a uniform distribution defined on the interval (0.1, 0.9).

The simulation was repeated 100 times. For each dataset, we applied RPKM, mseq, GPseq and POME, and calculated the mean square errors (MSE) of the four resulting estimates of expression index θ .

For all 439 transcripts, the POME method achieved the smallest MSE (1.2024, standard error 0.0379). The MSEs reported by RPKM, mseq and GPseq were much larger: 65.4605 (standard error 4.2217), 87.1559 (standard error 23.3511) and 16.0718 (standard error 0.0692), respectively. The RPKM method overestimated the true expression level when data showed strong over-dispersion. On the other hand, mseq used 40 base pairs in the neighborhood of each nucleotide as local sequence features that affect the base-level read coverage rate, which may have caused it to over fit the data. Although GPseq is capable of modeling both over-dispersion and under-dispersion patterns in the data, it does not take into account the spatial dependence between adjacent base pairs which may result in less accurate estimates.

3.2.2 Simulated data from the generalized Poisson model

Next we conducted another simulation study with a different simulation strategy. We simulated the read coverage data from the generalized Poisson distribution $GP(\theta, \lambda)$. We again used the 439 highly expressed transcripts in the previous simulation study, and fitted a generalized Poisson model using GPseq (Srivastava and Chen, 2010) to obtain the empirical estimates $\hat{\theta}$ and $\hat{\lambda}$ in each transcript.

We then simulated the read coverage profiles from $GP(\hat{\theta}, \hat{\lambda})$ for all 439 transcripts. We applied the RPKM method, mseq and the POME method to estimate θ , separately. We did not apply GPseq in this simulation study since the data was simulated from the generalized Poisson distribution. The simulation was repeated 100 times, and the mean square errors (MSEs) of the three resulting estimates of θ were calculated.

For all 439 transcripts, the MSEs achieved by POME, mseq and RPKM were 20.6736 (standard error 0.4633), 160.7138 (standard error 33.5309) and 166.1414 (standard error 7.9313), respectively. The results were similar to what we obtained in the previous simulation study. Although the data was simulated from generalized Poisson distribution with over-dispersion, POME nevertheless was able to provide more accurate estimate of θ through explicitly modeling the position specific variation.

3.3 Real data analysis

We next analyzed real RNA-Seq data in the twelve prostate cancer samples. Because highly expressed transcripts often display high level of over-dispersion, we selected the 5,000 most highly expressed transcripts according to Helicos measures in each sample and removed those that show extreme low read coverage in Illumina (<0.3 RPKM noise level as in Sam et al., 2011). The numbers of corresponding transcripts measured by Illumina sequencing were listed in Table 1.

We applied RPKM (Mortazavi, et al., 2008), mseq (Li, et al., 2010), GPseq (Srivastava and Chen, 2010) and POME as before. For mseq, we used 40 base pairs in the neighborhood of each nucleotide, and used the top 500 highly expressed transcripts as the training dataset. To avoid complication of missing data due to unmappable regions or dubious annotation, we removed all positions with zero coverage from each transcript in the data pre-processing step.

We used Helicos measure as the gold standard and compared the Spearman's rank correlation coefficients between the Helicos measure and the estimates of transcript-level gene expression generated by the four tested methods, respectively.

Table 1 shows the overall performances of the four tested methods. POME achieved the highest Spearman correlation coefficients in ten out of the twelve samples except for sample "aT34N" and sample "VCaP", where GPseq and RPKM were the best, respectively.

To further investigate the differences between POME and the other three competing methods, we focused on a subset of transcripts with high over-dispersion and high spatial dependence. The magnitudes of over-dispersion and spatial dependence were measured by variation-to-mean ratio (also called "Fano factor") and lag one autocorrelation between base-level reads count, respectively. In each sample around 1,500 transcripts were selected with both above-median over-dispersion and above-median spatial dependence. The numbers of selected transcripts in each sample were listed in Table 2.

We again compared the Spearman's rank correlation coefficients between the Helicos measure and the estimates of transcript-level gene expression generated by the four tested methods, respectively. Table 2 shows the overall performances of the four tested methods in the selected subset of transcripts with high over-dispersion and high spatial dependence. The POME method achieved the highest Spearman correlation coefficients in all twelve samples. Based on the real data results above, we believe that POME provides more

accurate quantification of transcript-level expression than the other three competing methods, especially for transcripts with high over-dispersion level and high spatial dependence.

Table 1. Comparison of Spearman correlation coefficients between four different expression measures (RPKM, mseq, GPseq and POME) in twelve real datasets. In each dataset, the method with the highest Spearman correlation coefficient is highlighted in bold.

Sample	# ¹	RPKM	mseq	GPseq	POME
LnCaP_0	4,964	0.6246	0.6155	0.5265	0.6887
LnCaP_24	4,956	0.6186	0.6035	0.5466	0.6728
LnCaP_48	4,946	0.6001	0.5955	0.4964	0.6407
VCaP_0	4,939	0.5801	0.5607	0.5822	0.6046
VCaP_24	4,948	0.5988	0.5474	0.5936	0.6234
VCaP_48	4,941	0.6222	0.4577	0.6129	0.6569
aT34	4,869	0.5789	0.5668	0.4351	0.5958
aT34N	4,747	0.4281	0.4245	0.4624	0.3944
DU145F	4,947	0.5945	0.5793	0.4575	0.6350
DU145F2	4,943	0.5939	0.5783	0.4608	0.6263
VCaP	4,944	0.5204	0.4024	0.4398	0.5175
RWPE	4,969	0.5600	0.5514	0.4762	0.6013

¹The number of transcripts measured in Illumina sequencing.

Table 2. Comparison of Spearman correlation coefficients between four different expression measures (RPKM, mseq, GPseq and POME) on transcripts with high over-dispersion and high spatial dependence in twelve real datasets. In each dataset, the method with the highest Spearman correlation coefficient is highlighted in bold.

Sample	# ¹	RPKM	mseq	GPseq	POME
LnCaP_0	1,495	0.6454	0.6232	0.6071	0.7362
LnCaP_24	1,534	0.6248	0.5990	0.5828	0.7118
LnCaP_48	1,518	0.6059	0.5963	0.5152	0.6795
VCaP_0	1,525	0.6641	0.6452	0.6496	0.7106
VCaP_24	1,523	0.6259	0.5783	0.6691	0.6841
VCaP_48	1,500	0.6354	0.5094	0.6820	0.7022
aT34	1,563	0.6705	0.6470	0.5214	0.6914
aT34N	1,476	0.6322	0.6268	0.6203	0.6523
DU145F	1,525	0.6254	0.6017	0.5326	0.6749
DU145F2	1,529	0.6239	0.5984	0.5375	0.6809
VCaP	1,689	0.6211	0.5534	0.4853	0.6301
RWPE	1,548	0.6250	0.6079	0.5173	0.6730

¹The number of transcripts measured in Illumina sequencing with high over-dispersion and high spatial dependence.

We next repeated the above analyses using the Pearson correlation coefficients and obtained similar results. The POME method achieved the highest Pearson correlation coefficients in seven out of twelve samples using highly expressed transcripts (Table S2). Using a subset of transcripts with high over-dispersion and high spatial dependence, the POME method achieved the highest Pear-

son correlation coefficients in nine out of twelve samples (Table S3).

To understand the differences between POME and the other three competing methods (RPKM, mseq and GPseq), we zoomed in on those transcripts for which the tested methods gave dramatically different expression measures. In the discussion below, sample LnCaP_0 was used as an illustrative example. To be specific, we first transferred the estimated transcript-level expression values and the Helicos measures (gold standard) into ranks, since we used Spearman rank correlation coefficient to measure the performance of a tested method. Next we calculated the rank differences between POME and other three competing methods, and selected the top ten transcripts with the largest rank differences. We used the ranks provided by the Helicos measure as the gold standard and assumed that the better method provides a closer rank to the Helicos rank. The selected transcripts were listed in Table S4 in which we also reported the variance-to-mean ratio and the lag one auto-correlation for each transcript.

First we looked at the ten transcripts with the largest rank differences between POME and RPKM. We found that POME outperformed RPKM in all ten transcripts. The RPKM method overestimated the expression levels of these ten highly over-dispersed transcripts. In contrast, the two random effect terms in the POME model were able to account for extra variability, which contributed to POME's improved accuracy.

For the ten transcripts with the largest rank differences between POME and GPseq, POME outperformed GPseq in eight of the ten transcripts. All of these ten transcripts are extremely highly over-dispersed (variance-to-mean ratios are larger than 10). GPseq was not able to accommodate such huge variability by the two parameters of the assumed generalized Poisson model and in most cases underestimated the expression level.

For the ten transcripts with the largest rank differences between POME and mseq, POME outperformed mseq in eight of the ten transcripts. POME produced more accurate estimates than mseq when the data showed high spatial dependence, since POME explicitly incorporated the spatial dependence while mseq assumed spatial independence between reads counts covering different base pairs.

4 DISCUSSION

In microarray data analysis, it is now widely accepted that estimates of the expression levels based on parametric models such as the model-based expression index (MBEI) (Li and Wong, 2001) are more accurate in reflecting the underlying expression levels than summary statistics of raw intensity values. Inspired by the success of model-based methods in microarray data analysis, in this study, we strived to develop a model-based method for analyzing RNA-Seq data.

When exploring the properties of base-level sequencing depth in RNA-Seq data, we found substantial variation in sequencing depth within most transcripts, especially for those that are highly expressed. We also found that a large proportion of transcripts (around 25%) shows over-dispersion in base-level sequencing depth (Figure S1), which is consistent with what has been reported in the literature (Li, et al., 2010). Additionally, we found that base-level sequencing depth displays substantial dependence between base pairs that are close to each other. This type of dependence resembles spatial correlation between neighboring areas considered in spatial statistics research (Waller and Gotway, 2004), and to the

best of our knowledge, has not yet been reported in the literature on RNA-Seq data analysis. Based on these observations, we believe that spatial models that can incorporate correlations such as POME have advantages over existing models in characterizing base-level RNA-Seq data.

In this study, we used POME to estimate the expression level of an individual transcript. Two types of random effects are introduced to characterize two types of variation in read counts, which are between adjacent base pairs (i.e. spatial correlation) and specific to each base pair (i.e. non-spatial variation), respectively. Analyses of both simulated and real RNA-Seq data demonstrate that the expression indices estimated by POME reflect the underlying expression levels more accurately compared to existing methods. We believe that the improvement of estimation accuracy of POME comes from modeling the dependence between the read counts of base pairs adjacent to each other. The POME model can also be used for differential gene expression analysis by incorporating biological or experimental conditions or other covariates. We will pursue research in these directions in the future.

POME model assumes over-dispersion in the observed count data which is the case for a large proportion of transcripts in RNA-Seq data, especially those highly expressed. For this reason, we focused on highly expressed transcripts in this study. From our experience, about one quarter of all transcripts belong to this category. For these transcripts, our analysis on simulated as well as real data demonstrated POME offers more accurate expression measure than other methods we compared. We have also performed small scale tests on transcripts with medium level expression, and found POME again outperforms RPKM, mseq and GPseq methods (data not shown). For lowly expressed transcripts we do not recommend to use POME for inference because observed data may be under-dispersed.

Like many other model-based methods, the POME method is more computational intensive than read enumeration methods. For a 100 bp transcript without replicate, it takes POME 23 seconds to complete 10,000 MCMC iterations on a Dell PowerEdge 1950 computing node (2.83 GHz CPU processors and 8 GB RAM). The computation time increases almost linearly with the length of transcripts. However, because model fitting of individual transcript is independent of each other and thus can be performed simultaneously, the computational intensity of the POME method can be much mitigated by parallel computing. The power of modern cluster computers will also help in this regard. After all, we believe that accuracy in statistical inference outweighs computation cost as long as the latter is affordable.

Although we only applied the POME method to RNA-Seq data obtained from the Illumina platform in this study, we believe it can also be applied to RNA-Seq data collected from other sequencing platforms such as SOLiD with little or no modification, because these data possess similar data structures.

ACKNOWLEDGEMENTS

We thank the associate editor and three anonymous reviewers for their constructive and helpful suggestions, which substantially improved our manuscript. We thank members of the Arul Chinnaiyan lab, especially Mr. Lee Sam for sharing the prostate cancer RNA-Seq data and helpful discussion. We thank Dr. Hao Wu for

helpful discussion. We thank Mr. Han Wu for his help during the revision of this manuscript.

Funding: NSF grants DMS 1000617 and DMS 1000443. NIH grant 5R01GM080625.

REFERENCES

- Besag, J. (1974) Spatial interaction and the statistical analysis of lattice systems (with Discussion), *Journal of the Royal Statistical Society, Ser. B*, **36**, 192-236.
- Besag, J. *et al.* (1991) Bayesian image restoration, with two applications in spatial statistics, *Annals of the Institute of Statistical Mathematics*, **43**, 1-20.
- Best, N. *et al.* (2005) A comparison of Bayesian spatial models for disease mapping, *Statistical Methods in Medical Research*, **14**, 35-59.
- Bullard, J.H. *et al.* (2010) Evaluation of statistical methods for normalization and differential expression in mRNA-Seq experiments, *BMC Bioinformatics*, **11**, 94.
- Clayton, D. and Kaldor, J. (1987) Empirical Bayes estimates of age-standardized relative risks for use in disease mapping, *Biometrics*, **43**, 671-681.
- Cloonan, N. *et al.* (2008) Stem cell transcriptome profiling via massive-scale mRNA sequencing, *Nat Methods*, **5**, 613-619.
- Cressie, N. and Chan, N. (1989) Spatial Modeling of Regional Variables, *Journal of the American Statistical Association*, **84**, 393-401.
- Diggle, P. *et al.* (1998) Model-based geostatistics, *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, **47**, 299-350.
- Gilks, W. and Wild, P. (1992) Adaptive rejection sampling for Gibbs sampling, *Applied Statistics*, **41**, 337-348.
- Gilks, W.R. *et al.* (1995) Adaptive Rejection Metropolis Sampling Within Gibbs Sampling, *Applied Statistics-Journal of the Royal Statistical Society Series C*, **44**, 455-472.
- Gilks, W.R. *et al.* (1998) *Markov chain Monte Carlo in practice*. Chapman & Hall, Boca Raton, Fla.
- Li, C. and Wong, W.H. (2001) Model-based analysis of oligonucleotide arrays: expression index computation and outlier detection, *Proc Natl Acad Sci U S A*, **98**, 31-36.
- Li, J. *et al.* (2010) Modeling non-uniformity in short-read rates in RNA-Seq data, *Genome biology*, **11**, R50.
- Lister, R. *et al.* (2008) Highly integrated single-base resolution maps of the epigenome in Arabidopsis, *Cell*, **133**, 523-536.
- Liu, J.S. (2001) *Monte Carlo Strategies in Scientific Computing*. Springer Verlag, Berlin.
- Lockhart, D.J. *et al.* (1996) Expression monitoring by hybridization to high-density oligonucleotide arrays, *Nature biotechnology*, **14**, 1675-1680.
- Maher, C.A. *et al.* (2009) Transcriptome sequencing to detect gene fusions in cancer, *Nature*, **458**, 97-101.
- Marioni, J.C. *et al.* (2008) RNA-seq: an assessment of technical reproducibility and comparison with gene expression arrays, *Genome Res*, **18**, 1509-1517.

-
- Morin, R.D. *et al.* (2008) Application of massively parallel sequencing to microRNA profiling and discovery in human embryonic stem cells, *Genome Res*, **18**, 610-621.
- Mortazavi, A. *et al.* (2008) Mapping and quantifying mammalian transcriptomes by RNA-Seq, *Nat Methods*, **5**, 621-628.
- Nagalakshmi, U. *et al.* (2008) The transcriptional landscape of the yeast genome defined by RNA sequencing, *Science*, **320**, 1344-1349.
- Prensner, J.R. *et al.* (2011) Transcriptome sequencing across a prostate cancer cohort identifies PCAT-1, an unannotated lincRNA implicated in disease progression, *Nature biotechnology*, **29**, 742-749.
- Sam, L.T. *et al.* (2011) A Comparison of Single Molecule and Amplification Based Sequencing of Cancer Transcriptomes, *PLoS ONE*, **6(3): e17305**.
- Schena, M. *et al.* (1995) Quantitative monitoring of gene expression patterns with a complementary DNA microarray, *Science*, **270**, 467-470.
- Srivastava, S. and Chen, L. (2010) A two-parameter generalized Poisson model to improve the analysis of RNA-seq data, *Nucleic Acids Res*, **38**, e170.
- Trapnell, C. *et al.* (2010) Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation, *Nature biotechnology*, **28**, 511-515.
- Wakefield, J. (2007) Disease mapping and spatial regression with count data, *Biostatistics*, **8**, 158-183.
- Waller, L.A. and Gotway, C.A. (2004) *Applied spatial statistics for public health data*. John Wiley & Sons, Hoboken, N.J.
- Wang, Z. *et al.* (2009) RNA-Seq: a revolutionary tool for transcriptomics, *Nature reviews genetics*, **10**, 57-63.
- Wilhelm, B.T. *et al.* (2008) Dynamic repertoire of a eukaryotic transcriptome surveyed at single-nucleotide resolution, *Nature*, **453**, 1239-1243.
- Zhu, H. *et al.* (2009) Stochastic approximation algorithms for estimation of spatial mixed models, *Manuscript*.