

Magnetic flatness and E. Hopf's theorem for magnetic systems

Valerio Assenza, James Marshall Reber, and Ivo Terek

ABSTRACT. Using the notion of magnetic curvature recently introduced by the first author, we extend E. Hopf's theorem to the setting of magnetic systems. Namely, we prove that if the magnetic flow on the s -sphere bundle is without conjugate points, then the total magnetic curvature is non-positive, and vanishes if and only if the magnetic system is magnetically flat. We then prove that magnetic flatness is a rigid condition, in the sense that it only occurs when either the magnetic form is trivial and the metric is flat, or when the magnetic system is Kähler, the metric has constant negative sectional holomorphic curvature, and s equals the Mañé critical value.

1. Introduction

A classical rigidity theorem in Riemannian geometry is due to E. Hopf [26]: *if a closed Riemannian surface is without conjugate points, then its total Gaussian curvature is non-positive, and vanishes if and only if the metric is flat.* The Gauss-Bonnet theorem then implies that every metric without conjugate points on the two-dimensional torus \mathbb{T}^2 is flat. Hopf's theorem was generalized to higher dimensions by Green [24], with the scalar curvature replacing the Gaussian curvature. However, in absence of a higher dimensional argument à la Gauss-Bonnet, the question of whether a metric without conjugate points on a torus \mathbb{T}^n is flat remained open for decades. The problem was finally solved using a different array of techniques by Burago and Ivanov [14].

In this paper, we generalize Green's argument to the setting of magnetic systems without conjugate points. This generalization naturally leads us to studying flatness in a magnetic sense. A *magnetic system* on a closed smooth manifold M is a pair (g, σ) , where $g = \langle \cdot, \cdot \rangle$ is a Riemannian metric and σ is a closed 2-form on M . In this context, σ is referred to as the *magnetic form*, and the requirement that σ

2010 *Mathematics Subject Classification.* 53C15 (Primary), 37D40 (Secondary).

Key words and phrases. Magnetic systems · Hopf's theorem · Green bundles.

is closed corresponds to the well-known fact that magnetic fields are divergence-free, courtesy of Maxwell's equation. The skew-adjoint endomorphism Ω of TM corresponding to σ under g is referred to as the *Lorentz force*, and it is defined by

$$(1.1) \quad \sigma(v, w) = \langle v, \Omega(w) \rangle \text{ for all } v, w \in TM.$$

Recall that a magnetic system is the mathematical formalism used to describe the motion of a charged particle moving on a Riemannian manifold under the action of a magnetic force field. More precisely, trajectory $\gamma: \mathbb{R} \rightarrow M$ of a charged particle is described by the second-order ordinary differential equation

$$(1.2) \quad \frac{D\dot{\gamma}}{dt} = \Omega(\dot{\gamma}),$$

where D/dt denotes the covariant derivative along γ induced by the Levi-Civita connection ∇ of g . Solutions of (1.2) are called (g, σ) -geodesics. Lifting (1.2) to TM , we obtain the *magnetic flow* $\Phi^{(g, \sigma)}: \mathbb{R} \times TM \rightarrow TM$ associated with (g, σ) . Observe that in absence of a magnetic interaction (that is, when $\sigma = 0$), (1.2) reduces to the standard equation for geodesics, and $\Phi^{(g, 0)}$ is the geodesic flow of the metric g . Since (g, σ) -geodesics have constant speed, the infinitesimal generator of $\Phi^{(g, \sigma)}$ is tangent to the s -sphere bundles $\Sigma_s = \{v \in TM \mid \|v\| = s\}$ for $s > 0$. Thus, Σ_s is an invariant set for the flow $\Phi^{(g, \sigma)}$, and hereafter we denote by $\Phi_s^{(g, \sigma)}: \mathbb{R} \times \Sigma_s \rightarrow \Sigma_s$ the restriction of $\Phi^{(g, \sigma)}$ to Σ_s . The dynamics of magnetic systems have been studied for many decades; for some historical background, see [6, 8, 21].

Linearizing (1.2) along a (g, σ) -geodesic γ , we obtain the differential equation

$$(1.3) \quad \frac{D^2 J}{dt^2} + R(J, \dot{\gamma})\dot{\gamma} = (\nabla_J \Omega)(\dot{\gamma}) + \Omega\left(\frac{DJ}{dt}\right)$$

for vector fields J along γ . A solution J of (1.3) is called a (g, σ) -Jacobi field. In addition, a (g, σ) -Jacobi field J is *normal* if it satisfies

$$(1.4) \quad \left\langle \frac{DJ}{dt}, \dot{\gamma} \right\rangle = 0.$$

A point $\gamma(\tau)$ is said to be *conjugate* to $\gamma(0)$ along γ if there exists a nontrivial normal Jacobi field J along γ such that $J(0) = 0$ and $J(\tau)$ is proportional to $\dot{\gamma}(\tau)$. A (g, σ) -geodesic γ is *without conjugate points* if $\gamma(\tau)$ is not conjugate to $\gamma(0)$ for every $\tau \neq 0$, and the flow $\Phi_s^{(g, \sigma)}$ is *without conjugate points* if all of (g, σ) -geodesics with speed s are without conjugate points.¹

In [10], the first author introduced a notion of magnetic curvature related to the second variation of the action which extends to higher dimensions the well-known Gaussian magnetic curvature introduced by G. and M. Paternain in [31]

¹The above definition is related to critical values of a suitable exponential map defined for magnetic systems. For more details, see [5, 25].

for surfaces. We briefly recall its definition. Let E and E^1 be the bundles of complementary and unit-complementary directions over the unit sphere bundle SM , whose fibers over $v \in SM$ are given by $E_v = v^\perp$ and $E_v^1 = v^\perp \cap SM$. Denote by R the Riemann curvature tensor, and for $s > 0$, consider the bundle endomorphisms $R_s^\Omega, A^\Omega : E \rightarrow E$ given by

$$(1.5) \quad (R_s^\Omega)_v(w) = s^2 R(w, v)v - s(\nabla_w \Omega)(v) + \frac{s}{2} ((\nabla_v \Omega)(w) - \langle (\nabla_v \Omega)(w), v \rangle v),$$

$$(1.6) \quad (A^\Omega)_v(w) = \frac{3}{4} \langle w, \Omega(v) \rangle \Omega(v) - \frac{1}{4} \Omega^2(w) - \frac{1}{4} \langle \Omega(w), \Omega(v) \rangle v.$$

The *magnetic curvature operator at level s* is the section M_s^Ω of $\text{End}(E)$ given by

$$(1.7) \quad (M_s^\Omega)_v(w) = (R_s^\Omega)_v(w) + (A^\Omega)_v(w).$$

The *magnetic sectional curvature* $\text{sec}_s^\Omega : E^1 \rightarrow \mathbb{R}$ and the *magnetic Ricci curvature* $\text{Ric}_s^\Omega : SM \rightarrow \mathbb{R}$ at level s are given by, respectively:

$$(1.8) \quad (\text{sec}_s^\Omega)_v(w) = \langle (M_s^\Omega)_v(w), w \rangle \quad \text{and} \quad \text{Ric}_s^\Omega(v) = \text{tr}(M_s^\Omega)_v.$$

We note that the magnetic curvature function captures relevant dynamical information for the magnetic system. For instance, it can be deduced from [34, Theorem 4.1] that if $\text{sec}_s^\Omega < 0$, then $\Phi_s^{(\mathbf{g}, \sigma)}$ is Anosov. Under the assumption that $\text{Ric}_s^\Omega > 0$, it was shown in [10] that there exists a closed contractible (\mathbf{g}, σ) -geodesic for small s .²

In Proposition 3.1, we relate M_s^Ω to (1.3), and in Proposition 3.2 we show that if $\text{sec}_s^\Omega \leq 0$, then $\Phi_s^{(\mathbf{g}, \sigma)}$ is without conjugate points. As in the Riemannian setting, the converse does not hold in general. An example of a magnetic flow without conjugate points whose magnetic curvature changes sign was constructed [15, Section 7]. Nevertheless, our main result gives a partial converse, generalizing [23, Theorem 1] from the case where $\nabla \sigma = 0$:

THEOREM A. *Let (\mathbf{g}, σ) be a magnetic system on a closed manifold M , and let $s > 0$. If $\Phi_s^{(\mathbf{g}, \sigma)}$ is without conjugate points, then*

$$(1.9) \quad \int_{SM} \text{Ric}_s^\Omega(v) \, d\mu_{\mathbf{g}}(v) \leq 0,$$

where $d\mu_{\mathbf{g}}$ denotes the Liouville measure on SM . Moreover, equality holds if and only if $M_s^\Omega = 0$.

The strategy of the proof is similar to the one given in [24]. With the aid of an appropriate twisted connection introduced in Section 2, we construct the *magnetic Green bundles* along (\mathbf{g}, σ) -geodesics without conjugate points in Section 4. The

²See [11] for previous results in this direction.

existence of such bundles guarantees that we are able to find non-trivial solutions to a suitable *magnetic Riccati equation* (see (5.3)), and the result follows quickly. We note that similar geometric constructions were previously introduced in the classical Riemannian case by Green in [24], for Finsler metrics without conjugate points in [20], and in the general context of Hamiltonian dynamics in [7, 13, 18].

We now observe some consequences of Theorem A. Using the linearity of Ω , one can deduce

$$(1.10) \quad \int_{SM} \text{tr}(R_s^\Omega)_v \, d\mu_{\mathbf{g}}(v) = s^2 \frac{\text{vol}(\mathbb{S}^{n-1})}{n} \int_M \text{scal}(x) \, d\nu_{\mathbf{g}}(x),$$

where $d\nu_{\mathbf{g}}$ is the Riemannian volume measure on M . Combining (1.10) with the fact that $\text{tr}(A^\Omega)_v$ is always non-negative and is positive on some open subset of SM as soon as σ is not identically zero (cf. [10, Lemma 5] or (1.6)), the following corollary holds:

COROLLARY B. *If σ is not identically zero, then for every sufficiently small s the flow $\Phi_s^{(\mathbf{g}, \sigma)}$ has at least one pair conjugate points. Moreover, if \mathbf{g} is a Riemannian metric on M such that*

$$\int_M \text{scal}(x) \, d\nu_{\mathbf{g}}(x) > 0,$$

then for every magnetic form σ and every $s > 0$ the flow $\Phi_s^{(\mathbf{g}, \sigma)}$ has at least one pair conjugate points.

This can be interpreted as saying that if the magnetic deformation is strong, then the magnetic flow must have at least one pair conjugate points. Naturally, this leads us to the following question: *given a Riemannian metric \mathbf{g} with at least one pair conjugate points, does there exist a closed 2-form σ and a value s such that the corresponding flow $\Phi_s^{(\mathbf{g}, \sigma)}$ is without conjugate points?* While Corollary B points to a negative answer for small values of s , the question is open to our best knowledge. Some progress towards this question has been made by Bialy in [12], where it is shown that if \mathbf{g} is conformally flat on the n -torus \mathbb{T}^n , then for every s the magnetic flow $\Phi_s^{(\mathbf{g}, \sigma)}$ is without conjugate points if and only if σ is identically zero and the metric is flat. Bialy also conjectured that such a result should hold for arbitrary metrics, which would generalize [14] to the magnetic case.

If M is an orientable surface, then every closed 2-form σ is related to the area form $\nu_{\mathbf{g}}$ by a smooth function b , in the sense that $\sigma = b\nu_{\mathbf{g}}$. We denote the magnetic system $(\mathbf{g}, b\nu_{\mathbf{g}})$ by (\mathbf{g}, b) . In this setting, the magnetic curvature functions sec_s^Ω and Ric_s^Ω reduce to the *Gaussian magnetic curvature at level s* (cf. [10, Lemma 8]), which is given by

$$K_s^{\mathbf{g}, b}(x, v) = s^2 K^{\mathbf{g}}(x) - s \, db_x(iv) + b(x)^2.$$

Here, iv is the vector obtained by rotating v by $\pi/2$ according to the fixed orientation, and $K^g: M \rightarrow \mathbb{R}$ is the Gaussian curvature. As a consequence of the Gauss-Bonnet theorem, we have

$$\int_{SM} K_s^{g,b}(x, v) d\mu_g(x, v) = 2\pi \left[s^2 \chi(M) + \int_M b^2(x) dv_g(x) \right],$$

where $\chi(M)$ is the Euler characteristic of M . As pointed out in Corollary B, every magnetic flow on S^2 has conjugate points. In the case of a magnetic flow on \mathbb{T}^2 , we are able to recover [12, Theorem 1]:

COROLLARY C. *The only magnetic flows on \mathbb{T}^2 without conjugate points are of the form $\Phi_s^{(g_0, 0)}$, where g_0 is a flat metric.*

Summarizing, the genus of the surface must be at least two for there to be a non-trivial magnetic system without conjugate points. If we assume that $(g, b) \neq (g_0, 0)$, then $K_s^{g,b} \equiv 0$ if and only if b is constant, g has constant negative curvature k , and $s = |b|/\sqrt{-k}$. Observe that if this happens, then $\Phi_s^{(g,b)}$ is the horocycle flow [16, Section 5.2]. In particular, this allows us to view the horocycle flow on hyperbolic surfaces as the magnetic analogue of \mathbb{T}^2 equipped with a flat metric.

In order to bring the discussion to higher dimensions, we recall that the *Mañé critical value* $s_0 = s_0(g, \sigma) \in [0, +\infty]$ of the pair (g, σ) is defined as

$$(1.11) \quad s_0 = \begin{cases} \inf_{d\theta = p^*\sigma} \sup_{x \in M} \|\theta_x\| & \text{if } p^*\sigma \text{ is exact,} \\ +\infty & \text{otherwise,} \end{cases}$$

where $p: \tilde{M} \rightarrow M$ is the universal covering map and $\|\cdot\|$ denotes the dual norm induced by g . One can immediately see that s_0 is finite if and only if $p^*\sigma$ admits a bounded primitive, and $s_0 = 0$ if and only if $\sigma = 0$. The Mañé critical value marks a drastic change of the dynamics of $\Phi_s^{(g, \sigma)}$, and its computation is challenging in general. For more details and alternative definitions of s_0 , we point the reader to [16, 29], and references within. In the special case when (M, g) is a Kähler manifold with Kähler form ω and constant holomorphic sectional curvature $\text{sec}^{\text{hol}} \equiv k < 0$, we have

$$(1.12) \quad s_0(g, \lambda\omega) = \frac{|\lambda|}{\sqrt{-k}};$$

see Appendix A.

We say that a magnetic system (g, σ) is *magnetically flat at level* $s > 0$ if $M_s^\Omega \equiv 0$. Just like in the surface case, it turns out that being magnetically flat is a rigid condition, and there are only two types of magnetic systems (g, σ) and levels s for which it is satisfied: when $\sigma = 0$ and g is flat (in which case $s > 0$ can be

anything), or when $(g, \lambda\omega)$ is a Kähler magnetic system on a Kähler manifold with constant negative holomorphic sectional curvature and $s = s_0(g, \lambda\omega)$. This is a consequence of the following theorem, which we prove in Section 6:

THEOREM D. *Let (g, σ) be a magnetic system on a (not necessarily closed) manifold M with $\sigma \neq 0$. If $\sec_s^\Omega \equiv c$ for some $s > 0$, then $\nabla\sigma = 0$, σ is nowhere vanishing, and one of the following conclusions must hold:*

- (i) (M, g) is an oriented surface whose Gaussian curvature is constant and equal to $(c - \|\Omega\|^2)/s^2$, and $\sigma = -\|\Omega\|v_g$;
- (ii) $\dim M \geq 4$, $c = 0$, and $\|\Omega\|^{-1}\Omega$ makes (M, g) a Kähler manifold with constant holomorphic sectional curvature equal to $-\|\Omega\|^2/s^2$.

Moreover, assuming that $c = 0$ in (i), we have that $s = s_0(g, \sigma)$ in either case.

We conclude by connecting the above discussion with the problem of existence of closed (g, σ) -geodesics with speed s . The literature around this problem is vast; for more details, see [9, 17, 29]. In general, existence depends on whether s is above or below s_0 . Indeed, for every $s > s_0$, there exists a closed (g, σ) -geodesic with speed s [19, Theorem 27]. Moreover, if M is not simply connected and $s > s_0$, then for every non-trivial free homotopy class contains a (g, σ) -geodesic with speed s [9]. On the other hand, for almost every sufficiently small $s < s_0$, there exists a contractible closed (g, σ) -geodesic with speed s [32, Corollary 3.5]. It is conjectured that the latter conclusion in fact holds for all $s \in (0, s_0)$. To our best knowledge, the only examples of magnetic flows without closed (g, σ) -geodesics are the ones of the form $\Phi_{s_0}^{(g, \lambda\omega)}$, where $(g, \lambda\omega)$ is a Kähler magnetic system with constant negative holomorphic sectional curvature and s_0 is the Mañé critical value. This raises the following question: *if there are no closed (g, σ) -geodesics with speed s , is the system (g, σ) magnetically flat at level s ?*

Acknowledgements and funding

We would like to thank Jacopo de Simoi and Andrey Gogolev for their support throughout this project. We would also like to thank Johanna Bimmerman for her notes on closed orbits for Kähler manifolds. Finally, we would like to thank Gabriele Benedetti, Andrey Gogolev, Leonardo Macarini, Gabriel Paternain, and Jacopo de Simoi for feedback on early drafts of the paper.

All authors were supported by NSERC Discovery Grant RGPIN-2022-04188. The first author was supported by Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project-ID 281071066 — TRR 191 and by the Instituto Serapilheira. The second author was supported by NSF DMS-2247747. The third author was supported by FAPESP–OSU Regular Research Award 2015/50265-6.

2. The anisotropic objects $\tilde{\Omega}$ and $\tilde{\mathcal{D}}$

In this section, we fix a magnetic system (g, σ) on a (not necessarily closed) manifold M , and denote by $\pi: TM \rightarrow M$ the natural projection. We construct the anisotropic Lorentz force and the anisotropic twisted connection, which will be useful tools for the calculations done in Sections 3, 4, and 5.

2.1. The anisotropic Lorentz force. The *anisotropic Lorentz force* $\tilde{\Omega}$ is defined by assigning to every $v \in SM$ the skew-adjoint operator $\tilde{\Omega}_v: T_{\pi(v)}M \rightarrow T_{\pi(v)}M$ given by

$$(2.1) \quad \tilde{\Omega}_v(w) = \Omega(w)^\top + \Omega(w^\top) + \frac{1}{2}\Omega(w^\perp)^\perp,$$

where $[\cdot]^\top$ and $[\cdot]^\perp$ denote the projections onto $\mathbb{R}v$ and v^\perp , according to the direct sum decomposition $T_{\pi(v)}M = \mathbb{R}v \oplus v^\perp$. A straightforward computation yields the alternative formula

$$(2.2) \quad \tilde{\Omega}_v(w) = \frac{1}{2} (\langle \Omega(w), v \rangle v + \langle w, v \rangle \Omega(v) + \Omega(w))$$

for $\tilde{\Omega}$. In particular, note that $\tilde{\Omega}_v$ is completely characterized by the relations

$$(2.3) \quad \text{i) } \tilde{\Omega}_v(v) = \Omega(v) \quad \text{ii) } \langle \tilde{\Omega}_v(w), v \rangle = \langle \Omega(w), v \rangle \quad \text{iii) } \tilde{\Omega}_v(w^\perp)^\perp = \frac{1}{2}\Omega(w^\perp)^\perp$$

imposed on all $w \in T_{\pi(v)}M$. Repeatedly applying (2.2) and recalling (1.6) yields

$$(2.4) \quad \tilde{\Omega}_v^2(w) = -(A^\Omega)_v(w^\perp) + \frac{1}{2}(\Omega^2(w^\top) + \Omega^2(w)^\top).$$

2.2. The anisotropic twisted connection. Let $\gamma: I \rightarrow M$ be a (g, σ) -geodesic with constant speed $s > 0$. We define a covariant derivative operator $\tilde{\mathcal{D}}$ acting on vector fields V along γ by

$$(2.5) \quad \tilde{\mathcal{D}}V = \frac{DV}{dt} - \tilde{\Omega}_{\dot{\gamma}/s}(V),$$

where D/dt is the Levi-Civita covariant derivative along γ , and $\tilde{\Omega}$ is as in (2.1). Note that as $\tilde{\Omega}_{\dot{\gamma}/s}$ is skew-adjoint, $\tilde{\mathcal{D}}$ is metric-compatible [10, Lemma 10], and so it commutes with tangent and orthogonal projections along γ . Furthermore, in view of (2.3-i) and (1.2), we have that $\tilde{\mathcal{D}}\dot{\gamma} = 0$; in fact, this is equivalent to (1.2).

Warning: For economy of notation, we will write simply $\tilde{\Omega}$ instead of $\tilde{\Omega}_{\dot{\gamma}/s}$ when using (2.5) in the next sections.

3. Curvature identities

The main goal of this section is to establish the following result:

PROPOSITION 3.1. *Let (\mathbf{g}, σ) be a magnetic system on a (not necessarily closed) manifold M , $\gamma: I \rightarrow M$ be a (\mathbf{g}, σ) -geodesic with speed $s > 0$, and J be a normal (\mathbf{g}, σ) -Jacobi field along γ . Then*

$$(3.1) \quad \tilde{\mathcal{D}}^2 J^\perp + (M_s^\Omega)_{\dot{\gamma}/s}(J^\perp) = 0$$

on I .

Here, we will need to consider the projections P^\top and P^\perp , acting on vector fields V along γ via

$$P^\top(V) = V^\top = s^{-2}\langle V, \dot{\gamma} \rangle \dot{\gamma} \quad \text{and} \quad P^\perp(V) = V^\perp = V - s^{-2}\langle V, \dot{\gamma} \rangle \dot{\gamma}.$$

Denoting Levi-Civita covariant derivatives with dots, a straightforward computation shows that $(P^\top)^\cdot = [\Omega, P^\top]$ and $(P^\perp)^\cdot = [\Omega, P^\perp]$. With these commutator identities in place, it follows from (2.1) and from differentiating $\tilde{\Omega}(\dot{\gamma}) = \Omega(\dot{\gamma})$ that

$$(3.2) \quad \text{i) } P^\perp \circ \dot{\tilde{\Omega}} \circ P^\perp = \frac{1}{2}P^\perp \circ \dot{\Omega} \circ P^\perp \quad \text{and} \quad \text{ii) } \dot{\tilde{\Omega}}(\dot{\gamma}) = \dot{\Omega}(\dot{\gamma}) + \frac{1}{2}\Omega^2(\dot{\gamma})^\perp,$$

respectively.

PROOF OF PROPOSITION 3.1. We start by computing

$$(3.3) \quad \tilde{\mathcal{D}}^2 J = \tilde{\mathcal{D}}(J - \tilde{\Omega}(J)) = \dot{J} - 2\tilde{\Omega}(J) - \dot{\tilde{\Omega}}(J) + \tilde{\Omega}^2(J),$$

and substituting the (\mathbf{g}, σ) -Jacobi equation (1.3) together with (2.4) to obtain

$$(3.4) \quad \begin{aligned} \tilde{\mathcal{D}}^2 J &= R(\dot{\gamma}, J)\dot{\gamma} + (\nabla_J \Omega)(\dot{\gamma}) + \Omega(J) - 2\tilde{\Omega}(J) \\ &\quad - \dot{\tilde{\Omega}}(J) - (A^\Omega)_{\dot{\gamma}/s}(J^\perp) + \frac{1}{2}(\Omega^2(J^\top) + \Omega^2(J)^\top). \end{aligned}$$

At the same time, using that (3.2-ii) remains valid when replacing $\dot{\gamma}$ with J^\top , we may evaluate (1.5) as

$$(3.5) \quad (R_s^\Omega)_{\dot{\gamma}/s}(J^\perp) = -R(\dot{\gamma}, J^\perp)\dot{\gamma} - (\nabla_{J^\perp} \Omega)(\dot{\gamma}) + \dot{\tilde{\Omega}}(J^\perp)^\perp.$$

As $R(\dot{\gamma}, J)\dot{\gamma} = R(\dot{\gamma}, J^\perp)\dot{\gamma}$ and $(\nabla_J \Omega)(\dot{\gamma}) = \dot{\Omega}(J^\top) + (\nabla_{J^\perp} \Omega)(\dot{\gamma})$, we may substitute (3.5) into (3.4) and use (1.7) to write

$$(3.6) \quad \begin{aligned} \tilde{\mathcal{D}}^2 J &= -(M_s^\Omega)_{\dot{\gamma}/s}(J^\perp) + \dot{\tilde{\Omega}}(J^\perp)^\perp + \dot{\Omega}(J^\top) + \Omega(J) \\ &\quad - 2\tilde{\Omega}(J) - \dot{\tilde{\Omega}}(J) + \frac{1}{2}(\Omega^2(J^\top) + \Omega^2(J)^\top). \end{aligned}$$

Applying P^\perp to both sides of (3.6) and noting that $\Omega(J)^\perp - 2\tilde{\Omega}(J)^\perp = 0$, by normality of J and (2.3-iii), it follows that

$$(3.7) \quad \tilde{\mathcal{D}}^2 J^\perp = -(M_s^\Omega)_{\dot{\gamma}/s}(J^\perp) + \check{\Omega}(J^\perp)^\perp + \dot{\Omega}(J^\top) - \check{\Omega}(J)^\perp + \frac{1}{2}\Omega^2(J^\top)^\perp.$$

Decomposing $\check{\Omega}(J)^\perp = \check{\Omega}(J^\top)^\perp + \check{\Omega}(J^\perp)^\perp$ and using (3.2-ii) for a second time, a complete cancellation occurs and (3.7) reduces to (3.1), as required. \square

We conclude this section with one interesting consequence of Proposition 3.1.

PROPOSITION 3.2. *Let (g, σ) be a magnetic system on a (not necessarily closed) manifold M . If $s > 0$ is such that $\sec_s^\Omega \leq 0$, then $\Phi_s^{(g, \sigma)}$ is without conjugate points.*

PROOF. Let $v \in \Sigma_s$, and assume that J is any normal (g, σ) -Jacobi field along the geodesic γ with initial velocity v . Using metric compatibility of $\tilde{\mathcal{D}}$ together with (3.1), we find that

$$(3.8) \quad \begin{aligned} \frac{d^2}{dt^2} \frac{\|J^\perp\|^2}{2} &= \|\tilde{\mathcal{D}}J^\perp\|^2 + \langle \tilde{\mathcal{D}}^2 J^\perp, J^\perp \rangle \\ &= \|\tilde{\mathcal{D}}J^\perp\|^2 - \langle (M_s^\Omega)_{\dot{\gamma}/s}(J^\perp), J^\perp \rangle \\ &= \|\tilde{\mathcal{D}}J^\perp\|^2 - (\sec_s^\Omega)_{\dot{\gamma}/s} \left(\frac{J^\perp}{\|J^\perp\|} \right) \|J^\perp\|^2 \geq 0, \end{aligned}$$

where the \sec_s^Ω term is understood to be zero at all instants on which J^\perp vanishes. Hence, the function $t \mapsto \|J(t)^\perp\|^2$ is increasing whenever J is normal. If, in addition, we suppose that $J(0) = 0$ and that $J(t_0)$ is proportional to $\dot{\gamma}(t_0)$ for some $t_0 > 0$, then it must be the case that $J^\perp|_{[0, t_0]} = 0$ identically, making $J|_{[0, t_0]}$ tangential. As a tangential and normal (g, σ) -Jacobi field is necessarily a constant multiple of $\dot{\gamma}$, $J(0) = 0$ implies that $J|_{[0, t_0]} = 0$, and so $\gamma(t_0)$ is not conjugate to $\gamma(0)$ along γ . \square

4. Magnetic Green bundles

In this section, we construct the so-called Green bundles associated with a magnetic flow $\Phi_s^{(g, \sigma)}$. As described in the Introduction, this is a useful tool for obtaining non-trivial normal (g, σ) -Jacobi fields along any given (g, σ) -geodesic. More precisely, in Section 4.2 we introduce a symplectic vector bundle Q over Σ_s , whose fibers are in correspondence with suitable spaces of normal (g, σ) -Jacobi fields (see Lemma 4.2). We conclude by constructing the Green bundles in Lemma 4.4.

4.1. General setup. Let (g, σ) be a magnetic system on a closed manifold M . On TM , we may consider the *twisted symplectic form* ω^σ defined by

$$(4.1) \quad \omega^\sigma(\xi, \eta) = \langle \pi_* \xi, K(\eta) \rangle - \langle K(\xi), \pi_* \eta \rangle - \sigma(\pi_* \xi, \pi_* \eta),$$

for all $\xi, \eta \in T(TM)$. Here, $\pi: TM \rightarrow M$ is the natural bundle projection, and $K: TTM \rightarrow TM$ is the *Levi-Civita connector* of (M, g) .

It is well-known that

$$(4.2) \quad \text{the magnetic flow } \Phi^{(g, \sigma)}: \mathbb{R} \times TM \rightarrow TM \text{ is Hamiltonian,}$$

as its generator $X^{g, \sigma} \in \mathfrak{X}(TM)$ satisfies $\omega^\sigma(X^{g, \sigma}, \cdot) = dL$, where $L: TM \rightarrow \mathbb{R}$ is given by $L(v) = \|v\|^2/2$. Indeed, a direct consequence of (4.1) together with

$$(4.3) \quad \text{(i) } dL_v(\eta) = \langle K(\eta), v \rangle \quad \text{and} \quad \text{(ii) } K(X_v^{g, \sigma}) = \Omega(v).$$

Clearly (4.3-i) holds, while (4.3-ii) follows from the easily verified general relation $X_v^{g, \sigma} = X_v^g + \Omega^\vee(v)$, where $X^g \in \mathfrak{X}(TM)$ denotes the geodesic vector field of (M, g) and $\Omega^\vee(v) \in T_v(TM)$ is the vertical lift of $\Omega(v)$, characterized by $\pi_*(\Omega^\vee(v)) = 0$ and $K(\Omega^\vee(v)) = \Omega(v)$.

In the following, instead of K we will use

$$(4.4) \quad \begin{aligned} &\text{the twisted connector } K^\sigma: TTM \setminus \{0\} \rightarrow TM, \text{ de-} \\ &\text{fined by } K^\sigma(\xi) = K(\xi) - \tilde{\Omega}(\pi_* \xi), \text{ for } \xi \in T_v(TM), \end{aligned}$$

where $\tilde{\Omega}$ is the anisotropic Lorentz force given by (2.1). Note that the zero section must be removed from TTM in (4.4) so that $\tilde{\Omega}$ may be evaluated. In addition, observe that $K^\sigma(X^{g, \sigma}) = 0$ as a consequence of (4.3-ii) and (2.3-i).

4.2. Reduced magnetic Green bundles over s -sphere bundles. Let $s > 0$ and fix $\Sigma_s = \{v \in TM \mid \|v\| = s\}$. We denote by π and K^σ the restrictions to Σ_s of the corresponding objects in TM , and by $\Phi_s^{(g, \sigma)}$ the magnetic flow restricted to Σ_s .

Combining (4.3-i) with (4.4) and (2.3-i), we see that

$$(4.5) \quad T_v \Sigma_s = \{\xi \in T_v(TM) \mid \langle K^\sigma(\xi) + \Omega(\pi_* \xi), v \rangle = 0\},$$

for every $v \in \Sigma_s$. Two distinguished subspaces of (4.5) are the (g, σ) -horizontal and vertical spaces at v , defined by $H_v^\sigma = \ker K^\sigma|_{T_v \Sigma_s}$ and $V = \ker \pi_*|_{T_v \Sigma_s}$, respectively, and they provide

$$(4.6) \quad \text{a direct sum decomposition } T\Sigma_s = H^\sigma \oplus V.$$

Indeed, it is straightforward to check that $H^\sigma \cap V = \{0\}$ by using that K and K^σ agree on vertical vectors, while the restrictions

$$(4.7) \quad \text{(i) } K^\sigma: V_v \rightarrow v^\perp \quad \text{and} \quad \text{(ii) } \pi_*: H_v^\sigma \rightarrow T_{\pi(v)}M$$

are isomorphisms, so that $\dim T_v \Sigma_s = \dim H_v^\sigma + \dim V_v$. Formula (4.1) can now be made simpler:

LEMMA 4.1. *With the above setup,*

$$(4.8) \quad \omega^\sigma(\zeta, \eta) = \langle \pi_* \zeta, K^\sigma(\eta) \rangle - \langle K^\sigma(\zeta), \pi_* \eta \rangle,$$

for all $\zeta, \eta \in T\Sigma_s$. In particular, ω^σ restricted to Σ_s is degenerate and has its kernel spanned by $X^{\mathfrak{g}, \sigma}$.

PROOF. For $\zeta, \eta \in T_v \Sigma_s$, decompose $T_{\pi(v)} M = \mathbb{R}v \oplus v^\perp$, and denote by $[\cdot]^\top$ and $[\cdot]^\perp$ the orthogonal projections onto $\mathbb{R}v$ and v^\perp , respectively. Using (2.2) and (1.1), (4.1) becomes

$$(4.9) \quad \begin{aligned} \omega^\sigma(\zeta, \eta) &= \langle \pi_* \zeta, K^\sigma(\eta) \rangle - \langle K^\sigma(\zeta), \pi_* \eta \rangle \\ &\quad + \langle \Omega(\pi_* \eta), [\pi_* \zeta]^\top \rangle - \langle \Omega(\pi_* \zeta), [\pi_* \eta]^\top \rangle. \end{aligned}$$

As $\zeta, \eta \in T_v \Sigma_s$, self-adjointness of $[\cdot]^\top$ and (4.5) together lead to the relation $\omega^\sigma(\zeta, \eta) = \langle \pi_* \zeta, K^\sigma(\eta)^\perp \rangle - \langle K^\sigma(\zeta)^\perp, \pi_* \eta \rangle$. Writing $\eta = \eta^h + \eta^v$ according to (4.6), note that $K^\sigma(\eta)^\perp = K^\sigma(\eta^v)^\perp = K^\sigma(\eta^v) = K^\sigma(\eta)$, as (4.7-i) is v^\perp -valued. The same calculation holds for ζ , and (4.8) follows.

Finally, if $\zeta \in T_v \Sigma_s$ is such that $\omega^\sigma(\zeta, T_v \Sigma_s) = 0$, then $\omega^\sigma(\zeta, H_v^\sigma) = 0$ implies that $K^\sigma(\zeta) = 0$, while $\omega^\sigma(\zeta, V_v) = 0$ means that $\pi_* \zeta = \lambda v$ for some $\lambda \in \mathbb{R}$. Both conditions together yield $\zeta = \lambda X_v^{\mathfrak{g}, \sigma}$, as required. \square

By modding out $\mathbb{R}X^{\mathfrak{g}, \sigma}$ in (4.6), we obtain a symplectic vector bundle Q over Σ_s , whose fiber over $v \in \Sigma_s$ is given by $Q_v = T_v \Sigma_s / \mathbb{R}X_v^{\mathfrak{g}, \sigma}$ and whose symplectic structure is naturally induced by ω^σ . As $X_v^{\mathfrak{g}, \sigma} \notin V_v$, we may also identify Q_v with the direct sum $[H_v^\sigma / \mathbb{R}X_v^{\mathfrak{g}, \sigma}] \oplus V_v$. In addition, due to (4.2) and $X^{\mathfrak{g}, \sigma}$ being $\Phi_s^{(\mathfrak{g}, \sigma)}$ -related to itself, we see that

$$(4.10) \quad \begin{aligned} &\text{for each } t \in \mathbb{R}, \text{ the quotient flow } \Psi^{\mathfrak{g}, \sigma}: \mathbb{R} \times Q \rightarrow Q \text{ induced by the} \\ &\text{derivative } d\Phi_s^{(\mathfrak{g}, \sigma)}: \mathbb{R} \times T\Sigma_s \rightarrow T\Sigma_s \text{ is a bundle symplectomorphism.} \end{aligned}$$

For notational simplicity, we write $\Psi_t^{(\mathfrak{g}, \sigma)} = \Psi^{\mathfrak{g}, \sigma}(t, \cdot)$. For every $v \in \Sigma_s$ and $t \in \mathbb{R}$, we define $E_v^\sigma(t) = (\Psi_t^{(\mathfrak{g}, \sigma)})^{-1}[V_{\Phi_s^{(\mathfrak{g}, \sigma)}(t, v)}]$, so that $(\Psi_t^{(\mathfrak{g}, \sigma)})_v[E_v^\sigma(t)] = V_{\Phi_s^{(\mathfrak{g}, \sigma)}(t, v)}$. Here, we identify each V_v with its isomorphic image under the quotient projection $T\Sigma_s \rightarrow Q$. It follows from (4.8) that V is a Lagrangian subbundle of Q , and thus, by (4.10),

$$(4.11) \quad \mathbb{R} \ni t \mapsto E_v^\sigma(t) \text{ is a curve of Lagrangian subspaces of } Q_v.$$

To study the behavior of $E_v^\sigma(t)$ as $t \rightarrow \pm\infty$, it will be convenient to use a different characterization of Q . For each $v \in \Sigma_s$, we let $\mathcal{J}_0^{\mathfrak{g}, \sigma, n}(v)$ be the space of all normal (\mathfrak{g}, σ) -Jacobi fields along γ_v such that $J(0)^\top = 0$.

LEMMA 4.2. *The mapping*

$$(4.12) \quad Q_v \ni \zeta + \mathbb{R}X_v^{\mathfrak{g},\sigma} \mapsto J_\zeta \in \mathcal{J}_0^{\mathfrak{g},\sigma,n}(v),$$

where J_ζ is the unique normal (\mathfrak{g},σ) -Jacobi field determined by the initial conditions $J_\zeta(0) = [\pi_*\zeta]^\perp$ and $(\tilde{\mathcal{D}}J_\zeta)(0) = K^\sigma(\zeta)$, is a well-defined linear isomorphism.

PROOF. It is easy to see that if one replaces ζ with $\zeta + \lambda X_v^{\mathfrak{g},\sigma}$ for some $\lambda \in \mathbb{R}$, then such initial conditions remain unchanged, while J_ζ is automatically normal in view of $\zeta \in T_v\Sigma_s$ combined with (4.5). Finally, (4.12) is injective for reasons explained in the proof of Lemma 4.1: namely, if $[\pi_*\zeta]^\perp = 0$, so that $\pi_*\zeta = \lambda v$ for some $\lambda \in \mathbb{R}$, then $K^\sigma(\zeta) = 0$ implies that $\zeta = \lambda X_v^{\mathfrak{g},\sigma}$, and thus $\zeta + \mathbb{R}X_v^{\mathfrak{g},\sigma}$ is zero in Q_v . As $\dim Q_v = \dim \mathcal{J}_0^{\mathfrak{g},\sigma,n}(v) = 2(n-1)$, it follows that (4.12) is an isomorphism. \square

As the next result shows, the condition of $\Phi_s^{(\mathfrak{g},\sigma)}$ being without conjugate points is what will allow us to proceed.

PROPOSITION 4.3. *Let γ be the (\mathfrak{g},σ) -geodesic with initial velocity $v \in \Sigma_s$. For $\tau \in \mathbb{R}$, $E_v^\sigma(\tau) \cap V_v \neq \{0\}$ if and only if $\gamma(\tau)$ is conjugate to $\gamma(0) = \pi(v)$ along γ .*

PROOF. Note that, under the identification (4.12), we may express the vertical space as $V_v \cong \{J \in \mathcal{J}_0^{\mathfrak{g},\sigma,n}(v) \mid J(0) = 0\}$. The quotient flow introduced in (4.10), in turn, appears as the mapping $\Psi_\tau^{\mathfrak{g},\sigma} : \mathcal{J}_0^{\mathfrak{g},\sigma,n}(v) \rightarrow \mathcal{J}_0^{\mathfrak{g},\sigma,n}(\Phi_s^{(\mathfrak{g},\sigma)}(\tau, v))$ given by

$$(4.13) \quad [\Psi_\tau^{\mathfrak{g},\sigma}(J)](t) = J(t + \tau) - s^{-2}\langle J(\tau), \dot{\gamma}(\tau) \rangle \dot{\gamma}(t + \tau).$$

Indeed, that the derivative of the magnetic flow pushes J to $t \mapsto J(t + \tau)$ is obtained exactly as in [30, Lemma 1.40], while the term subtracted in (4.13) amounts to the natural projection of a normal (\mathfrak{g},σ) -Jacobi field along γ onto $\mathcal{J}_0^{\mathfrak{g},\sigma,n}(v)$. As $[\Psi_\tau^{\mathfrak{g},\sigma}(J)](0) = J(\tau)^\perp$, we see that $E_v^\sigma(\tau) \cong \{J \in \mathcal{J}_0^{\mathfrak{g},\sigma,n}(v) \mid J(\tau)^\perp = 0\}$ under (4.12), so that

$$(4.14) \quad E_v^\sigma(\tau) \cap V_v \cong \{J \in \mathcal{J}_0^{\mathfrak{g},\sigma,n}(v) \mid J(0) = J(\tau)^\perp = 0\}.$$

\square

From here on, we assume that $\Phi_s^{(\mathfrak{g},\sigma)}$ is without conjugate points. Observe the isomorphism $H_v^\sigma / \mathbb{R}X_v^{\mathfrak{g},\sigma} \cong v^\perp$ induced by (4.7-ii), in view of $\pi_*(X_v^{\mathfrak{g},\sigma}) = v$ and $T_{\pi(v)}M / \mathbb{R}v \cong v^\perp$. With this isomorphism and (4.7-i) in mind, we have

$$(4.15) \quad \text{each } E_v^\sigma(t) \text{ corresponds to the graph of a linear operator } S_v(t) : v^\perp \rightarrow v^\perp$$

which, by (4.8) and (4.11), must be self-adjoint. Now, recall the *Loewner order* on the space of self-adjoint operators on v^\perp : we write $A < B$ if $B - A$ is positive-definite. For this partial order, it is well-known that a *monotone convergence theorem*

holds: a bounded and monotone sequence (or curve) of self-adjoint operators must converge.

LEMMA 4.4. *For every $v \in \Sigma_s$, the limits*

$$(4.16) \quad S_v^+ = \lim_{t \rightarrow +\infty} S_v(t) \quad \text{and} \quad S_v^- = \lim_{t \rightarrow -\infty} S_v(t)$$

exist, and their graphs define $\Psi_t^{(\mathbf{g}, \sigma)}$ -invariant subbundles $E^{\sigma, +}$ and $E^{\sigma, -}$ of Q .

PROOF. The arguments for S_v^+ and S_v^- are identical, so it suffices to show that S_v^+ exists and its graph defines an invariant subbundle. We first show that the limit exists, following the argument in [27]. Assume for simplicity that $s = 1$, so $\Sigma_s = SM$. We claim that

$$(4.17) \quad \begin{aligned} &\text{the difference } S_v(t, s) = S_v(t) - S_v(s) \text{ has constant sig-} \\ &\text{nature in the open regions (i) } 0 < s < t \text{ and (ii) } s < 0 < t. \end{aligned}$$

To verify (4.17-i), assume that there exists $w \in \ker S_v(t_0, s_0) \setminus \{0\}$, for suitable $0 < s_0 < t_0$. Then, the field $J \in \mathcal{J}_0^{\mathbf{g}, \sigma, n}(v)$ with initial conditions $J(0) = w$ and $(\tilde{\mathcal{D}}J)(0) = S_v(t_0)w$ necessarily has $J(t_0)^\perp = J(s_0)^\perp = 0$, due to (4.14) and (4.15). It follows from (4.13) that $\Psi_{s_0}^{(\mathbf{g}, \sigma)}(J) \in \mathcal{J}_0^{\mathbf{g}, \sigma, n}(\Phi_1^{(\mathbf{g}, \sigma)}(s_0, v))$ satisfies $\Psi_{s_0}^{(\mathbf{g}, \sigma)}(J)(0) = 0$ and $[\Psi_{s_0}^{(\mathbf{g}, \sigma)}(J)(t_0 - s_0)]^\perp = 0$, contradicting our assumption that $\Phi_1^{(\mathbf{g}, \sigma)}$ is without conjugate points. The same argument also establishes (4.17-ii).

In view of (4.17-i), to show that $t \mapsto S_v(t)$ is increasing for $t > 0$, it suffices to exhibit $0 < s_0 < t_0$ such that $S_v(t_0, s_0)$ is positive-definite. This is done with a local argument: for every $t > 0$, let $P_{-t}: \dot{\gamma}(t)^\perp \rightarrow v^\perp$ be the $\tilde{\mathcal{D}}$ -parallel transport along γ_v and, for every $w \in v^\perp$ let $J \in E_v^\sigma(t)$ have initial conditions $J(0) = w$ and $(\tilde{\mathcal{D}}J)(0) = S_v(t)w$ — here, we use (4.12). Now the Taylor expansion

$$(4.18) \quad \begin{aligned} 0 &= P_{-t}(J(t)^\perp) = J(0)^\perp + t(\tilde{\mathcal{D}}J^\perp)(0) + \mathcal{O}(t^2) \\ &= w + tS_v(t)w + \mathcal{O}(t^2) = (\text{Id}_{v^\perp} + tS_v(t) + \mathcal{O}(t^2))w \end{aligned}$$

leads to $S_v(t) = -t^{-1}\text{Id}_{v^\perp} + \mathcal{O}(t)$, meaning that $S_v(t, s)$ is positive-definite for $0 < s < t$ sufficiently small.

Finally, it follows from (4.17-ii) that $t \mapsto S_v(t)$ is bounded for $t > 0$ as, for instance, $S_v(t) < S_v(-1)$. The existence of $\lim_{t \rightarrow +\infty} S_v(t)$ is now a consequence of the monotone convergence theorem mentioned after (4.15).

To prove invariance of the bundle, fix $\tau \neq 0$ and $v \in SM$. Observe that if $J \in \mathcal{J}_0^{\mathbf{g}, \sigma, n}(v)$ has initial conditions $J(0) = w$ and $(\tilde{\mathcal{D}}J)(0) = S_v(t)w$, then $\Psi_\tau^{(\mathbf{g}, \sigma)}(J) \in \mathcal{J}_0^{\mathbf{g}, \sigma, n}(\Phi_1^{(\mathbf{g}, \sigma)}(\tau, v))$ is a new Jacobi field satisfying $\Psi_\tau^{(\mathbf{g}, \sigma)}(J)(0) = J(\tau)^\perp$ and $[\Psi_\tau^{(\mathbf{g}, \sigma)}(J)(t - \tau)]^\perp = 0$. This shows that we have

$$(\tilde{\mathcal{D}}\Psi_\tau^{(\mathbf{g}, \sigma)}(J))(0) = S_{\Phi_1^{(\mathbf{g}, \sigma)}(\tau, v)}(t - \tau)J(\tau)^\perp.$$

We let t tend to $+\infty$ to get that $\Psi_\tau^{(\mathbf{g},\sigma)}(E_v^{\sigma,+}) \subseteq E_{\Phi_1^{(\mathbf{g},\sigma)}(\tau,v)}^{\sigma,+}$. The choice of v and τ were arbitrary, so the same argument also shows that $\Psi_{-\tau}^{(\mathbf{g},\sigma)}(E_{\Phi_1^{(\mathbf{g},\sigma)}(\tau,v)}^{\sigma,+}) \subseteq E_v^{\sigma,+}$, and we conclude invariance. \square

5. Proof of Theorem A

Let (\mathbf{g}, σ) be a magnetic system on a closed manifold M , and $s > 0$ be such that $\Phi_s^{(\mathbf{g},\sigma)}: \mathbb{R} \times \Sigma_s \rightarrow \Sigma_s$ is without conjugate points. We use the notation and setup from Section 4. For every $v \in \Sigma_s$ and $t \in \mathbb{R}$, we define an operator $Y_v(t): v^\perp \rightarrow v^\perp$ by $Y_v(t)w = P_{-t}(J_w(t)^\perp)$, where $P_{-t}: \dot{\gamma}(t)^\perp \rightarrow v^\perp$ is the $\tilde{\mathcal{D}}$ -parallel transport operator along γ_v , and $J_w \in \mathcal{J}_0^{\mathbf{g},\sigma,n}(v)$ has the initial conditions $J_w(0) = w$ and $(\tilde{\mathcal{D}}J_w)(0) = S_v^+ w$, for S_v^+ given as in Lemma 4.4. We first claim that

$$(5.1) \quad P_{-\tau} \circ Y_{\Phi_s^{(\mathbf{g},\sigma)}(\tau,v)}(t) \circ P_\tau \circ Y_v(\tau) = Y_v(t + \tau)$$

holds for all $t, \tau \in \mathbb{R}$. Indeed, if we denote by $L_v(t): v^\perp \rightarrow \dot{\gamma}(t)^\perp$ the linear mapping taking w to $J_w(t)^\perp$, the cocycle relation $L_{\Phi_s^{(\mathbf{g},\sigma)}(\tau,v)}(t) \circ L_v(\tau) = L_v(t + \tau)$ obviously holds; substituting $L_v(t) = P_t \circ Y_v(t)$ in it yields (5.1).

By construction of S_v^+ , we have that J_w is nowhere vanishing whenever $w \neq 0$, and thus each $Y_v(t)$ is invertible. This allows us to consider the self-adjoint operator $U_v(t) = \dot{Y}_v(t) \circ Y_v(t)^{-1}$ on v^\perp . Using Proposition 3.1, we have that

$$(5.2) \quad \ddot{Y}_v(t)w = P_{-t}(\tilde{\mathcal{D}}^2(J_w)(t)^\perp) = -P_{-t} \left((M_s^\Omega)_{\Phi_s^{(\mathbf{g},\sigma)}(t,v)/s} (J_w(t)^\perp) \right)$$

or, equivalently, $\ddot{Y}_v(t) = -P_{-t} \circ (M_s^\Omega)_{\Phi_s^{(\mathbf{g},\sigma)}(t,v)/s} \circ L_v(t)$. Thus,

$$(5.3) \quad \begin{aligned} \dot{U}_v(t) &= \ddot{Y}_v(t)Y_v(t)^{-1} - \dot{Y}_v(t)Y_v(t)^{-1}\dot{Y}_v(t)Y_v(t)^{-1} \\ &= -P_{-t} \circ (M_s^\Omega)_{\Phi_s^{(\mathbf{g},\sigma)}(t,v)/s} \circ P_t - U_v(t)^2. \end{aligned}$$

Taking the trace of (5.3), it follows that

$$(5.4) \quad \text{tr}(\dot{U}_v(t)) + \text{tr}(U_v(t)^2) + \text{Ric}_s^\Omega(\Phi_s^{(\mathbf{g},\sigma)}(t,v)/s) = 0.$$

Setting $t = 0$ and integrating both sides of (5.4) with respect to the Liouville measure $d\mu_{\mathbf{g},s} = d\nu_{\mathbf{g}} \otimes dm_s$ induced in Σ_s , where $d\nu_{\mathbf{g}}$ is the Riemannian measure of (M, \mathbf{g}) and dm_s is the Lebesgue measure on a round sphere of radius s , we see that

$$(5.5) \quad \int_{\Sigma_s} \left(\text{tr}(\dot{U}_v(0)) + \text{tr}(U_v(0)^2) \right) d\mu_{\mathbf{g},s}(v) + s^{n-1} \int_{SM} \text{Ric}_s^\Omega(v) d\mu_{\mathbf{g},1}(v) = 0.$$

For the integral of Ric_s^Ω above, note that $dm_s(v) = s^{n-1}dm(w)$ for $w = v/s$ and apply Fubini's theorem.

Letting $F: \Sigma_s \rightarrow \mathbb{R}$ be given by $F(v) = \text{tr}(U_v(0))$, we claim that $\text{tr}(\dot{U}_v(0)) = X^{\mathfrak{g},\sigma}(F)(v)$. Indeed, in view of the relation

$$(5.6) \quad U_{\phi_s^{\mathfrak{g},\sigma}(\tau,v)}(t) = P_\tau \circ U_v(t + \tau) \circ P_{-\tau},$$

which is a direct consequence of (5.1), we have

$$(5.7) \quad X^{\mathfrak{g},\sigma}(F)(v) = \left. \frac{d}{dt} \right|_{t=0} \text{tr} U_{\phi_s^{\mathfrak{g},\sigma}(t,v)}(0) = \left. \frac{d}{dt} \right|_{t=0} \text{tr} U_v(t) = \text{tr} \dot{U}_v(0).$$

As the magnetic flow is volume-preserving, cf. [34, Proposition 1.5], it follows that the integral of $\text{tr}(\dot{U}_v(0))$ over Σ_s vanishes, and so (5.5) leads us to

$$(5.8) \quad \int_{SM} \text{Ric}_s^\Omega(v) d\mu_{\mathfrak{g},1}(v) = -s^{1-n} \int_{\Sigma_s} \text{tr}(U_v(0)^2) d\mu_{\mathfrak{g},s}(v) \leq 0,$$

concluding the first part of the argument.

Assuming now that the equality holds in (5.8), measurability and non-negativity of the function $\Sigma_s \ni v \mapsto \text{tr}(U_v(0)^2)$ tells us that $\text{tr}(U_v(0)^2) = 0$ for almost every $v \in \Sigma_s$. As $U_v(0)$ is self-adjoint, it must then be the case that $U_v(0) = 0$ for almost every $v \in \Sigma_s$. Using that the flow is volume preserving along with (5.6), we see that for every $t \in \mathbb{R}$ the set $B_t = \{v \in \Sigma_s \mid U_v(t) = 0\}$ has full $\mu_{\mathfrak{g},s}$ -measure. Hence, $B = B_0 \cap (\bigcap_{n \geq 1} B_{1/n})$ also has full measure, so for almost every $v \in \Sigma_s$ we have $\dot{U}_v(0) = \lim_{n \rightarrow \infty} n[U_v(1/n) - U_v(0)] = 0$. Substituting this into (5.3) with $t = 0$, we obtain that $M_s^\Omega = 0$ as required.

6. The meaning of magnetic flatness and proof of Theorem D

Recall that a Kähler manifold may be defined as a Riemannian manifold (M, \mathfrak{g}) equipped with an almost-complex structure J such that $\mathfrak{g}(J \cdot, J \cdot) = \mathfrak{g}$ and $\nabla J = 0$, with ∇ denoting the Levi-Civita connection of (M, \mathfrak{g}) . This latter condition then implies that J is integrable. Restricting the sectional curvature function of (M, \mathfrak{g}) to J -invariant planes, one has the well-known notion of *holomorphic sectional curvature*, which may be regarded as the smooth function $\text{sec}^{\text{hol}}: SM \rightarrow \mathbb{R}$ given by $\text{sec}^{\text{hol}}(v) = \text{sec}(v, Jv)$. Note that sec^{hol} also equals the Gaussian curvature of (M, \mathfrak{g}) when $\dim M = 2$.

Given any $\lambda \in (0, \infty)$, we may consider the magnetic system $(\mathfrak{g}, \lambda\omega)$ on M , where $\omega = \mathfrak{g}(J \cdot, \cdot)$ is the Kähler form of (M, \mathfrak{g}) . Systems of this form are called *Kähler magnetic systems*, and they have been extensively studied by Adachi [1–4, 33].

Whenever the holomorphic sectional curvature of (M, \mathfrak{g}) is constant and equal to k , the Riemann curvature tensor is given by

$$(6.1) \quad R(X, Y)Z = \frac{k}{4}(\langle Y, Z \rangle X - \langle X, Z \rangle Y - \langle Y, JZ \rangle JX + \langle X, JZ \rangle JY + 2\langle X, JY \rangle JZ),$$

for all vector fields X, Y , and Z on M (cf. [28, Proposition 7.3, p. 167]). In this case, a straightforward computation shows that

$$(6.2) \quad (\sec_s^\Omega)_v(w) = \frac{s^2k + \lambda^2}{4}(1 + 3\langle v, Jw \rangle^2),$$

for all $s > 0$ and all pairs of orthonormal vectors v, w tangent to M . In the case where $\dim M = 2$, (6.2) reads $\sec_s^\Omega \equiv s^2k + \lambda^2$.

Our Theorem D serves as a rough “converse” of the above discussion.

PROOF OF THEOREM D. Combining (1.6) with the easily-verified relation

$$(6.3) \quad (\sec_s^\Omega)_v(w) = s^2\sec(v, w) - s\langle (\nabla_w \Omega)(v), w \rangle + \langle (A^\Omega)_v(w), w \rangle,$$

the constant \sec_s^Ω condition can be written as

$$(6.4) \quad s^2\sec(v, w) - s\langle (\nabla_w \Omega)(v), w \rangle + \frac{3}{4}\langle v, \Omega(w) \rangle^2 + \frac{1}{4}\|\Omega(w)\|^2 = c,$$

for all pairs of orthonormal vectors $v, w \in TM$. We first claim that

$$(6.5) \quad (\nabla_w \Omega)(w) = 0, \text{ for every } w \in TM.$$

To establish (6.5), we may assume without loss of generality that $\|w\| = 1$. Replacing $v \mapsto -v$ in (6.4) and invoking skew-adjointness of $\nabla_w \Omega$ immediately yields $\langle (\nabla_w \Omega)(w), v \rangle = 0$, for all unit vectors v orthogonal to w . This forces $(\nabla_w \Omega)(w)$ to be proportional to w , while at the same time $(\nabla_w \Omega)(w)$ is orthogonal to w ; (6.5) thus holds.

Polarizing (6.5), it follows that $\nabla \Omega$ is skew-symmetric, and as a consequence $\nabla \sigma$ is alternating. However, the exterior derivative of a differential form equals the alternator of its covariant differential, up to a dimensional constant depending on conventions. Therefore, $\nabla \sigma = 0$ follows from $d\sigma = 0$.

As a nontrivial parallel tensor field on any connected manifold equipped with an affine connection is necessarily nowhere-vanishing, we now claim that

$$(6.6) \quad \begin{aligned} &\Omega \text{ is a conformal endomorphism of } TM, \text{ and} \\ &\text{thus } \|\Omega\|^{-1}\Omega \text{ is a complex structure on } M. \end{aligned}$$

Indeed, replacing $(v, w) \mapsto (w, v)$ in (6.4) gives us that $\|\Omega(v)\| = \|\Omega(w)\|$ for all pairs of orthonormal vectors v, w tangent to M , and such condition immediately implies that Ω is conformal, with constant conformal factor $\|\Omega\| > 0$. Namely, we have that $\langle \Omega(v), \Omega(w) \rangle = \|\Omega\|^2 \langle v, w \rangle$ for all vectors v, w tangent to M . As Ω is skew-adjoint, nondegeneracy of g now implies that $\Omega^2 = -\|\Omega\|^2 \text{Id}$, yielding (6.6).

Since Ω is parallel, we have that $J = \|\Omega\|^{-1}\Omega$ is parallel and preserves g , so (6.6) makes (M, g) a Kähler manifold. Let \sec^{hol} be its holomorphic sectional

curvature. Setting $w = Jv$ in (6.4) leads to $s^2 \sec^{\text{hol}}(v) + \|\Omega\|^2 = c$ — yielding (i) — and so (6.2) with $\lambda = -\|\Omega\|$ implies that

$$(6.7) \quad \frac{c}{4}(1 + 3\langle v, Jw \rangle^2) = c,$$

for all pairs of orthonormal vectors v, w tangent to M . Now, it follows from (6.7) that $c = 0$ whenever $\dim M > 2$. Indeed, if $c \neq 0$, then $1 + 3\langle v, Jw \rangle^2 = 4$ leads to $|\langle v, Jw \rangle| = 1$, and the equality case in the Cauchy-Schwarz inequality implies that

$$(6.8) \quad \begin{array}{l} \text{the set } \{v, Jw\} \text{ is linearly dependent, for all} \\ \text{pairs of orthonormal vectors } v, w \text{ tangent to } M. \end{array}$$

If $\dim M \geq 3$, (6.8) easily leads to a contradiction by fixing w and choosing two mutually orthogonal unit vectors v, v' , both orthogonal to w .

□

Appendix A. The Mañé critical value of Kähler magnetic systems

The goal of this appendix is to mimic the proof of [16, Lemma 6.1] in order to establish (1.12). Namely:

LEMMA A.1. *Let (M, \mathfrak{g}) be a compact Kähler manifold with Kähler form ω , whose holomorphic sectional curvature is constant and equal to $k < 0$. Then for every constant $\lambda \in \mathbb{R}$, the Mañé critical value of the Kähler magnetic system $(\mathfrak{g}, \lambda\omega)$ is given by $s_0(\mathfrak{g}, \lambda\omega) = |\lambda|/\sqrt{-k}$.*

Here, we will use an alternative description of the Mañé critical value (see [16, Section 5.1]). Let (\mathfrak{g}, σ) be any magnetic system on M , $s > 0$, and $p: \tilde{M} \rightarrow M$ be the universal covering of M . If $p^*\sigma$ is exact, we consider the Lagrangian $L_s: T\tilde{M} \rightarrow \mathbb{R}$ given by

$$(A.1) \quad L_s(v) = \frac{1}{2}(\|v\|^2 + s^2) - \theta(v),$$

where $\|\cdot\|$ is the norm induced by $p^*\mathfrak{g}$ and θ is any primitive of $p^*\sigma$. Writing $\Lambda\tilde{M}$ for the space of absolutely continuous loops in \tilde{M} , we also consider the action functional $A_s: \Lambda\tilde{M} \rightarrow \mathbb{R}$ given by

$$(A.2) \quad A_s(\gamma) = \int_0^T L_s(\dot{\gamma}(t)) dt,$$

where $T > 0$ is the smallest period of γ . Evidently, (A.2) is independent of the choice of primitive θ . With this in place, we have that

$$(A.3) \quad s_0(\mathfrak{g}, \sigma) = \begin{cases} \inf\{s > 0 \mid A_s(\gamma) \geq 0 \text{ for all } \gamma \in \Lambda\tilde{M}\} & \text{if } p^*\sigma \text{ is exact,} \\ +\infty & \text{otherwise.} \end{cases}$$

PROOF OF LEMMA A.1. As it follows from (1.11) that

$$(A.4) \quad s_0(-k\mathfrak{g}, \lambda\omega) = \frac{|\lambda|}{\sqrt{-k}} s_0(\mathfrak{g}, \omega),$$

it suffices to prove that $s_0(\mathfrak{g}, \omega) = 1$ under the assumption that $k = -1$. By [28, Theorems 7.8 and 7.9], the universal covering of M is holomorphically isometric to the unit ball $\mathbb{B}^n = \{z \in \mathbb{C}^n \mid \|z\| < 1\}$ equipped with the Kähler metric

$$(A.5) \quad \tilde{\mathfrak{g}} = -4 \sum_{j,k=1}^n \left(\frac{(1 - \|z\|^2)\delta_{jk} + \bar{z}_j z_k}{(1 - \|z\|^2)^2} \right) dz_j d\bar{z}_k.$$

The Kähler form $\tilde{\omega}$ of $(\mathbb{B}^n, \tilde{\mathfrak{g}})$ is given by

$$(A.6) \quad \tilde{\omega} = -4i \sum_{j,k=1}^n \left(\frac{(1 - \|z\|^2)\delta_{jk} + \bar{z}_j z_k}{(1 - \|z\|^2)^2} \right) dz_j \wedge d\bar{z}_k,$$

and admits

$$(A.7) \quad \theta = \frac{-2i}{1 - \|z\|^2} \sum_{j=1}^n (\bar{z}_j dz_j - z_j d\bar{z}_j)$$

as a primitive. A routine computation using (A.5), (A.7), and (1.11), shows that $\|\theta_z\| = \|z\| < 1$ for all $z \in \mathbb{B}^n$, so that $s_0(\mathbf{g}, \omega) \leq \sup_{z \in \mathbb{B}^n} \|\theta_z\| = 1$. It remains to establish the reverse inequality. To do so, we consider the totally geodesic embedding $\iota: \mathbb{B}^1 \rightarrow \mathbb{B}^n$ given by $\iota(\zeta) = (\zeta, 0, \dots, 0)$. From [22, Theorem 3.19], we have

$$(A.8) \quad \iota^* \tilde{\mathbf{g}} = -\frac{4 d\zeta d\bar{\zeta}}{(1 - |\zeta|^2)^2} \quad \text{and} \quad \iota^* \tilde{\omega} = -\frac{4i d\zeta \wedge d\bar{\zeta}}{(1 - |\zeta|^2)^2}.$$

Let $s > 0$. For each $r > 0$, let $\gamma_r: [0, \sinh r] \rightarrow \mathbb{B}^1$ be the circle of radius r with respect to $\iota^* \tilde{\mathbf{g}}$, parametrized with constant speed s . Denoting by $\ell(\gamma_r) = 2\pi \sinh r$ and $\mathfrak{a}(D_r) = 2\pi(\cosh r - 1)$ the $\iota^* \tilde{\mathbf{g}}$ -length of γ_r and the $\iota^* \tilde{\mathbf{g}}$ -area of the disk bounded by γ_r , respectively, we proceed to compute the action (A.2) with the aid of θ in (A.7) and Stokes' theorem:

$$(A.9) \quad \begin{aligned} A_s(\gamma_r) &= \int_0^{\ell(\gamma_r)/s} \frac{1}{2} \left((\iota^* \tilde{\mathbf{g}})(\dot{\gamma}_r(t), \dot{\gamma}_r(t)) + s^2 \right) - (\iota^* \theta)(\dot{\gamma}_r(t)) dt \\ &= s\ell(\gamma_r) - \mathfrak{a}(D_r) \\ &= 2\pi(s \sinh r - \cosh r + 1). \end{aligned}$$

It follows from (A.9) that if $s < 1$, the value $A_s(\gamma_r)$ becomes negative for sufficiently large r . We conclude from (A.3) that $s_0(\mathbf{g}, \omega) \geq 1$, as desired. \square

References

- [1] T. Adachi. Kähler magnetic flows for a manifold of constant holomorphic sectional curvature. *Tokyo J. Math.*, 18(2):473–483, 1995.
- [2] T. Adachi. A comparison theorem on magnetic Jacobi fields. *Proc. Edinburgh Math. Soc.* (2), 40(2):293–308, 1997.
- [3] T. Adachi. A theorem of Hadamard-Cartan type for Kähler magnetic fields. *J. Math. Soc. Japan*, 64(3):969–984, 2012.
- [4] T. Adachi. A comparison theorem on harp-sectors for Kähler magnetic fields. *Southeast Asian Bull. Math.*, 38(5):619–626, 2014.
- [5] T. Adachi and P. Bai. An estimate of the spread of trajectories for Kähler magnetic fields. *Hokkaido Math. J.*, 42(3):445–462, 2013.
- [6] D. Anosov and Y. Sinai. Some smooth ergodic systems. *Russian Mathematical Surveys*, 22(5):103, 1967.
- [7] M. Arnaud. Fibrés de Green et régularité des graphes C^0 -lagrangiens invariants par un flot de Tonelli. 9(5):881–926, 2008.
- [8] V. I. Arnol'd. Some remarks on flows of line elements and frames. *Sov. Math., Dokl.*, 2:562–564, 1961.

- [9] L. Asselle and G. Benedetti. The Lusternik–Fet theorem for autonomous Tonelli Hamiltonian systems on twisted cotangent bundles. *Journal of Topology and Analysis*, 8(03):545–570, 2016.
- [10] V. Assenza. Magnetic curvature and existence of closed magnetic geodesics on low energy levels. Preprint available at <https://arxiv.org/abs/2309.03159>.
- [11] A. Bahri and I. A. Taimanov. Periodic orbits in magnetic fields and Ricci curvature of Lagrangian systems. *Trans. Amer. Math. Soc.*, 350(7):2697–2717, 1998.
- [12] M. Bialy. Rigidity for periodic magnetic fields. *Ergodic Theory and Dynamical Systems*, 20(6):1619–1626, 2000.
- [13] M. Bialy and R. MacKay. Symplectic twist maps without conjugate points. *Israel Journal of Mathematics*, 141:235–247, 2004.
- [14] D. Burago and S. Ivanov. Riemannian tori without conjugate points are flat. *Geom. Funct. Anal.*, 4(3):259–269, 1994.
- [15] K. Burns and G. P. Paternain. Anosov magnetic flows, critical values and topological entropy. *Nonlinearity*, 15(2):281–314, 2002.
- [16] K. Cieliebak, U. Frauenfelder, and G. P. Paternain. Symplectic topology of Mañé’s critical values. *Geometry & Topology*, 14(3):1765–1870, 2010.
- [17] G. Contreras. The Palais–Smale condition on contact type energy levels for convex Lagrangian systems. *Calculus of Variations and Partial Differential Equations*, 27(3):321–395, 2006.
- [18] G. Contreras and R. Iturriaga. Convex Hamiltonians without conjugate points. *Ergodic Theory Dyn. Syst.*, 19(4):901–952, 1999.
- [19] G. Contreras, R. Iturriaga, G. P. Paternain, and M. Paternain. The palais-smale condition and mañé’s critical values. In *Annales Henri Poincaré*, volume 1, pages 655–684. Springer, 2000.
- [20] P. Foulon. Estimation de l’entropie des systèmes lagrangiens sans points conjugués. 57(2):117–146, 1992.
- [21] V. Ginzburg. A charge in a magnetic field: Arnold’s problems 1981–9, 1982–24, 1984–4, 1994–14, 1994–35, 1996–17, 1996–18. *Arnold’s problems*, pages 395–401, 2004.
- [22] W. Goldman. *Complex hyperbolic geometry*. Oxford Math. Monogr. Oxford: Clarendon Press, 1999.
- [23] N. Gouda. The theorem of E. Hopf under uniform magnetic fields. *J. Math. Soc. Japan*, 50(3):767–779, 1998.
- [24] L. W. Green. A theorem of E. Hopf. *Michigan Math. J.*, 5:31–34, 1958.
- [25] P. Herreros. Scattering boundary rigidity in the presence of a magnetic field. *Comm. Anal. Geom.*, 20(3):501–528, 2012.
- [26] E. Hopf. Closed surfaces without conjugate points. *Proc. Nat. Acad. Sci. U.S.A.*, 34:47–51, 1948.
- [27] R. Iturriaga. A geometric proof of the existence of the Green bundles. *Proc. Amer. Math. Soc.*, 130(8):2311–2312, 2002.
- [28] S. Kobayashi and K. Nomizu. *Foundations of differential geometry. Vol. II*, volume Vol. II of *Interscience Tracts in Pure and Applied Mathematics, No. 15*. Interscience Publishers John Wiley & Sons, Inc., New York-London-Sydney, 1969.
- [29] W. Merry. Closed orbits of a charge in a weakly exact magnetic field. *Pacific journal of mathematics*, 247(1):189–212, 2010.
- [30] G. P. Paternain. *Geodesic flows*, volume 180 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, 1999.
- [31] G. P. Paternain and M. Paternain. Anosov geodesic flows and twisted symplectic structures. In *International Conference on Dynamical Systems (Montevideo, 1995)*, volume 362 of *Pitman Res. Notes Math. Ser.*, pages 132–145. Longman, Harlow, 1996.

- [32] F. Schlenk. Applications of hofer's geometry to hamiltonian dynamics. *Commentarii Mathematici Helvetici*, 81(1):105–121, 2006.
- [33] Q. Shi and T. Adachi. Comparison theorems on trajectory-harps for Kähler magnetic fields which are holomorphic at their arches. *Hokkaido Math. J.*, 48(2):427–441, 2019.
- [34] M. P. Wojtkowski. Magnetic flows and Gaussian thermostats on manifolds of negative curvature. *Fund. Math.*, 163(2):177–191, 2000.

INSTITUTO DE MATEMÁTICA PURA E APLICADA, RIO DE JANEIRO, RJ 22460-320, BRAZIL

Email address: `valerio.assenza@impa.br`

DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY, COLUMBUS, OH 43210, USA

Email address: `marshallreber.1@osu.edu`

Email address: `terekcouth.1@osu.edu`