

Name _____ KEY _____ Lab TA/time _____

1. (5 pts) Calculate the **mass percent** composition of **ALL elements** in $\text{Ca}_3(\text{PO}_4)_2$.
(At. wts: Ca = 40.08, P = 30.97, O = 16.00) (**Must show all work.**)

$$\text{Mass \% of element} = \frac{\text{mass element}}{\text{total mass}} \times 100\%$$

$$\text{From a molecular or : empirical formula} \quad \text{mass \%} = \frac{\text{mass element}}{\text{MW (or FW)}} \times 100\%$$

$$\text{Ca : } 3 \times 40.08 \text{ amu} = 120.24 \text{ amu}$$

$$\text{P : } 2 \times 30.97 \text{ amu} = 61.94 \text{ amu}$$

$$\text{O : } 8 \times 16.00 \text{ amu} = 128.00 \text{ amu}$$

$$\text{FW for } \text{Ca}_3(\text{PO}_4)_2 = \frac{\quad}{\quad} = 310.18 \text{ amu (5 s.f.)}$$

$$\% \text{ Ca} = \frac{120.24 \text{ amu}}{310.18 \text{ amu}} \times 100 \% = 38.765 \% \text{ Ca (5 s.f.)}$$

$$\% \text{ P} = \frac{61.94 \text{ amu}}{310.18 \text{ amu}} \times 100 \% = 19.97 \% \text{ P (4 s.f.)}$$

$$\% \text{ O} = \frac{128.00 \text{ amu}}{310.18 \text{ amu}} \times 100 \% = 41.266 \% \text{ O (5 s.f.)}$$

2. (3 pts) Cisplatin, an anticancer drug, has the molecular formula $\text{Pt}(\text{NH}_3)_2\text{Cl}_2$. How many moles of hydrogen atoms are in 2.8×10^{-4} g of cisplatin?
(At. Wts.: H = 1.008, N = 14.01, Cl = 35.45, Pt = 195.1 ; Mol. wt: 300.07)

$$1 \text{ mol Pt}(\text{NH}_3)_2\text{Cl}_2 = 6.02 \times 10^{23} \text{ Pt}(\text{NH}_3)_2\text{Cl}_2 \text{ molecules} = 300.07 \text{ g Pt}(\text{NH}_3)_2\text{Cl}_2$$

$$\text{Also, } 1 \text{ mol Pt}(\text{NH}_3)_2\text{Cl}_2 = 6 \text{ mol H atoms}$$

$$\begin{aligned} ? \text{ mol H atoms} &= 2.8 \times 10^{-4} \text{ g Pt}(\text{NH}_3)_2\text{Cl}_2 \times \frac{1 \text{ mol Pt}(\text{NH}_3)_2\text{Cl}_2}{300.07 \text{ g Pt}(\text{NH}_3)_2\text{Cl}_2} \times \frac{6 \text{ mol H atoms}}{1 \text{ mol Pt}(\text{NH}_3)_2\text{Cl}_2} \\ &= 5.5986 \times 10^{-6} \text{ moles H atoms} \\ &= 5.6 \times 10^{-6} \text{ moles H atoms} \end{aligned}$$

3. (3 pts) Elements represented by atomic symbols **A** and **Z** form molecular compounds **AZ** (g) and **A₂Z** (l). For 6.00 g samples of each of the two compounds at 25°C,
- one can state that there are the same number of molecules in both samples.
 - one can state that there are fewer molecules of **AZ** than **A₂Z**.
 - * one can state that there are more molecules of **AZ** than **A₂Z**.
 - one can not make a statement about the number of molecules because the states of the samples are different.
 - one can not make a statement about the number of molecules because the atomic weights of **A** and **Z** are not given.

There are two molecules **AZ**(g) and **A₂Z**(l) composed of atoms **A** and **Z** (the states of the two molecules is immaterial). There are 6.00 g of each. Based on the formulas the **A₂Z** molecule has a larger MW (molar mass). The molar mass of a substance is inversely proportional to the moles ($\text{MM} = \text{g/mol}$) and thus the moles is inversely proportional to the MM,

$$\text{MM} = \text{g/mol}$$

$$\text{mol} = \text{g/MM}$$

Thus, since the mass of each cmpd is the same (6.00 g) the substance with the smaller molar mass will have the larger number of moles and thus the larger number of molecules (since molecules and moles are directly proportional). **AZ** has a smaller molar mass than **A₂Z** and thus **AZ** has the larger number of moles and molecules.

4. (7 pts) An analysis of a compound containing only carbon and fluorine gives a mass percent composition of 21.32% C and 78.68% F. The experimentally determined molecular weight is 507 amu. (At. Wt.: C = 12.011, F = 18.998)

Must determine empirical formula first

Assume a 100 g sample

21.32 g C 78.68 g F

Step 1: convert to moles

$$\text{C: } 21.32 \text{ g C} \times \frac{1 \text{ mole C}}{12.011 \text{ g C}} = 1.77\textbf{50} \text{ mol C (4 s.f.)}$$

$$\text{F: } 78.68 \text{ g F} \times \frac{1 \text{ mole F}}{18.998 \text{ g F}} = 4.14\textbf{14} \text{ mol F (4 s.f.)}$$

Step 2: find mole ratio (divide each by smallest # moles)- this is also atom ratio in empirical formula.

$$\text{C: } \frac{1.77\textbf{50}}{1.77\textbf{50}} = 1.000$$

$$\text{F: } \frac{4.14\textbf{14 mol F}}{1.77\textbf{50 mol C}} = 2.333$$

$\text{C}_1 \text{F}_{2.333}$ can't have fractional subscripts, so multiply subscripts by 3
(if you try multiplying by 2 you are not close enough to a whole number - you get 4.666 F atoms)

EF is C_3F_7

EFW = 169.02 amu (empirical formula weight)

b) (2 pts) What is the **molecular formula**? Not asked for on the quiz.

Step 3: determine the molecular formula. The MF is always an integer multiple of the empirical formula, EF. Thus the MW is the same multiple of the EFW.

$$\text{MW} = n \times \text{EFW}$$

$$n = \frac{\text{MW}}{\text{EFW}} = \frac{507 \text{ amu}}{169.02 \text{ amu}} = 2.99 = 3$$

$$\text{MF} = (\text{EF})_n = (\text{C}_3\text{F}_7)_3 = \boxed{\text{C}_9\text{F}_{21}}$$

5. (7 pts) A 0.589 g sample of an organic compound containing only carbon, hydrogen and oxygen was burned completely in air to produce 0.733 g of CO₂ and 0.299 g of H₂O. What is the empirical formula of the compound? (Atomic weights: C = 12.01, H = 1.008, O = 16.00)

This is an empirical formula problem using combustion analysis. In CA the sample undergoes complete combustion. All the Carbon winds up in the CO₂ and the Hydrogen winds up in the H₂O. If there is one other atom its mass can be determined by remembering the conservation of mass, in this case oxygen.

In this problem a 0.589 g sample produces 0.733 g of CO₂ and 0.299 g H₂O.

Can do this in a couple of ways. In lecture I converted the masses of CO₂ and H₂O into the masses of C and H atoms present in the original sample by using mass fractions. Then I got the mass of the third atom (in this case oxygen) by using the Law of Conservation of mass. Then I converted all the masses to moles.

Here I'm converting the masses of CO₂ and H₂O into the moles of C and H atoms present in the original sample since we need that anyway and then converting that to the masses of C and H atoms present in the original sample. Then those are used to obtain the mass of oxygen which is then converted to moles of oxygen.

Determine moles and mass of C and H in the sample:

$$? \text{ mol C} = 0.733 \text{ g CO}_2 \times \frac{1 \text{ mol CO}_2}{44.01 \text{ g CO}_2} \times \frac{1 \text{ mol C}}{1 \text{ mol CO}_2} = 0.016\textbf{65} \text{ mol C (3 s.f.)}$$

$$? \text{ mol H} = 0.299 \text{ g H}_2\text{O} \times \frac{1 \text{ mol H}_2\text{O}}{18.016 \text{ g H}_2\text{O}} \times \frac{2 \text{ mol H}}{1 \text{ mol H}_2\text{O}} = 0.033\textbf{19} \text{ mol H (3 s.f.)}$$

Need moles of oxygen but have to get mass first. Need to calculate mass of C and H and subtract from the total mass of sample to get the mass of oxygen.

$$? \text{ g C} = 0.016\textbf{65} \text{ mol C} \times \frac{12.01 \text{ g C}}{1 \text{ mol C}} = 0.20\textbf{00} \text{ g C (3 s.f.)}$$

$$? \text{ g H} = 0.033\textbf{19} \text{ mol H} \times \frac{1.008 \text{ g H}}{1 \text{ mol H}} = 0.033\textbf{45} \text{ g H (3 s.f.)}$$

$$? \text{ g O} = 0.589 \text{ g sample} - (0.20\textbf{00} \text{ g C} + 0.033\textbf{45} \text{ g H}) = 0.35\textbf{55} \text{ g O}$$

Find moles of O:

$$? \text{ mol O} = 0.35\textbf{55} \text{ g O} \times \frac{1 \text{ mol O}}{16.00 \text{ g O}} = 0.022\textbf{21} \text{ mol O (3 s.f.)}$$

***** continued on next page *****

5. (Cont.)

Empirical Formula calculations:


Divide each of the moles by the smallest number of moles (in this case O).

$$\text{C: } \frac{0.016\textbf{65} \text{ mol C}}{0.016\textbf{65} \text{ mol C}} = 1.00 \times 3 = 3.00 = 3$$


$$\text{H: } \frac{0.033\textbf{45} \text{ mol H}}{0.016\textbf{65} \text{ mol C}} = 1.99 \times 3 = 5.97 = 6$$

$$\text{O: } \frac{0.022\textbf{21} \text{ mol O}}{0.016\textbf{65} \text{ mol C}} = 1.33 \times 3 = 3.99 = 4$$

Can't have non-integer ratios (subscripts). Have to multiply by a factor to get integers. Must multiply each one by 3 (multiplying by 2 won't give a number close enough to an integer for oxygen).




6. (4 pts) Given the balanced equation below, how many moles of hydrogen can be produced from the complete reaction of 3.860×10^{-1} mol of Fe with excess water? (At. Wts.: H = 1.008, O = 16.00, Fe = 55.85)



mole-to-mole stoichiometry problem

$$? \text{ mol H}_2 = 0.386\textbf{0} \text{ mol Fe} \times \frac{4 \text{ mol H}_2}{3 \text{ mol Fe}} = 0.514666 = 0.5147 \text{ mole H}_2.$$

7. (5 pts) How many **grams** of oxygen (O_2), reacting with excess C_2H_6 , are required to form 35.0 g of carbon dioxide (CO_2), according to the following equation? (At. Wt.: H = 1.01, O = 16.00, C = 12.01; Mol. Wt: C_2H_6 = 30.08, O_2 = 32.00, CO_2 = 44.01, H_2O = 18.02)




gram \rightarrow gram stoich

$$\begin{aligned} ? \text{ g O}_2 &= 35.0 \text{ g CO}_2 \times \frac{1 \text{ mol CO}_2}{44.01 \text{ g CO}_2} \times \frac{7 \text{ mol O}_2}{4 \text{ mol CO}_2} \times \frac{32.00 \text{ g O}_2}{1 \text{ mol O}_2} = 44.\textbf{535} \text{ g O}_2 \\ &= 44.5 \text{ g O}_2 \\ &\quad (3 \text{ s.f.}) \end{aligned}$$

$$\frac{\text{grams desired}}{\text{grams desired}} = \frac{\text{grams given}}{\text{grams given}} \times \frac{\text{moles given}}{\text{grams given}} \times \frac{\text{moles desired}}{\text{moles given}} \times \frac{\text{grams desired}}{\text{moles desired}}$$

$$\text{grams given} \Rightarrow \text{mol given} \Rightarrow \text{mol desired} \Rightarrow \text{grams desired}$$

8. (6 pts) Calcium hydroxide reacts with phosphoric acid according to the following equation. Which substance is the limiting reagent when 1.00 mol of Ca(OH)_2 reacts with 0.50 mol of H_3PO_4 ? How many moles of the excess reagent remain after completion of the reaction?



This is a limiting reactant problem. These are just stoichiometry problems. There is more than one way to do a LR problem. In this case since it's asking for the limiting reactant and the mass of excess reactant remaining after completion of the reaction.

1) Method 1: Calculate which reactant gives the smallest number of moles of product, and then use to determine how much of the excess reactant would be used (and remains).

Calculate mol $\text{Ca}_3(\text{PO}_4)_2$ from Ca(OH)_2 and H_3PO_4 :

$$? \text{ mol } \text{Ca}_3(\text{PO}_4)_2 = 1.00 \text{ mol } \text{Ca(OH)}_2 \times \frac{1 \text{ mol } \text{Ca}_3(\text{PO}_4)_2}{3 \text{ mol } \text{Ca(OH)}_2} = 0.33\bar{3} \text{ mol } \text{Ca}_3(\text{PO}_4)_2 \text{ (3 s.f.)}$$

$$? \text{ mol } \text{Ca}_3(\text{PO}_4)_2 = 0.50 \text{ mol } \text{H}_3\text{PO}_4 \times \frac{1 \text{ mol } \text{Ca}_3(\text{PO}_4)_2}{2 \text{ mol } \text{H}_3\text{PO}_4} = 0.2500 \text{ mol } \text{Ca}_3(\text{PO}_4)_2 \text{ (3 s.f.) } \mathbf{LR}$$

Fewer moles of $\text{Ca}_3(\text{PO}_4)_2$ obtained from the H_3PO_4 so the $\mathbf{H}_3\text{PO}_4$ is the **LR** and only 0.2500 moles of $\text{Ca}_3(\text{PO}_4)_2$ are formed. Calculate the moles of Ca(OH)_2 required to produce 0.2500 mol of $\text{Ca}_3(\text{PO}_4)_2$

$$? \text{ mol } \underset{\text{used}}{\text{Ca(OH)}_2} = 0.2500 \text{ mol } \text{Ca}_3(\text{PO}_4)_2 \times \frac{3 \text{ mol } \text{Ca(OH)}_2}{1 \text{ mol } \text{Ca}_3(\text{PO}_4)_2} = 0.750 \text{ mol } \text{Ca(OH)}_2 \text{ used (2 s.f.)}$$

$$? \text{ mol } \text{Ca(OH)}_2 = 1.00 \text{ mol } \text{Ca(OH)}_2 - 0.750 \text{ mol } \text{Ca(OH)}_2 = 0.250 \text{ mol } \text{Ca(OH)}_2 \text{ remaining}$$

H_3PO_4 is LR 0.25 mol Ca(OH)_2 remaining

2) Method 2: Determine mole ratio of the two reactants & compare it to the mole ratio in the bal. eqn.

$$\frac{1.00 \text{ mol } \text{Ca(OH)}_2}{0.50 \text{ mol } \text{H}_3\text{PO}_4} > \frac{3 \text{ mol } \text{Ca(OH)}_2}{2 \text{ mol } \text{H}_3\text{PO}_4}$$

Since the actual ratio of Ca(OH)_2 to H_3PO_4 (ratio = 2) is greater than that from the bal. eqn. (ratio = 1.5) this means Ca(OH)_2 is in excess and H_3PO_4 is the LR. Can then calculate the moles of Ca(OH)_2 required to react with the H_3PO_4 .

$$? \text{ mol } \text{Ca(OH)}_2 = 0.50 \text{ mol } \text{H}_3\text{PO}_4 \times \frac{3 \text{ mol } \text{Ca(OH)}_2}{2 \text{ mol } \text{H}_3\text{PO}_4} = 0.750 \text{ mol } \text{Ca(OH)}_2 \text{ used (3 s.f.)}$$

9. (3 pts) Which of the following are **strong electrolytes**?

HF HCl Cu(ClO₃)₂ Ca(OH)₂ C₂H₅OH

All soluble ionic substances (salts) are strong electrolytes. All strong acids and bases are strong electrolytes. You need to memorize these. All other molecular substances that dissolve are either weak or non-electrolytes.

The 7 strong acids are: HCl, HBr, HI binary acids

HNO₃, HClO₃, HClO₄, H₂SO₄ (1st H⁺ only) oxyacids (ternary)

Strong bases are group 1 A and 2 A hydroxides are the most common. We will learn others in Ch 16.

The following are strong electrolytes:

HCl strong acid

Cu(ClO₃)₂ soluble ionic compd.

Ca(OH)₂ strong base (ionic as well)

The following are weak electrolytes:

HF weak acid

C₂H₅OH molecular (nonelectrolyte)