

# Chapter 7

## Periodic Properties of the Elements

### I) Development of the P.T.

Generally, the electronic structure of atoms correlates w. the prop. of the elements

- reflected by the arrangement of the elements in the P.T.

A number of elements were discovered based on expected prop. of the “missing” elements.

## A) Noble Gases

$ns^2np^6$  - very stable


## B) Representative Elements

“last”  $e^-$  added to s & p orbitals

distinct & fairly regular variations  
in prop. w. changes in atomic #

## C) d-Transition Elements

$e^-$  added to d orbitals



## II) Effective Nuclear Charge

Net (+) charge attracting an  $e^-$

$$Z_{\text{eff}} = Z - S$$

$S$  = screening constant

- avg. number of  $e^-$ 's between nucleus & any particular  $e^-$
- depends on specific orbitals

Subsets of  $e^-$ :

- 1) core  $e^-$
- 2) valence  $e^-$

Inner  $e^-$  screen or shield outer  $e^-$  from full (+) charge


Primary interaction of  $e^-$  & nucleus is due to charge:

Coulomb's Law:


$$F = \frac{k (Q_e Q_n)}{r^2}$$

However, valence shell  $e^-$  do not experience full nuclear charge

- partially shielded by the core  $e^-$



(a)



(b)

Copyright © 2006 Pearson Prentice Hall, Inc.

Value of  $S$  is usually close to # core  $e^-$



Valence shell  $e^-$  do not screen each other effectively.

- same distance from nucleus

The “p”  $e^-$  do not screen “s”  $e^-$

The “s”  $e^-$  do screen “p”  $e^-$  somewhat due to a probability for these  $e^-$  to be nearer the nucleus

- penetration



Copyright © 2006 Pearson Prentice Hall, Inc.

## A) General Trends

### 1) Across Row

$Z_{\text{eff}}$  inc. by  $\sim 1$  as each atom has added 1 proton to nucleus and 1  $e^-$  to valence shell (which does not screen)



$Z_{\text{eff}}$  inc.

### 2) Down Column

$Z_{\text{eff}}$  inc. slightly as valence shell  $e^-$  can penetrate better



### III) Atomic and Ionic Radii

#### A) Atomic Radii

##### 1) Nonbonding

Closest approach of atoms based on gas phase collisions or crystal structures

##### 2) Bonding Atomic Radius


Bond Length:

Distance between atoms in a covalently bound compd., averaged over many compds.

b.a.r =  $\frac{1}{2}$  bond length

Electron  
distribution  
in molecule

Nonbonding  
atomic radius



Copyright © 2006 Pearson Prentice Hall, Inc.

### 3) Size inc. down a group


$e^-$  occupy a higher energy level w. each element down a group &  $n$  determines size of orbital and avg. radius

$\therefore$  Inc.  $n \Rightarrow$  Inc. atomic size

### 4) Size generally dec. across a period from left to right

- $e^-$  added to same shell
- nuclear charge,  $Z_{\text{eff}}$ , inc. which pulls whole shell closer


### 5) Overall Trend



## B) Ionic Radii

Determined from crystal structure of ionic cmpds.

Averaged interatomic distance from multiple cmpds.




## 1) Cations

always smaller than parent atom

$\text{Cs}^+$  smaller than Cs

a) Size decreases with increasing ionic charge



## 2) Anions

always larger than parent atom











































## 3) Isoelectronic Series

Same #  $e^-$



$\xrightarrow{\hspace{10em}}$   
 $Z_{\text{eff}}$  inc., radius dec.

| Group 1A                                                                                                                                                                                              | Group 2A                                                                                                                                                                                                 | Group 3A                                                                                                                                                                                                 | Group 6A                                                                                                                                                                                                   | Group 7A                                                                                                                                                                                                 |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $\text{Li}^+$ 0.90 <br> Li 1.34     | $\text{Be}^{2+}$ 0.59 <br> Be 0.90     | $\text{B}^{3+}$ 0.41 <br> B 0.82       |  $\text{O}^{2-}$ 1.26<br> O 0.73      |  $\text{F}^-$ 1.19<br> F 0.71      |
|  $\text{Na}^+$ 1.16<br> Na 1.54     | $\text{Mg}^{2+}$ 0.86 <br> Mg 1.30     | $\text{Al}^{3+}$ 0.68 <br> Al 1.18     |  $\text{S}^{2-}$ 1.70<br> S 1.02       |  $\text{Cl}^-$ 1.67<br> Cl 0.99    |
|  $\text{K}^+$ 1.52<br> K 1.96    | $\text{Ca}^{2+}$ 1.14 <br> Ca 1.24  | $\text{Ga}^{3+}$ 0.76 <br> Ga 1.26  |  $\text{Se}^{2-}$ 1.84<br> Se 1.16  |  $\text{Br}^-$ 1.82<br> Br 1.14 |
|  $\text{Rb}^+$ 1.66<br> Rb 2.11 |  $\text{Sr}^{2+}$ 1.32<br> Sr 1.92 | $\text{In}^{3+}$ 0.94 <br> In 1.44 |  $\text{Te}^{2-}$ 2.07<br> Te 1.35 |  $\text{I}^-$ 2.06<br> I 1.33  |



= cation



= anion



= neutral atom

## IV) Ionization Energy, I.E.

Ionization: removal of an  $e^-$

I.E. : energy required to remove  $e^-$   
from gaseous atom or ion



$e^-$  removed is from highest  
energy level (highest  $n$  &  $l$ )

I.E. depends on avg. distance  
from the nucleus.

First I.E.,  $I_1$

Energy req. to remove the highest energy  $e^-$  from neutral atom



Second I.E  $I_2$

Energy req. to remove the next highest energy  $e^-$  from ion



Successive I.E. inc. in magnitude

- #  $e^-$  dec. (less repulsion)
- $Z$  (#  $p^+$ ) same (greater attraction)



|    | $I_1$               | $I_2$          | $I_3$          | $I_4$        | $I_5$  |
|----|---------------------|----------------|----------------|--------------|--------|
| Na | 496<br>[Ne]         | 4560           |                |              |        |
| Mg | 738<br>$3s^1$       | 1450<br>[Ne]   | 7730           |              |        |
| Al | 577<br>$3s^2$       | 1816<br>$3s^1$ | 2744<br>[Ne]   | 11,600       |        |
| Si | 786<br>$3s^2 3p^1$  | 1577<br>$3s^2$ | 3228<br>$3s^1$ | 4354<br>[Ne] | 16,100 |
| P  | 1060<br>$3s^2 3p^2$ | 1890           | 2905           | 4950         | 6270   |
| S  | 999<br>$3s^2 3p^3$  | 2260           | 3375           | 4565         | 6950   |

inner-shell  $e^-$

I.E. for removing  $e^-$  beyond valence  $e^-$   
greater than energy involved in  
chem. rxns & bonding

- only  $e^-$  outside noble-gas core  
involved in chem. change

Remember:

Atoms tend to lose or gain  $e^-$   
to get filled outer shell

- $e^-$  config. of a noble gas

Note

I.E. depends on avg. distance  
from nucleus &  $Z_{\text{eff}}$

$$\text{I.E.} \propto Z_{\text{eff}}$$

$$\text{I.E.} \propto 1/r$$

## A) Up a Group

Dec. atomic radius

$e^-$  held more tightly

I.E. Inc.

## B) Across a Period


Dec. atomic radius

$Z_{\text{eff}}$  inc.  $e^-$  held more tightly

I.E. Inc

## C) Summary





Copyright © 2006 Pearson Prentice Hall, Inc.

## D) Irregularities

$e^-$  config. accounts for irregularities

Li  $\rightarrow$  Ne, generally inc.

However,



$1/2$ -filled & filled subshells more stable

Elements at end of each transition series, Zn, Cd & Hg have higher I.E. than following element

pseudo-noble-gas

Highest I.E. for noble gases  
- filled s & p subshells

## E) Electron Config. of Ions

### 1) Representative Ions


#### a) Metals

Form Cations

#### 1) s - block

Groups 1A & 2A

All valence  $e^-$  removed  
- noble-gas config.



# 1) p - block

Groups 3A - 5A

Lose p e<sup>-</sup> fairly readily

(group # - 2)

Often req. too much energy  
to remove all val. e<sup>-</sup>

(group #)



$\text{Pb}^{2+}$  more common than  $\text{Pb}^{4+}$

## b) NonMetals

Monatomic anions

$$\text{charge} = (\text{group \#} - 8)$$

- add  $e^-$  to obtain  
noble-gas  $e^-$  config.





## 2) Transition Metal Ions

Generally, only highest energy  $e^-$  lost

Outer s-subshell  $e^-$

Many tran. metals form +2 cations

- lose both s-subshell  $e^-$

For ions of higher charge  
d-subshell  $e^-$  are lost

a) Ex 1:

Group 2B

Zn, Cd, Hg



b) Ex 2:



## V) Electron Affinity, EA

Energy associated with the gain of an  $e^-$  by a gaseous atom or ion

### A) First EA



Energy released for most neutral atoms & all positive ions

greater attraction for  $e^- \Rightarrow$  more neg. EA

### B) Second EA



$2^{\text{nd}}$   $e^-$  must be forced onto a neg. charged ion which requires energy

|                  |                  |                  |                   |                   |                   |                   |                  |
|------------------|------------------|------------------|-------------------|-------------------|-------------------|-------------------|------------------|
| <b>H</b><br>-73  |                  |                  |                   |                   |                   |                   | <b>He</b><br>> 0 |
| <b>Li</b><br>-60 | <b>Be</b><br>> 0 | <b>B</b><br>-27  | <b>C</b><br>-122  | <b>N</b><br>> 0   | <b>O</b><br>-141  | <b>F</b><br>-328  | <b>Ne</b><br>> 0 |
| <b>Na</b><br>-53 | <b>Mg</b><br>> 0 | <b>Al</b><br>-43 | <b>Si</b><br>-134 | <b>P</b><br>-72   | <b>S</b><br>-200  | <b>Cl</b><br>-349 | <b>Ar</b><br>> 0 |
| <b>K</b><br>-48  | <b>Ca</b><br>-2  | <b>Ga</b><br>-30 | <b>Ge</b><br>-119 | <b>As</b><br>-78  | <b>Se</b><br>-195 | <b>Br</b><br>-325 | <b>Kr</b><br>> 0 |
| <b>Rb</b><br>-47 | <b>Sr</b><br>-5  | <b>In</b><br>-30 | <b>Sn</b><br>-107 | <b>Sb</b><br>-103 | <b>Te</b><br>-190 | <b>I</b><br>-295  | <b>Xe</b><br>> 0 |
| 1A               | 2A               | 3A               | 4A                | 5A                | 6A                | 7A                | 8A               |

Copyright © 2006 Pearson Prentice Hall, Inc.

## A) Periodic Trends in EA

Generally, parallels variation  
in atomic size

- not as well-established  
as other trends (exceptions)

$e^-$  placed into outer shell

- closer it gets to nucleus  
& greater  $Z_{\text{eff}}$

larger neg. EA

## C) Summary



# 1) Exceptions

## a) 2<sup>nd</sup> period

F: -328 kJ/mol      Cl: -349 kJ/mol

True for other 2<sup>nd</sup> period elements

Small size of 2<sup>nd</sup> period elements

$e^-$  enters small outer shell

Adding an  $e^-$  places it very close to other 2s and 2p  $e^-$  resulting in stronger  $e^- - e^-$  repulsions.

## b) Other Exceptions

Adding an  $e^-$  to stable  $e^-$  - config.

### 1) Group 2A

full s subshell

added  $e^-$  goes into p subshell

### 2) Group 5A

$\frac{1}{2}$  - filled p valence subshell


added  $e^-$  pairs w. another  $e^-$  in occupied p orbital & experiences repulsions

### 3) Group 8A

filled valence shell


$e^-$  goes into next higher shell

# VI) Metals, Nonmetals, Metalloids







# Summary of Periodic Trends




increasing  
metallic  
character



increasing  
atomic  
radius



increasing  
ionization  
energy



increasing  
electron  
affinity